

# Improved error estimates of the splitting methods for the long time dynamics for the weakly nonlinear Schrödinger equations

Yongyong Cai

Beijing Normal University

joint with Weizhu Bao and Yue Feng

Guiyang, Guizhou, 08/08/2023

- 1 Weakly nonlinear/perturbed Schrödinger equations
- 2 Splitting methods and main results
- 3 Improved estimates-linear case
- 4 Improved estimates-nonlinear case

# Weakly nonlinear/perturbed Schrödinger equation

- Schrödinger equation (SE) with small potential on torus

$$i\partial_t\psi(\mathbf{x}, t) = -\Delta\psi(\mathbf{x}, t) + \varepsilon V(\mathbf{x})\psi(\mathbf{x}, t)$$

- Weakly nonlinear Schrödinger equation (NLSE) on torus

$$i\partial_t\psi(\mathbf{x}, t) = -\Delta\psi(\mathbf{x}, t) \pm \varepsilon^2|\psi(\mathbf{x}, t)|^2\psi(\mathbf{x}, t)$$

- $0 < \varepsilon \ll 1$  a small parameter
- Initial value  $\psi(\mathbf{x}, 0) = O(1)$

- Hamiltonian perturbation of linear PDEs on torus (J. Bourgain, Ann. Math., 98')

$$iu_t + Au + \varepsilon \partial_{\bar{u}} H(u, \bar{u}) = 0$$

-2D linear Schrödinger equation: quasi-periodic solutions

- 2D weakly nonlinear Schrödinger equation on torus (E. Faou, P. Germain, Z. Hani, JAMS, 16')

$$iu_t = -\Delta u \pm |u|^2 u$$

– global solution with small initial data

– small data  $u(0) = O(\varepsilon)$ ,  $u = \varepsilon \psi$ ,  $\psi(0) = O(1)$

$$i\partial_t \psi = -\Delta \psi \pm \varepsilon^2 |\psi|^2 \psi$$

# Property as $\varepsilon \rightarrow 0$

- SE with small potential:  $i\partial_t\psi = -\Delta\psi + \varepsilon V\psi$

trivial —  $[0, T/\varepsilon^\alpha]$  ( $\alpha \in [0, 1)$ )

non-trivial behavior happens  $\alpha = 1$

$$t \rightarrow t\varepsilon: i\partial_t\psi = -\frac{\Delta}{\varepsilon}\psi + V\psi$$

- Weakly NLSE  $i\partial_t\psi = -\Delta\psi \pm \varepsilon^2|\psi|^2\psi$

trivial linear dynamics  $[0, T/\varepsilon^\alpha]$  ( $\alpha \in [0, 2)$ )

non-trivial dynamics for  $\alpha = 2$

$$t \rightarrow t\varepsilon^2: i\partial_t\psi = -\frac{\Delta}{\varepsilon^2}\psi \pm |\psi|^2\psi$$

# Limit as $\varepsilon \rightarrow 0$

- Nonlinear case:  $i\partial_t \psi = -\frac{\Delta}{\varepsilon^2} \psi \pm |\psi|^2 \psi$
- $\tilde{\psi} = e^{-it\frac{\Delta}{\varepsilon^2}} \psi$ :

$$i\partial_t \tilde{\psi} = \pm F(t/\varepsilon^2, \tilde{\psi}), \quad F(s, \tilde{\psi}) = e^{-is\Delta} \left( \left| e^{is\Delta} \tilde{\psi} \right|^2 e^{is\Delta} \tilde{\psi} \right)$$

RHS:  $F(s, \tilde{\psi})$  periodic in  $s$

- in 1D torus
- 2D  $[0, L_x] \times [0, L_y]$  with  $(L_y/L_x)^2$  rational
- 3D  $[0, L_x] \times [0, L_y] \times [0, L_z]$  with  $(L_y/L_x)^2, (L_z/L_x)^2$  rational
- 1D  $[0, L]$ ,  $\tilde{\psi} = \sum_{k \in \mathbb{Z}} a_k e^{2\pi i k x / L}$

$$F(s, \tilde{\psi}) = \pm \sum_k \left( \sum_{\mathcal{I}_k} e^{-4\pi^2 s i \delta / L^2} a_{k_1} \bar{a}_{k_2} a_{k_3} \right) e^{2\pi i k x / L}$$

$$\mathcal{I}_k = \{(k_1, k_2, k_3) \in \mathbb{Z}^3 : k_1 - k_2 + k_3 = k\}, \quad \delta = k^2 - k_1^2 + k_2^2 - k_3^2$$

- Averaging as  $\varepsilon \rightarrow 0$ : formally, only  $\delta = 0$  term in the limit  $\varepsilon \rightarrow 0$
- averaging based methods (P. Chartier, A. Murua, J.M. Sanz-Serna, FOCM, 10', 12')

- Linear  $i\partial_t\psi = -\Delta\psi + \varepsilon V\psi$  over interval  $[0, T/\varepsilon]$
- Nonlinear  $i\partial_t\psi = -\Delta\psi \pm \varepsilon^2|\psi|^2\psi$  over interval  $[0, T/\varepsilon^2]$
- Sufficiently regular initial data  $\rightarrow$  uniform Sobolev norms
- Numerical methods (splitting methods) and performance?
  - splitting methods known for good performance over long time

# Splitting methods for linear case

- 1D linear on periodic  $[a, b]$ :  $i\partial_t\psi(x, t) = -\Delta\psi(x, t) + \varepsilon V(x)\psi(x, t)$
- split into two subproblems:
  - $i\partial_t\psi(x, t) = -\Delta\psi(x, t)$  solve exactly in phase space  
 $\psi(\cdot, t) = e^{it\Delta}\psi_0(\cdot)$
  - $i\partial_t\psi(x, t) = \varepsilon V(x)\psi(x, t)$  integrate exactly in physical space  
 $\psi(x, t) = e^{-i\varepsilon tV(x)}\psi_0(x)$
- Strang splitting  $\psi^{[n+1]}(x) = \mathcal{S}_\tau(\psi^{[n]}) = e^{i\frac{\tau}{2}\Delta}e^{-i\varepsilon\tau V(x)}e^{i\frac{\tau}{2}\Delta}\psi^{[n]}(x)$
- Strang splitting with Fourier pseudo-spectral discretization:  
 $\psi_j^n \approx \psi^{[n+1]}(x_j) \quad (x_j = a + jh, \quad h = (b - a)/N)$ 

$$\psi_j^{(1)} = \sum_{l=-N/2}^{N/2-1} e^{-i\frac{\tau\mu_l^2}{2}} (\widetilde{\psi^n})_l e^{i\mu_l(x_j-a)}, \quad \psi_j^{(2)} = e^{-i\varepsilon\tau V(x_j)}\psi_j^{(1)}$$

$$\psi_j^{n+1} = \sum_{l=-N/2}^{N/2-1} e^{-i\frac{\tau\mu_l^2}{2}} (\widetilde{\psi^{(2)}})_l e^{i\mu_l(x_j-a)}$$

$$(\widetilde{\psi^n})_l \text{ the discrete Fourier transform of } \psi_j^n$$



# Splitting methods for nonlinear case

- 1D nonlinear on periodic  $[a, b]$ :  $i\partial_t\psi(x, t) = -\Delta\psi \pm \varepsilon^2|\psi|^2\psi$
- Strang splitting  $\psi^{[n+1]}(x) = \mathcal{S}_\tau(\psi^{[n]}) = e^{i\frac{\tau}{2}\Delta} e^{\mp i\varepsilon^2\tau} \left| e^{i\frac{\tau}{2}\Delta} \psi^{[n]}(x) \right|^2 e^{i\frac{\tau}{2}\Delta} \psi^{[n]}(x)$
- Strang splitting with Fourier pseudo-spectral discretization:  
 $\psi_j^n \approx \psi^{[n+1]}(x_j) \quad (x_j = a + jh, \quad h = (b - a)/N)$

$$\psi_j^{(1)} = \sum_{l=-N/2}^{N/2-1} e^{-i\frac{\tau\mu_l^2}{2}} \widetilde{(\psi^n)}_l e^{i\mu_l(x_j-a)}$$

$$\psi_j^{(2)} = e^{\mp i\varepsilon^2\tau\lambda|\psi_j^{(1)}|^2} \psi_j^{(1)}$$

$$\psi_j^{n+1} = \sum_{l=-N/2}^{N/2-1} e^{-i\frac{\tau\mu_l^2}{2}} \widetilde{(\psi^{(2)})}_l e^{i\mu_l(x_j-a)}$$

# Some known properties of Splitting

- $\varepsilon = 1$

Convergence in the linear case  $O(\tau^2)$  (T. Jahnke, C. Lubich, BIT, 00'; S. Blanes, F. Casas, A. Murua, 14', ...) )

Convergence in the nonlinear case  $O(\tau^2)$  (C. Lubich, MCOM, 08', ...) )

Convergence in the nonlinear fully discrete case (Z. Wang, J. Shen, FOCM, 13'; W. Bao, Y. Cai, 13', ...) )

Long time near conservation (L. Gauckler, C. Lubich, FOCM, 10', ...) )

Convergence in low regularity/filtered case: (A. Ostermann, F. Rousset, K. Schratz, 22', ...) )

- semi-classical case

Best resolution in observables, etc. (W. Bao, S. Jin, P.A. Markowich, 02', ...) )

General splitting linear setup (P. Bader, A. Iserles, K. Kropielnicka, P. Singh, FOCM, 14', ...) )

- what to be expected when  $0 < \varepsilon \ll 1$ ?

# Direct computation-linear case

- Local error:  $\mathcal{S}_\tau(\psi^n) = e^{i\frac{\tau}{2}\Delta} e^{-i\varepsilon\tau V(x)} e^{i\frac{\tau}{2}\Delta} \psi^n$

$$\begin{aligned}\mathcal{S}_\tau(\psi(t_n)) &= e^{i\tau\Delta} \psi(t_n) - i\varepsilon\tau \left( e^{i\frac{\tau}{2}\Delta} V e^{i\frac{\tau}{2}\Delta} \psi(t_n) \right) \\ &\quad - \frac{\varepsilon^2\tau^2}{2} e^{i\frac{\tau}{2}\Delta} V^2 e^{i\frac{\tau}{2}\Delta} \psi(t_n) + O(\varepsilon^3\tau^3), \\ \psi(t_{n+1}) &= e^{i\tau\Delta} \psi(t_n) - i\varepsilon \int_0^\tau \left( e^{i(\tau-s)\Delta} V \psi(t_n + s) \right) ds \\ &= e^{i\tau\Delta} \psi(t_n) - i\varepsilon \int_0^\tau \left( e^{i(\tau-s)\Delta} V e^{is\Delta} \psi(t_n) \right) ds \\ &\quad - \varepsilon^2 \int_0^\tau \int_0^s \left( e^{i(\tau-s)\Delta} V e^{i(s-w)\Delta} V e^{iw\Delta} \psi(t_n) \right) dw ds + O(\varepsilon^3\tau^3)\end{aligned}$$

- $\mathcal{S}_\tau(\psi(t_n)) - \psi(t_{n+1}) = O(\varepsilon\tau^3[\Delta, [\Delta, V]]) + O(\varepsilon^2\tau^3) + O(\varepsilon^3\tau^3)$
- stability  $\|\mathcal{S}_\tau\psi\|_{H^1} \lesssim (1 + \varepsilon\tau)\|\psi\|_{H^1}$ ,  $\|\mathcal{S}_\tau\psi\|_{L^2} = \|\psi\|_{L^2}$
- over  $[0, T/\varepsilon]$ , global error  $\frac{T}{\varepsilon\tau} O(\varepsilon\tau^3[\Delta, [\Delta, V]]) = O(\tau^2)$

# Extension to weakly NLSE

- Local error

$$e^{i\frac{\tau}{2}\Delta} e^{\mp i\varepsilon^2\tau} \left| e^{i\frac{\tau}{2}\Delta} \psi(t_n) \right|^2 e^{i\frac{\tau}{2}\Delta} \psi(t_n) - \psi(t_{n+1}) = -i\varepsilon^2\tau f\left(\frac{\tau}{2}\right) + i\varepsilon^2 \int_0^\tau f(s) ds + O(\varepsilon^4\tau^3)$$

- $f(s) = \pm e^{i(\tau-s)\Delta} |e^{is\Delta}\psi(t_n)|^2 e^{is\Delta}\psi(t_n)$
- size:  $O(\varepsilon^2\tau^3[\Delta, [\Delta, |\psi|^2]]) + O(\varepsilon^4\tau^3) = O(\varepsilon^2\tau^3)$
- Direct extension would be  $O(\tau^2) = O(\varepsilon^2\tau^2) \frac{T}{\tau\varepsilon^2}$  over  $[0, T/\varepsilon^2]$

# Known results in the nonlinear case

- **Improved** error in weakly NLSE  $0 < \varepsilon \ll 1$  (P. Chartier, F. Méhats, M. Thalhammer, Y. Zhang, MCOM, 16')
- $e^{it\Delta}$  periodic with period  $\Delta T$  ( $L^2/4\pi^2$ , 1D),  $\tau = \frac{\Delta T}{N}$
- Strang splitting error improved!

$$\|\psi^{[n]} - \psi(t_n)\| \lesssim \varepsilon^2 \tau^2 + N^{-m}$$

$m$ -regularity dependent parameter

- A factor of  $\varepsilon^2$  improved!
- Drawback: periodicity highly involved

# Main results

For weakly NLSE, Strang splitting with Fourier pseudo-spectral

## Theorem

Let  $\psi^n$  be the numerical approximation obtained from the TSFP. For sufficiently regular data, for any  $0 < \varepsilon \leq 1$ , when  $0 < \tau \leq \alpha \frac{\tau_0^2(b-a)^2}{4\pi(1+\tau_0)^2} < 1$  with a constant  $\alpha \in (0, 1)$ , there holds

$$\|\psi(x, t_n) - I_N \psi^n\|_{H^1} \lesssim h^{m-1} + \varepsilon^2 \tau^2 + \tau_0^{m-1},$$

$$\|I_N \psi^n\|_{H^1} \leq 1 + M, \quad 0 \leq n \leq \frac{T/\varepsilon^2}{\tau},$$

$M := \|\psi\|_{L^\infty([0, T_\varepsilon]; H^1)}$ . If the exact solution is smooth, i.e.  $\psi(x, t) \in H_{\text{per}}^\infty$ , the  $\tau_0^{m-1}$  can be ignored in practical computation when  $\tau_0$  is small but fixed, the estimates become

$$\|\psi(x, t_n) - I_N \psi^n\|_{H^1} \lesssim h^{m-1} + \varepsilon^2 \tau^2.$$

# Results for linear case

TSFP for SE with small potential

## Theorem

For  $\varepsilon \in (0, 1]$  and a fixed  $\tau_0 \in (0, 1)$ , for  $\tau \in \left(0, \alpha \frac{(b-a)^2}{2\pi(1+\tau_0)^2} \tau_0^2\right)$  ( $\alpha \in (0, 1)$ ) and sufficiently regular data, the error reads

$$\|\psi(x, t_n) - I_N \psi^n\|_{H^1} \lesssim h^{m-1} + \varepsilon \tau^2 + \tau_0^{m-1}, \quad 0 \leq n \leq \frac{T/\varepsilon}{\tau}.$$

Again for smooth solution, i.e.  $\psi(x, t) \in H_{\text{per}}^\infty$ , the  $\tau_0^{m-1}$  part can be ignored in practical computation, the improved error bounds read

$$\|\psi(x, t_n) - I_N \psi^n\|_{H^1} \lesssim h^{m-1} + \varepsilon \tau^2, \quad 0 \leq n \leq \frac{T/\varepsilon}{\tau}.$$

# Estimates revisited-linear case

- Local error-full discretization:  $\mathcal{S}_\tau : \psi \rightarrow e^{i\frac{\tau}{2}\Delta} e^{-i\varepsilon\tau V(x)} \psi$  (propagator of semi discretization)

$$\mathcal{E}^n(x) := P_N \mathcal{S}_\tau(P_N \psi(t_n)) - P_N \psi(t_{n+1}) = P_N \mathcal{F}(P_N \psi(t_n)) + R_n$$

$P_N$  projection onto Fourier modes  $k = -N/2, \dots, N/2 - 1$



$$\mathcal{F}(P_N \psi(t_n)) = -i\varepsilon\tau f^n\left(\frac{\tau}{2}\right) + i\varepsilon \int_0^\tau f^n(s) ds$$

with  $f^n(s) = e^{i(\tau-s)\Delta} V e^{is\Delta} P_N \psi(t_n)$

- $\|R_n\|_{H^1} \lesssim \varepsilon^2 \tau^3 + \varepsilon \tau h^{m-1}$ ,  $\|\mathcal{F}(P_N \psi(t_n))\|_{H^1} \lesssim \varepsilon \tau^3$
- Naive estimates over  $0 \leq n \leq \frac{T}{\tau\varepsilon}$  interval gives  $h^{m-1} + \varepsilon \tau^2 + \tau^2$
- Improved error?



# Estimates of global error

## Global error

- pseudo-spectral v.s. spectral (local error)

$$I_N \psi^{n+1} = e^{i\frac{\tau}{2}\Delta} (I_N \psi^{(2)}), \quad I_N(\psi^{(2)}) = I_N(e^{-i\varepsilon\tau V(x)} \psi^{(1)}), \quad I_N \psi^{(1)} = e^{i\frac{\tau}{2}\Delta} I_N \psi^n$$

$$P_N(\mathcal{S}_\tau(\psi(t_n))) = e^{i\frac{\tau}{2}\Delta} (P_N \psi^{(2)}), \quad \psi^{(2)} = e^{-i\varepsilon\tau V(x)} \psi^{(1)}, \quad \psi^{(1)} = e^{i\frac{\tau}{2}\Delta} P_N \psi(t_n)$$

- error:  $e^n := e^n(x) = I_N \psi^n - P_N \psi(t_n)$ :

$$e^{n+1} = e^{i\tau\Delta} e^n + Q^n(x) + \mathcal{E}^n$$

## $\mathcal{E}^n$ -local error

$$Q^n(x) = -i\varepsilon\tau e^{i\frac{\tau}{2}\Delta} \left( I_N(V(x)) \int_0^1 e^{-i\varepsilon\theta\tau V(x)} d\theta \psi^{(1)} \right) + i\varepsilon\tau e^{i\frac{\tau}{2}\Delta} \left( P_N(V(x)) \int_0^1 e^{-i\varepsilon\theta\tau V(x)} d\theta \psi^{(1)} \right)$$

- $\|Q^n(x)\|_{H^1} \lesssim \varepsilon\tau (h^{m-1} + \|e^n\|_{H^1})$

$$e^{n+1} = e^{i(n+1)\tau\Delta} e^0 + \sum_{k=0}^n e^{i(n-k)\tau\Delta} (Q^k(x) + \mathcal{E}^k)$$

- Global error

$$\|e^{n+1}\|_{H^1} \lesssim h^{m-1} + \varepsilon\tau^2 + \varepsilon\tau \sum_{k=0}^n \|e^k\|_{H^1} + \left\| \sum_{k=0}^n e^{i(n-k)\tau\Delta} P_N \mathcal{F}(P_N \psi(t_k)) \right\|_{H^1}$$

# Refined error analysis

- Refined estimates on  $\sum_{k=0}^n e^{i(n-k)\tau\Delta} P_N \mathcal{F}(P_N \psi(t_k))$

$$\mathcal{F}(P_N \psi(t_n)) = -i\varepsilon\tau f^n\left(\frac{\tau}{2}\right) + i\varepsilon \int_0^\tau f^n(s) ds, \quad f^n(s) = e^{i(\tau-s)\Delta} V e^{is\Delta} P_N \psi(t_n)$$

- Periodicity approach:  $\tau = L^2/4\pi^2/N^*$  ( $N^* \in \mathbb{Z}^+$ )

sum  $k = 0, \dots, N^* - 1$  is equivalent to a midpoint rule for a periodic function—spectral in time  $(N^*)^{-m}!$

$O(\varepsilon\tau^3)$  only accumulate at most in one period to  $O(\varepsilon\tau^2)$ —desired!

- Our approach: (regularity compensate oscillation)
  - Separate high frequency and low frequency:  $> 1/\tau_0$  and  $\leq 1/\tau_0$  ( $\tau_0$  parameter)
  - High modes controlled by regularity (projection):  $N_0 = 2\lceil 1/\tau_0 \rceil \in \mathbb{Z}^+$

$$\left\| P_{N_0} \mathcal{F}(P_{N_0} \psi(t_n)) - P_N \mathcal{F}(P_N \psi(t_n)) \right\|_{H^1} \lesssim \varepsilon\tau(h^{m-1} + \tau_0^{m-1})$$

- Calculate phase cancellation for low modes  $\leq 1/\tau_0$  (summation by parts), i.e. the phase in  $f^n(s)$

## RCO for phase cancellation

- gain order in  $\varepsilon$ :  $\partial_t \psi(x, t) - i\Delta \psi(x, t) = O(\varepsilon)$ , 'twisted variable' as

$$\phi(x, t) = e^{-it\Delta} \psi(x, t), \quad \partial_t \phi(x, t) = O(\varepsilon)$$

- low modes left:  $\|e^{n+1}\|_{H^1} \lesssim h^{m-1} + \tau_0^{m-1} + \varepsilon \tau^2 + \varepsilon \tau \sum_{k=0}^n \|e^k\|_{H^1} + \|\mathcal{R}^n\|_{H^1}$

$$\mathcal{R}^n(x) = \sum_{k=0}^n e^{-i(k+1)\tau\Delta} P_{N_0} \mathcal{F}(e^{it_k\Delta}(P_{N_0}\phi(t_k)))$$

- $\mathcal{T}_{N_0} = \{-N_0/2, -N_0/2+1, \dots, N_0/2-1\}$ ,  $\mathcal{I}_l^{N_0} = \{(l_1, l_2) \mid l_1 + l_2 = l, l_1 \in \mathbb{Z}, l_2 \in \mathcal{T}_{N_0}\}$

- $\phi(t) = \sum_{l \in \mathbb{Z}} \widehat{\phi}_l(t) e^{i\mu_l(x-a)} \quad (t \geq 0)$ ,  $P_{N_0}\phi(t) = \sum_{l \in \mathcal{T}_{N_0}} \widehat{\phi}_l(t) e^{i\mu_l(x-a)}$

- $\mathcal{R}^n(x) = i\varepsilon \sum_{k=0}^n \sum_{l \in \mathcal{T}_{N_0}} \sum_{(l_1, l_2) \in \mathcal{I}_l^{N_0}} \lambda_{k,l,l_1,l_2} e^{i\mu_l(x-a)}$

- $\lambda_{k,l,l_1,l_2} = -\tau \mathcal{G}_{k,l,l_1,l_2}(\tau/2) + \int_0^\tau \mathcal{G}_{k,l,l_1,l_2}(s) ds = r_{l,l_2} e^{it_k \delta_{l,l_2}} c_{k,l,l_1,l_2}$

- $\mathcal{G}_{k,l,l_1,l_2}(s) = e^{i(t_k+s)\delta_{l,l_2}} \widehat{V}_{l_1} \widehat{\phi}_{l_2}(t_k)$ ,  $\delta_{l,l_2} = \delta_l - \delta_{l_2}$ ,  $\delta_l = \mu_l^2$

$$r_{l,l_2} = -\tau e^{i\frac{\tau \delta_{l,l_2}}{2}} + \int_0^\tau e^{is\delta_{l,l_2}} ds = O(\tau^3(\delta_{l,l_2})^2)$$

## RCO

- summation by parts in time: exponential sums  $S_{n,l,l_2} = \sum_{k=0}^n e^{it_k \delta_{l,l_2}}$
- For  $0 < \tau \leq \alpha \frac{2\pi}{\mu_1^2(1+\tau_0)^2} \tau_0^2$ ,  $|S_{n,l,l_2}| \leq C/\tau |\delta_{l,l_2}|$
- $\sum_{k=0}^n \lambda_{k,l,l_1,l_2} = r_{l,l_2} \sum_{k=0}^{n-1} S_{k,l,l_2} (c_{k,l,l_1,l_2} - c_{k+1,l,l_1,l_2}) + S_{n,l,l_2} r_{l,l_2} c_{n,l,l_1,l_2}$
- $r_{l,l_2} = O(\tau^3 (\delta_{l,l_2})^2)$ ,  $c_{k,l,l_1,l_2} - c_{k+1,l,l_1,l_2} = \widehat{V}_{l_1} (\widehat{\phi}_{l_2}(t_k) - \widehat{\phi}_{l_2}(t_{k+1})) = O(\varepsilon \tau)$
- $|\sum_{k=0}^n \lambda_{k,l,l_1,l_2}| \lesssim \tau^2 |\delta_{l,l_2}| \left| \widehat{V}_{l_1} \right| \left[ \sum_{k=0}^{n-1} |\widehat{\phi}_{l_2}(t_k) - \widehat{\phi}_{l_2}(t_{k+1})| + |\widehat{\phi}_{l_2}(t_n)| \right]$
- $\|\mathcal{R}^n(x)\|_{H^1} \lesssim \varepsilon \tau^2$  for  $n\tau \leq T/\varepsilon$  (**Improved estimates by Gronwall inequality**)
- $\|\mathcal{R}^n(x)\|_{H^1}^2 = \varepsilon^2 \sum_{l \in \mathcal{T}_{N_0}} (1 + \mu_l^2) \left| \sum_{(l_1, l_2) \in \mathcal{I}_l^{N_0}} \sum_{k=0}^n \lambda_{k,l,l_1,l_2} \right|^2$
- Estimates sums instead by considering  $U(x) = \sum_{l \in \mathbb{Z}} (1 + \mu_l^2)^{3/2} \left| \widehat{V}_l \right| e^{i\mu_l(x-a)}$   
and  $\xi(x) = \sum_{l \in \mathbb{Z}} (1 + \mu_l^2)^{3/2} \left| \widehat{\phi}_l(t_n) \right| e^{i\mu_l(x-a)}$


- $\tau_0$ -artificial parameter (truncation of Fourier modes)-depend on the exact solution
- structure of Fourier functions  $e^{i\mu_l(x-a)}e^{i\mu_k(x-a)} = e^{i\mu_{l+k}(x-a)}$
- valid for non-periodic evolutionary Schrödinger operator
- more choices of time step size  $\tau$



- need  $|S_{n,l,l_2}| = \left| \sum_{k=0}^n e^{it_k \delta_{l,l_2}} \right| \lesssim \frac{1}{\tau |\delta_{l,l_2}|^\beta} \quad (\beta \leq 2)$



Diophantine condition ([Z. Shang, 00'](#))

$$\left| \frac{1 - e^{i\tau\mu_1^2 K}}{\tau} \right| \geq \frac{\gamma}{|K|^\nu}, \quad \forall K \in \mathbb{Z}, K \neq 0,$$

independent of  $\tau_0$ ,  $h$ , let  $\tau_0, h \rightarrow 0^+$ , recover  $\varepsilon\tau^2$  for the semi-discrete-in-time splitting error











# Extension to nonlinear case

- Diophantine type non-resonant time step size  $\tau$
- Higher dimensions with tensor grids—straightforward!
- Arbitrary aspect ratios in higher dimensions—no periodicity needed!
- Polynomial type nonlinearities—ok!
- extra larger step size ok—  $\tau = O(1/\varepsilon)$ -non-resonant for the nonlinear case  
 $\tau = O(1/\sqrt{\varepsilon})$ -non-resonant for the linear case
- Higher order splitting—ok but need work out



# Summary

- Improved error bounds of splitting methods for SE with small potential and weakly NLSE
- Resonance is important
- RCO approach efficient tool for analyzing Schrödinger type equations
- Larger time step size could be accurate

THANK YOU!