


## More linear classification

DSE 220

# The decision boundary




Decision boundary in  $\mathbb{R}^p$  is a **hyperplane**.

- How is this boundary parametrized?
- How can we learn a hyperplane from training data?

# Hyperplanes

Hyperplane  $\{x : w \cdot x = b\}$


- orientation  $w \in \mathbb{R}^p$
- offset  $b \in \mathbb{R}$



# Hyperplanes

Hyperplane  $\{x : w \cdot x = b\}$

- orientation  $w \in \mathbb{R}^p$
- offset  $b \in \mathbb{R}$



Can always normalize  $w$  to unit length:


$$(w, b) \longleftrightarrow \left( \hat{w} = \frac{w}{\|w\|}, \frac{b}{\|w\|} \right)$$

$$w \cdot x = b \longleftrightarrow \hat{w} \cdot x = \frac{b}{\|w\|}$$

# Hyperplanes

Hyperplane  $\{x : w \cdot x = b\}$

- orientation  $w \in \mathbb{R}^p$
- offset  $b \in \mathbb{R}$



Can always normalize  $w$  to unit length:

$$(w, b) \longleftrightarrow \left( \hat{w} = \frac{w}{\|w\|}, \frac{b}{\|w\|} \right)$$
$$w \cdot x = b \longleftrightarrow \hat{w} \cdot x = \frac{b}{\|w\|}$$

Equivalently: all points whose projection onto  $\hat{w}$  is  $b/\|w\|$ .

# Homogeneous linear separators

Hyperplanes that pass through the origin have no offset,  $b = 0$ .

# Homogeneous linear separators

Hyperplanes that pass through the origin have no offset,  $b = 0$ .

Reduce to this case by adding an extra feature to  $x$ :

$$\tilde{x} = (x, 1) \in \mathbb{R}^{p+1}$$

Then  $\{x : w \cdot x = b\} \equiv \{x : \tilde{w} \cdot \tilde{x} = 0\}$  where  $\tilde{w} = (w, -b)$ .



# Homogeneous linear separators

Hyperplanes that pass through the origin have no offset,  $b = 0$ .

Reduce to this case by adding an extra feature to  $x$ :

$$\tilde{x} = (x, 1) \in \mathbb{R}^{p+1}$$

Then  $\{x : w \cdot x = b\} \equiv \{x : \tilde{w} \cdot \tilde{x} = 0\}$  where  $\tilde{w} = (w, -b)$ .





# The learning problem: separable case

*Input:* training data  $(x^{(1)}, y^{(1)}), \dots, (x^{(n)}, y^{(n)}) \in \mathbb{R}^p \times \{-1, +1\}$

*Output:* linear classifier  $w \in \mathbb{R}^p$  such that

$$y^{(i)}(w \cdot x^{(i)}) > 0 \quad \text{for } i = 1, 2, \dots, n$$

# The learning problem: separable case

*Input:* training data  $(x^{(1)}, y^{(1)}), \dots, (x^{(n)}, y^{(n)}) \in \mathbb{R}^p \times \{-1, +1\}$

*Output:* linear classifier  $w \in \mathbb{R}^p$  such that

$$y^{(i)}(w \cdot x^{(i)}) > 0 \quad \text{for } i = 1, 2, \dots, n$$

This is linear programming:

- Each data point is a linear constraint on  $w$
- Want to find  $w$  that satisfies all these constraints

But we won't use generic linear programming methods, such as simplex.

# The learning problem: separable case

*Input:* training data  $(x^{(1)}, y^{(1)}), \dots, (x^{(n)}, y^{(n)}) \in \mathbb{R}^p \times \{-1, +1\}$

*Output:* linear classifier  $w \in \mathbb{R}^p$  such that

$$y^{(i)}(w \cdot x^{(i)}) > 0 \quad \text{for } i = 1, 2, \dots, n$$

This is linear programming:

- Each data point is a linear constraint on  $w$
- Want to find  $w$  that satisfies all these constraints

But we won't use generic linear programming methods, such as simplex.


A simple alternative: **Perceptron algorithm** (Rosenblatt, 1958)

- $w = 0$
- while some  $(x, y)$  is misclassified:
  - $w = w + yx$

# Perceptron: example

- $w = 0$
- while some  $(x, y)$  is misclassified:
  - $w = w + yx$

▲ +1  
● -1




**Separator:**  $w = 0$

# Perceptron: example

- $w = 0$
- while some  $(x, y)$  is misclassified:
  - $w = w + yx$

▲ +1  
● -1




**Separator:**  $w = -x^{(1)}$

# Perceptron: example

- $w = 0$
- while some  $(x, y)$  is misclassified:
  - $w = w + yx$

▲ +1  
● -1



**Separator:**  $w = -x^{(1)} + x^{(6)}$

## Perceptron: convergence

**Theorem:** Let  $R = \max \|x^{(i)}\|$ . Suppose there is a unit vector  $w^*$  and some (margin)  $\gamma > 0$  such that

$$y^{(i)}(w^* \cdot x^{(i)}) \geq \gamma \text{ for all } i.$$

Then the Perceptron algorithm converges after at most  $R^2/\gamma^2$  updates.


# Perceptron: convergence

**Theorem:** Let  $R = \max \|x^{(i)}\|$ . Suppose there is a unit vector  $w^*$  and some (margin)  $\gamma > 0$  such that

$$y^{(i)}(w^* \cdot x^{(i)}) \geq \gamma \text{ for all } i.$$

Then the Perceptron algorithm converges after at most  $R^2/\gamma^2$  updates.

**Proof idea.** Let  $w_t$  be the classifier after  $t$  updates.



**Track angle between  $w_t$  and  $w^*$ :**

$$\cos(\angle(w_t, w^*)) = \frac{w_t \cdot w^*}{\|w\|}.$$




# Perceptron: convergence

**Theorem:** Let  $R = \max \|x^{(i)}\|$ . Suppose there is a unit vector  $w^*$  and some (margin)  $\gamma > 0$  such that

$$y^{(i)}(w^* \cdot x^{(i)}) \geq \gamma \text{ for all } i.$$

Then the Perceptron algorithm converges after at most  $R^2/\gamma^2$  updates.

**Proof idea.** Let  $w_t$  be the classifier after  $t$  updates.



Track angle between  $w_t$  and  $w^*$ :

$$\cos(\angle(w_t, w^*)) = \frac{w_t \cdot w^*}{\|w\|}.$$

On each mistake, when  $w_t$  is updated to  $w_{t+1}$ ,

- $w_t \cdot w^*$  grows significantly.
- $\|w_t\|$  does not grow much.

## Perceptron convergence, cont'd

Perceptron update: if  $y(w_t \cdot x) < 0$  (misclassified) then  $w_{t+1} = w_t + yx$ .  
Target vector  $w^*$  has unit length, and margin condition  $y(w^* \cdot x) \geq \gamma$ .

## Perceptron convergence, cont'd

Perceptron update: if  $y(w_t \cdot x) < 0$  (misclassified) then  $w_{t+1} = w_t + yx$ .  
Target vector  $w^*$  has unit length, and margin condition  $y(w^* \cdot x) \geq \gamma$ .

- 1 Initial vector  $w_0 = 0$ .

## Perceptron convergence, cont'd

Perceptron update: if  $y(w_t \cdot x) < 0$  (misclassified) then  $w_{t+1} = w_t + yx$ .  
Target vector  $w^*$  has unit length, and margin condition  $y(w^* \cdot x) \geq \gamma$ .

- 1 Initial vector  $w_0 = 0$ .
- 2 When updating  $w_t$  to  $w_{t+1}$ :

$$w_{t+1} \cdot w^* = (w_t + yx) \cdot w^* = w_t \cdot w^* + y(w^* \cdot x) \geq w_t \cdot w^* + \gamma$$

## Perceptron convergence, cont'd

Perceptron update: if  $y(w_t \cdot x) < 0$  (misclassified) then  $w_{t+1} = w_t + yx$ .  
Target vector  $w^*$  has unit length, and margin condition  $y(w^* \cdot x) \geq \gamma$ .

- 1 Initial vector  $w_0 = 0$ .
- 2 When updating  $w_t$  to  $w_{t+1}$ :

$$\begin{aligned}w_{t+1} \cdot w^* &= (w_t + yx) \cdot w^* = w_t \cdot w^* + y(w^* \cdot x) \geq w_t \cdot w^* + \gamma \\ \|w_{t+1}\|^2 &= \|w_t + yx\|^2 = \|w_t\|^2 + \|x\|^2 + 2y(w_t \cdot x) \leq \|w_t\|^2 + R^2\end{aligned}$$

## Perceptron convergence, cont'd

Perceptron update: if  $y(w_t \cdot x) < 0$  (misclassified) then  $w_{t+1} = w_t + yx$ .  
Target vector  $w^*$  has unit length, and margin condition  $y(w^* \cdot x) \geq \gamma$ .

- 1 Initial vector  $w_0 = 0$ .
- 2 When updating  $w_t$  to  $w_{t+1}$ :

$$\begin{aligned}w_{t+1} \cdot w^* &= (w_t + yx) \cdot w^* = w_t \cdot w^* + y(w^* \cdot x) \geq w_t \cdot w^* + \gamma \\ \|w_{t+1}\|^2 &= \|w_t + yx\|^2 = \|w_t\|^2 + \|x\|^2 + 2y(w_t \cdot x) \leq \|w_t\|^2 + R^2\end{aligned}$$

- 3 After  $T$  updates, we have

$$\begin{aligned}w_T \cdot w^* &\geq T\gamma \\ \|w_T\|^2 &\leq TR^2\end{aligned}$$

## Perceptron convergence, cont'd

Perceptron update: if  $y(w_t \cdot x) < 0$  (misclassified) then  $w_{t+1} = w_t + yx$ .  
Target vector  $w^*$  has unit length, and margin condition  $y(w^* \cdot x) \geq \gamma$ .

- 1 Initial vector  $w_0 = 0$ .
- 2 When updating  $w_t$  to  $w_{t+1}$ :

$$\begin{aligned}w_{t+1} \cdot w^* &= (w_t + yx) \cdot w^* = w_t \cdot w^* + y(w^* \cdot x) \geq w_t \cdot w^* + \gamma \\ \|w_{t+1}\|^2 &= \|w_t + yx\|^2 = \|w_t\|^2 + \|x\|^2 + 2y(w_t \cdot x) \leq \|w_t\|^2 + R^2\end{aligned}$$

- 3 After  $T$  updates, we have

$$\begin{aligned}w_T \cdot w^* &\geq T\gamma \\ \|w_T\|^2 &\leq TR^2\end{aligned}$$


- 4 The angle between  $w_T$  and  $w^*$  is given by

$$\cos(\angle(w_T, w^*)) = \frac{w_T \cdot w^*}{\|w\|} \geq \frac{T\gamma}{R\sqrt{T}}.$$

This is at most 1, so  $T \leq R^2/\gamma^2$ .

## A better separator?


For a linearly separable data set, there are in general many possible separating hyperplanes, and Perceptron is guaranteed to find one of them.





## A better separator?

For a linearly separable data set, there are in general many possible separating hyperplanes, and Perceptron is guaranteed to find one of them.



But is there a better, more systematic choice of separator? The one with the most buffer around it, for instance?

## Maximizing the margin

Given training data  $(x^{(1)}, y^{(1)}), \dots, (x^{(n)}, y^{(n)}) \in \mathbb{R}^p \times \{-1, +1\}$ , find  $w \in \mathbb{R}^p$  and  $b \in \mathbb{R}$  such that  $y^{(i)}(w \cdot x^{(i)} + b) > 0$  for all  $i$ .

# Maximizing the margin

Given training data  $(x^{(1)}, y^{(1)}), \dots, (x^{(n)}, y^{(n)}) \in \mathbb{R}^p \times \{-1, +1\}$ , find  $w \in \mathbb{R}^p$  and  $b \in \mathbb{R}$  such that  $y^{(i)}(w \cdot x^{(i)} + b) > 0$  for all  $i$ .

By scaling  $w, b$ , can equally ask for


$$y^{(i)}(w \cdot x^{(i)} + b) \geq 1 \quad \text{for all } i.$$

# Maximizing the margin

Given training data  $(x^{(1)}, y^{(1)}), \dots, (x^{(n)}, y^{(n)}) \in \mathbb{R}^p \times \{-1, +1\}$ , find  $w \in \mathbb{R}^p$  and  $b \in \mathbb{R}$  such that  $y^{(i)}(w \cdot x^{(i)} + b) > 0$  for all  $i$ .

By scaling  $w, b$ , can equally ask for

$$y^{(i)}(w \cdot x^{(i)} + b) \geq 1 \quad \text{for all } i.$$




# Maximizing the margin

Given training data  $(x^{(1)}, y^{(1)}), \dots, (x^{(n)}, y^{(n)}) \in \mathbb{R}^p \times \{-1, +1\}$ , find  $w \in \mathbb{R}^p$  and  $b \in \mathbb{R}$  such that  $y^{(i)}(w \cdot x^{(i)} + b) > 0$  for all  $i$ .

By scaling  $w, b$ , can equally ask for


$$y^{(i)}(w \cdot x^{(i)} + b) \geq 1 \quad \text{for all } i.$$



Maximize the **margin**  $\gamma$ .


# What is the margin?

Close-up of a point  $z$  on the positive boundary.



# What is the margin?


Close-up of a point  $z$  on the positive boundary.



Let  $\hat{w}$  be the unit vector in the direction of  $w$ , i.e.  $\hat{w} = w/\|w\|$ .

# What is the margin?

Close-up of a point  $z$  on the positive boundary.



Let  $\hat{w}$  be the unit vector in the direction of  $w$ , i.e.  $\hat{w} = w/\|w\|$ .


Then  $z - \gamma\hat{w}$  is on the separator, so

$$w \cdot (z - \gamma\hat{w}) + b = 0 \Rightarrow \gamma w \cdot \hat{w} = w \cdot z + b = 1 \Rightarrow \gamma = 1/\|w\|$$



# What is the margin?

Close-up of a point  $z$  on the positive boundary.



Let  $\hat{w}$  be the unit vector in the direction of  $w$ , i.e.  $\hat{w} = w/\|w\|$ .

Then  $z - \gamma\hat{w}$  is on the separator, so

$$w \cdot (z - \gamma\hat{w}) + b = 0 \Rightarrow \gamma w \cdot \hat{w} = w \cdot z + b = 1 \Rightarrow \gamma = 1/\|w\|$$

In short: to maximize the margin, minimize  $\|w\|$ .

# Maximum-margin linear classifier

- Given  $(x^{(1)}, y^{(1)}), \dots, (x^{(n)}, y^{(n)}) \in \mathbb{R}^p \times \{-1, +1\}$ .

$$\begin{array}{ll} \text{(PRIMAL)} & \min_{w \in \mathbb{R}^p, b \in \mathbb{R}} \quad \frac{1}{2} \|w\|^2 \\ \text{s.t.:} & y^{(i)}(w \cdot x^{(i)} + b) \geq 1 \quad \text{for all } i = 1, 2, \dots, n \end{array}$$

# Maximum-margin linear classifier

- Given  $(x^{(1)}, y^{(1)}), \dots, (x^{(n)}, y^{(n)}) \in \mathbb{R}^p \times \{-1, +1\}$ .

$$\begin{array}{ll} \text{(PRIMAL)} & \min_{w \in \mathbb{R}^p, b \in \mathbb{R}} \quad \frac{1}{2} \|w\|^2 \\ \text{s.t.:} & y^{(i)}(w \cdot x^{(i)} + b) \geq 1 \quad \text{for all } i = 1, 2, \dots, n \end{array}$$

- This is a convex optimization problem:
  - Convex objective function
  - Linear constraints

# Maximum-margin linear classifier

- Given  $(x^{(1)}, y^{(1)}), \dots, (x^{(n)}, y^{(n)}) \in \mathbb{R}^p \times \{-1, +1\}$ .

$$\begin{aligned} \text{(PRIMAL)} \quad & \min_{w \in \mathbb{R}^p, b \in \mathbb{R}} \quad \frac{1}{2} \|w\|^2 \\ \text{s.t.:} \quad & y^{(i)}(w \cdot x^{(i)} + b) \geq 1 \quad \text{for all } i = 1, 2, \dots, n \end{aligned}$$

- This is a convex optimization problem:
  - Convex objective function
  - Linear constraints
- It has a dual maximization problem with the same optimum value.

$$\begin{aligned} \text{(DUAL)} \quad & \max_{\alpha \in \mathbb{R}^n} \quad \sum_{i=1}^n \alpha_i - \frac{1}{2} \sum_{i,j=1}^n \alpha_i \alpha_j y^{(i)} y^{(j)} (x^{(i)} \cdot x^{(j)}) \\ & \text{s.t.:} \quad \sum_{i=1}^n \alpha_i y^{(i)} = 0 \\ & \quad \quad \alpha \geq 0 \end{aligned}$$

## Complementary slackness

$$\begin{aligned} \text{(PRIMAL)} \quad & \min_{w \in \mathbb{R}^p, b \in \mathbb{R}} \quad \frac{1}{2} \|w\|^2 \\ \text{s.t.:} \quad & y^{(i)}(w \cdot x^{(i)} + b) \geq 1 \quad \text{for all } i = 1, 2, \dots, n \end{aligned}$$


$$\begin{aligned} \text{(DUAL)} \quad & \max_{\alpha \in \mathbb{R}^n} \quad \sum_{i=1}^n \alpha_i - \frac{1}{2} \sum_{i,j=1}^n \alpha_i \alpha_j y^{(i)} y^{(j)} (x^{(i)} \cdot x^{(j)}) \\ \text{s.t.:} \quad & \sum_{i=1}^n \alpha_i y^{(i)} = 0 \\ & \alpha \geq 0 \end{aligned}$$

At optimality,  $w = \sum_{i=1}^n \alpha_i y^{(i)} x^{(i)}$  and moreover

$$\alpha_i > 0 \quad \Rightarrow \quad y^{(i)}(w \cdot x^{(i)} + b) = 1$$


Points  $x^{(i)}$  with  $\alpha_i > 0$  are called **support vectors**.

# Support vectors




Linear classifier  $w = \sum_{i=1}^n \alpha_i y^{(i)} x^{(i)}$  is a function of just the support vectors.

## Small example: Iris data set



## Small example: Iris data set






# The non-separable case

Idea: allow each data point  $x^{(i)}$  some slack  $\xi_i$ .

$$\begin{aligned} \text{(PRIMAL)} \quad & \min_{w \in \mathbb{R}^p, b \in \mathbb{R}, \xi \in \mathbb{R}^n} \quad \frac{1}{2} \|w\|^2 + C \sum_{i=1}^n \xi_i \\ \text{s.t.:} \quad & y^{(i)}(w \cdot x^{(i)} + b) \geq 1 - \xi_i \quad \text{for all } i = 1, 2, \dots, n \\ & \xi \geq 0 \end{aligned}$$



## Dual for general case

$$\begin{aligned} \text{(PRIMAL)} \quad & \min_{w \in \mathbb{R}^p, b \in \mathbb{R}, \xi \in \mathbb{R}^n} \quad \frac{1}{2} \|w\|^2 + C \sum_{i=1}^n \xi_i \\ \text{s.t.:} \quad & y^{(i)}(w \cdot x^{(i)} + b) \geq 1 - \xi_i \quad \text{for all } i = 1, 2, \dots, n \\ & \xi \geq 0 \end{aligned}$$

$$\begin{aligned} \text{(DUAL)} \quad & \max_{\alpha \in \mathbb{R}^n} \quad \sum_{i=1}^n \alpha_i - \frac{1}{2} \sum_{i,j=1}^n \alpha_i \alpha_j y^{(i)} y^{(j)} (x^{(i)} \cdot x^{(j)}) \\ \text{s.t.:} \quad & \sum_{i=1}^n \alpha_i y^{(i)} = 0 \\ & 0 \leq \alpha_i \leq C \end{aligned}$$


At optimality,  $w = \sum_i \alpha_i y^{(i)} x^{(i)}$ , with

$$0 < \alpha_i < C \quad \Rightarrow \quad y^{(i)}(w \cdot x^{(i)} + b) = 1$$

$$\alpha_i = C \quad \Rightarrow \quad y^{(i)}(w \cdot x^{(i)} + b) = 1 - \xi_i$$


# Wine data set

Here  $C = 1.0$




# Wine data set

Here  $C = 1.0$




## Back to Iris

$C = 10$




## Back to Iris

$C = 10$




## Back to Iris

$C = 3$



## Back to Iris


$C = 2$






## Back to Iris

$C = 1$




## Back to Iris

$C = 0.5$




## Back to Iris

$C = 0.1$



## Back to Iris

$C = 0.01$



## Convex surrogates for 0-1 loss

Want a separator  $w$  that misclassifies as few training points as possible.

- 0-1 loss: charge  $1(y(w \cdot x) < 0)$  for each  $(x, y)$

Problem: this is NP-hard.

# Convex surrogates for 0-1 loss

Want a separator  $w$  that misclassifies as few training points as possible.

- 0-1 loss: charge  $1(y(w \cdot x) < 0)$  for each  $(x, y)$

Problem: this is NP-hard.

Instead, use **convex** loss functions.

- Hinge loss (SVM): charge  $(1 - y(w \cdot x))_+$
- Logistic loss: charge  $\ln(1 + e^{-y(w \cdot x)})$

# Convex surrogates for 0-1 loss


Want a separator  $w$  that misclassifies as few training points as possible.

- 0-1 loss: charge  $1(y(w \cdot x) < 0)$  for each  $(x, y)$

Problem: this is NP-hard.

Instead, use **convex** loss functions.

- Hinge loss (SVM): charge  $(1 - y(w \cdot x))_+$
- Logistic loss: charge  $\ln(1 + e^{-y(w \cdot x)})$



# A high-level view of optimization

## Unconstrained optimization

Logistic regression: find the vector  $w \in \mathbb{R}^p$  that minimizes

$$L(w) = \sum_{i=1}^n \ln(1 + \exp(-y^{(i)}(w \cdot x^{(i)}))).$$

We know how to check such problems for convexity and how to solve using gradient descent or Newton methods.



# A high-level view of optimization

## Unconstrained optimization

Logistic regression: find the vector  $w \in \mathbb{R}^p$  that minimizes

$$L(w) = \sum_{i=1}^n \ln(1 + \exp(-y^{(i)}(w \cdot x^{(i)}))).$$

We know how to check such problems for convexity and how to solve using gradient descent or Newton methods.

## Constrained optimization

Support vector machine: find  $w \in \mathbb{R}^p$  and  $b \in \mathbb{R}$  that minimize

$$L(w) = \|w\|^2$$

subject to the constraints

$$y^{(i)}(w \cdot x^{(i)} + b) \geq 1$$

What problems of this kind are easy to solve?

# Constrained optimization

Write the optimization problem in a standardized form:

$$\begin{aligned} \min f_o(z) \\ f_i(z) \leq 0 \quad \text{for } i = 1, \dots, m \\ h_i(z) = 0 \quad \text{for } i = 1, \dots, n \end{aligned}$$

Special cases that can be solved (relatively) easily:

- **Linear programs.**  
 $f_o, f_i, h_i$  are all linear functions.
- **Convex programs.**  
 $f_o, f_i$  are convex functions. The  $h_i$  are linear functions.

## Example: regression with $\ell_1$ loss

Given  $(x^{(1)}, y^{(1)}), \dots, (x^{(n)}, y^{(n)}) \in \mathbb{R}^p \times \mathbb{R}$ , find  $w \in \mathbb{R}^p$  minimizing

$$L(w) = \sum_{i=1}^n |y^{(i)} - (w \cdot x^{(i)})|.$$

## Example: regression with $\ell_1$ loss

Given  $(x^{(1)}, y^{(1)}), \dots, (x^{(n)}, y^{(n)}) \in \mathbb{R}^p \times \mathbb{R}$ , find  $w \in \mathbb{R}^p$  minimizing

$$L(w) = \sum_{i=1}^n |y^{(i)} - (w \cdot x^{(i)})|.$$

Equivalently: let  $X$  be the  $n \times p$  matrix with rows  $x^{(i)}$ , and  $y = (y^{(1)}, \dots, y^{(n)})$ . Then  $L(w) = \|Xw - y\|_1$ .

## Example: regression with $\ell_1$ loss

Given  $(x^{(1)}, y^{(1)}), \dots, (x^{(n)}, y^{(n)}) \in \mathbb{R}^p \times \mathbb{R}$ , find  $w \in \mathbb{R}^p$  minimizing

$$L(w) = \sum_{i=1}^n |y^{(i)} - (w \cdot x^{(i)})|.$$

Equivalently: let  $X$  be the  $n \times p$  matrix with rows  $x^{(i)}$ , and  $y = (y^{(1)}, \dots, y^{(n)})$ . Then  $L(w) = \|Xw - y\|_1$ .

Optimization problem in  $p + n$  variables,  $w \in \mathbb{R}^p$  and  $z \in \mathbb{R}^n$ :

$$\begin{aligned} \min \quad & \sum_{i=1}^n z_i \\ & y^{(i)} - w \cdot x^{(i)} \leq z_i, \quad i = 1, 2, \dots, n \\ & w \cdot x^{(i)} - y^{(i)} \leq z_i, \quad i = 1, 2, \dots, n \end{aligned}$$

## Example: regression with $\ell_1$ loss

Given  $(x^{(1)}, y^{(1)}), \dots, (x^{(n)}, y^{(n)}) \in \mathbb{R}^p \times \mathbb{R}$ , find  $w \in \mathbb{R}^p$  minimizing

$$L(w) = \sum_{i=1}^n |y^{(i)} - (w \cdot x^{(i)})|.$$

Equivalently: let  $X$  be the  $n \times p$  matrix with rows  $x^{(i)}$ , and  $y = (y^{(1)}, \dots, y^{(n)})$ . Then  $L(w) = \|Xw - y\|_1$ .

Optimization problem in  $p + n$  variables,  $w \in \mathbb{R}^p$  and  $z \in \mathbb{R}^n$ :

$$\begin{aligned} \min \quad & \sum_{i=1}^n z_i \\ & y^{(i)} - w \cdot x^{(i)} \leq z_i, \quad i = 1, 2, \dots, n \\ & w \cdot x^{(i)} - y^{(i)} \leq z_i, \quad i = 1, 2, \dots, n \end{aligned}$$

A linear program.

# The dual of an optimization problem

Take any optimization problem, convex or not:

$$\begin{aligned} \min f_o(z) \\ f_i(z) \leq 0 \quad \text{for } i = 1, \dots, m \\ h_i(z) = 0 \quad \text{for } i = 1, \dots, n \end{aligned}$$

Call this the *primal* problem.

# The dual of an optimization problem

Take any optimization problem, convex or not:

$$\begin{aligned} \min f_o(z) \\ f_i(z) &\leq 0 \quad \text{for } i = 1, \dots, m \\ h_i(z) &= 0 \quad \text{for } i = 1, \dots, n \end{aligned}$$

Call this the *primal* problem.

There is a *dual* optimization problem over  $n + m$  variables:

- $\lambda \in \mathbb{R}^m$ , one variable for each primal inequality
- $\nu \in \mathbb{R}^n$ , one variable for each primal equality

It is a maximization problem:

$$\begin{aligned} \max g(\lambda, \nu) \\ \lambda &\geq 0 \end{aligned}$$



# The dual of an optimization problem

Take any optimization problem, convex or not:

$$\begin{aligned} \min f_o(z) \\ f_i(z) &\leq 0 \quad \text{for } i = 1, \dots, m \\ h_i(z) &= 0 \quad \text{for } i = 1, \dots, n \end{aligned}$$

Call this the *primal* problem.

There is a *dual* optimization problem over  $n + m$  variables:

- $\lambda \in \mathbb{R}^m$ , one variable for each primal inequality
- $\nu \in \mathbb{R}^n$ , one variable for each primal equality

It is a maximization problem:

$$\begin{aligned} \max g(\lambda, \nu) \\ \lambda &\geq 0 \end{aligned}$$

Constructing the dual is straightforward. But interpreting it is not.

# Duality and complementary slackness

Let  $z^*$  and  $\lambda^*, \nu^*$  be the optimal primal and dual solutions.

# Duality and complementary slackness

Let  $z^*$  and  $\lambda^*, \nu^*$  be the optimal primal and dual solutions.

- Weak duality.

Dual solution is at most the primal solution:  $g(\lambda^*, \nu^*) \leq f_o(z^*)$ .

# Duality and complementary slackness

Let  $z^*$  and  $\lambda^*, \nu^*$  be the optimal primal and dual solutions.

- **Weak duality.**

Dual solution is at most the primal solution:  $g(\lambda^*, \nu^*) \leq f_o(z^*)$ .

- **Strong duality.**

If primal problem is convex then (almost always, with an easily checkable condition) the primal and dual solutions are equal.

# Duality and complementary slackness

Let  $z^*$  and  $\lambda^*, \nu^*$  be the optimal primal and dual solutions.

- **Weak duality.**

Dual solution is at most the primal solution:  $g(\lambda^*, \nu^*) \leq f_o(z^*)$ .

- **Strong duality.**

If primal problem is convex then (almost always, with an easily checkable condition) the primal and dual solutions are equal.

- **Complementary slackness.**

If primal and dual solutions are equal, then for any  $i = 1, \dots, m$ ,

$$\lambda_i^* > 0 \Rightarrow f_i(z^*) = 0.$$

# Duality and complementary slackness

Let  $z^*$  and  $\lambda^*, \nu^*$  be the optimal primal and dual solutions.

- **Weak duality.**

Dual solution is at most the primal solution:  $g(\lambda^*, \nu^*) \leq f_o(z^*)$ .

- **Strong duality.**

If primal problem is convex then (almost always, with an easily checkable condition) the primal and dual solutions are equal.

- **Complementary slackness.**

If primal and dual solutions are equal, then for any  $i = 1, \dots, m$ ,

$$\lambda_i^* > 0 \Rightarrow f_i(z^*) = 0.$$

- **KKT (Karush-Kuhn-Tucker) conditions.**

If the  $f_i$  and  $h_i$  are differentiable, these are first-derivative-equals-zero conditions that hold at optimality.