

# Some linear algebra background

CSE 250B

# Matrix-vector notation

Vector  $x \in \mathbb{R}^p$  and matrix  $M \in \mathbb{R}^{r \times p}$ :

$$x = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ \vdots \\ x_p \end{pmatrix}, \quad M = \begin{pmatrix} M_{11} & M_{12} & \cdots & M_{1p} \\ M_{21} & M_{22} & \cdots & M_{2p} \\ \vdots & \vdots & \ddots & \vdots \\ M_{r1} & M_{r2} & \cdots & M_{rp} \end{pmatrix}.$$

# Matrix-vector notation

Vector  $x \in \mathbb{R}^p$  and matrix  $M \in \mathbb{R}^{r \times p}$ :

$$x = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ \vdots \\ x_p \end{pmatrix}, \quad M = \begin{pmatrix} M_{11} & M_{12} & \cdots & M_{1p} \\ M_{21} & M_{22} & \cdots & M_{2p} \\ \vdots & \vdots & \ddots & \vdots \\ M_{r1} & M_{r2} & \cdots & M_{rp} \end{pmatrix}.$$

Transpose  $x^T$  and  $M^T \in \mathbb{R}^{p \times r}$ :

$$x^T = (x_1 \quad x_2 \quad \cdots \quad x_p), \quad M^T = \begin{pmatrix} M_{11} & \cdots & M_{r1} \\ M_{12} & \cdots & M_{r2} \\ M_{13} & \cdots & M_{r3} \\ \vdots & \ddots & \vdots \\ M_{1p} & \cdots & M_{rp} \end{pmatrix}.$$

# Matrix-vector notation

Vector  $x \in \mathbb{R}^p$  and matrix  $M \in \mathbb{R}^{r \times p}$ :

$$x = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ \vdots \\ x_p \end{pmatrix}, \quad M = \begin{pmatrix} M_{11} & M_{12} & \cdots & M_{1p} \\ M_{21} & M_{22} & \cdots & M_{2p} \\ \vdots & \vdots & \ddots & \vdots \\ M_{r1} & M_{r2} & \cdots & M_{rp} \end{pmatrix}.$$

Transpose  $x^T$  and  $M^T \in \mathbb{R}^{p \times r}$ :

$$x^T = (x_1 \quad x_2 \quad \cdots \quad x_p), \quad M^T = \begin{pmatrix} M_{11} & \cdots & M_{r1} \\ M_{12} & \cdots & M_{r2} \\ M_{13} & \cdots & M_{r3} \\ \vdots & \ddots & \vdots \\ M_{1p} & \cdots & M_{rp} \end{pmatrix}.$$

Properties of transpose:  $(A^T)^T = A$  and  $(AB)^T = B^T A^T$ .

# Dot product

Dot product of vectors  $x, y \in \mathbb{R}^p$ :


$$x \cdot y = x^T y = x_1 y_1 + \cdots + x_p y_p.$$

# Dot product

Dot product of vectors  $x, y \in \mathbb{R}^p$ :

$$x \cdot y = x^T y = x_1 y_1 + \cdots + x_p y_p.$$

This tells us the angle between  $x$  and  $y$ :




$$\cos \theta = \frac{x \cdot y}{\|x\| \|y\|}.$$

# Dot product

Dot product of vectors  $x, y \in \mathbb{R}^p$ :

$$x \cdot y = x^T y = x_1 y_1 + \cdots + x_p y_p.$$

This tells us the angle between  $x$  and  $y$ :



$$\cos \theta = \frac{x \cdot y}{\|x\| \|y\|}.$$


Easiest when  $x, y$  are *unit vectors* (length 1): then  $\cos \theta = x \cdot y$ .

# Dot product

Dot product of vectors  $x, y \in \mathbb{R}^p$ :

$$x \cdot y = x^T y = x_1 y_1 + \cdots + x_p y_p.$$

This tells us the angle between  $x$  and  $y$ :



$$\cos \theta = \frac{x \cdot y}{\|x\| \|y\|}.$$

Easiest when  $x, y$  are *unit vectors* (length 1): then  $\cos \theta = x \cdot y$ .

$x$  is orthogonal (at right angles) to  $y$  iff  $x \cdot y = 0$




# Dot product

Dot product of vectors  $x, y \in \mathbb{R}^p$ :

$$x \cdot y = x^T y = x_1 y_1 + \cdots + x_p y_p.$$

This tells us the angle between  $x$  and  $y$ :



$$\cos \theta = \frac{x \cdot y}{\|x\| \|y\|}.$$

Easiest when  $x, y$  are *unit vectors* (length 1): then  $\cos \theta = x \cdot y$ .

$x$  is orthogonal (at right angles) to  $y$  iff  $x \cdot y = 0$ .

What is  $x \cdot x$ ?

# Matrix-vector products

If  $M \in \mathbb{R}^{r \times p}$  and  $x \in \mathbb{R}^p$  then

$$Mx = \begin{pmatrix} \leftarrow M_1 \rightarrow \\ \leftarrow M_2 \rightarrow \\ \vdots \\ \leftarrow M_r \rightarrow \end{pmatrix} \begin{pmatrix} \uparrow \\ x \\ \downarrow \end{pmatrix} = \begin{pmatrix} M_1 \cdot x \\ M_2 \cdot x \\ \vdots \\ M_r \cdot x \end{pmatrix}$$

# Matrix-vector products

If  $M \in \mathbb{R}^{r \times p}$  and  $x \in \mathbb{R}^p$  then

$$Mx = \begin{pmatrix} \leftarrow M_1 \rightarrow \\ \leftarrow M_2 \rightarrow \\ \vdots \\ \leftarrow M_r \rightarrow \end{pmatrix} \begin{pmatrix} \uparrow \\ x \\ \downarrow \end{pmatrix} = \begin{pmatrix} M_1 \cdot x \\ M_2 \cdot x \\ \vdots \\ M_r \cdot x \end{pmatrix}$$

This mapping  $x \mapsto Mx$  is a **linear function** from  $\mathbb{R}^p$  to  $\mathbb{R}^r$ :

$$M(x + x') = Mx + Mx'.$$

# Matrix-vector products

If  $M \in \mathbb{R}^{r \times p}$  and  $x \in \mathbb{R}^p$  then

$$Mx = \begin{pmatrix} \leftarrow M_1 \rightarrow \\ \leftarrow M_2 \rightarrow \\ \vdots \\ \leftarrow M_r \rightarrow \end{pmatrix} \begin{pmatrix} \updownarrow \\ x \\ \updownarrow \end{pmatrix} = \begin{pmatrix} M_1 \cdot x \\ M_2 \cdot x \\ \vdots \\ M_r \cdot x \end{pmatrix}$$

This mapping  $x \mapsto Mx$  is a **linear function** from  $\mathbb{R}^p$  to  $\mathbb{R}^r$ :

$$M(x + x') = Mx + Mx'.$$

If  $M \in \mathbb{R}^{p \times p}$  and  $x \in \mathbb{R}^p$  then  $x \mapsto x^T Mx$  is a **quadratic function** from  $\mathbb{R}^p$  to  $\mathbb{R}$ :

$$x^T Mx = \sum_{i,j=1}^p M_{ij} x_i x_j.$$

## Quick quiz

- 1 Write the linear function  $f(x_1, x_2) = 3x_1 + 2x_2$  using vector notation (here,  $x_1, x_2 \in \mathbb{R}$ ).

## Quick quiz

- 1 Write the linear function  $f(x_1, x_2) = 3x_1 + 2x_2$  using vector notation (here,  $x_1, x_2 \in \mathbb{R}$ ).
- 2 Write the quadratic function  $f(x_1, x_2) = x_1^2 + 2x_1x_2 + 3x_2^2$  using matrices and vectors.

## Quick quiz

- 1 Write the linear function  $f(x_1, x_2) = 3x_1 + 2x_2$  using vector notation (here,  $x_1, x_2 \in \mathbb{R}$ ).
- 2 Write the quadratic function  $f(x_1, x_2) = x_1^2 + 2x_1x_2 + 3x_2^2$  using matrices and vectors.
- 3 A linear function from  $\mathbb{R}^3$  to  $\mathbb{R}^3$  is given by the matrix

$$M = \begin{pmatrix} 1 & 1 & 1 \\ 2 & 2 & 2 \\ 3 & 3 & 3 \end{pmatrix}.$$

As  $x$  varies, does  $Mx$  fill up all of  $\mathbb{R}^3$ ?

# A hierarchy of square matrices

Square

$$M \in \mathbb{R}^{p \times p}$$



Symmetric

$$M = M^T$$



Positive  
semidefinite

$$z^T M z \geq 0 \text{ for all } z \in \mathbb{R}^p$$



Positive  
definite

$$z^T M z > 0 \text{ for all } z \neq 0$$



## Quick quiz

Symmetric matrix  $M$  is positive semidefinite (psd) if  $z^T M z \geq 0$  for all  $z$ .

① PSD or not?

- $\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$

## Quick quiz

Symmetric matrix  $M$  is positive semidefinite (psd) if  $z^T M z \geq 0$  for all  $z$ .

① PSD or not?

- $\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$

- $\begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix}$

## Quick quiz

Symmetric matrix  $M$  is positive semidefinite (psd) if  $z^T M z \geq 0$  for all  $z$ .

① PSD or not?

- $\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$

- $\begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix}$

② A diagonal matrix is PSD if and only if ???

## Quick quiz

Symmetric matrix  $M$  is positive semidefinite (psd) if  $z^T M z \geq 0$  for all  $z$ .

① PSD or not?

- $\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$

- $\begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix}$

② A diagonal matrix is PSD if and only if ???

③ Show: If  $M, N$  are of the same size and PSD and  $M + N$  is PSD.

# Checking if a matrix is PSD

Useful fact: a matrix  $M$  is PSD iff it can be written in the form  $UU^T$  for some matrix  $U$ .

# Checking if a matrix is PSD

Useful fact: a matrix  $M$  is PSD iff it can be written in the form  $UU^T$  for some matrix  $U$ .

Quick check: say  $U \in \mathbb{R}^{r \times p}$  and  $M = UU^T$ .

# Checking if a matrix is PSD

Useful fact: a matrix  $M$  is PSD iff it can be written in the form  $UU^T$  for some matrix  $U$ .

Quick check: say  $U \in \mathbb{R}^{r \times p}$  and  $M = UU^T$ .

- 1  $M$  is square.

# Checking if a matrix is PSD

Useful fact: a matrix  $M$  is PSD iff it can be written in the form  $UU^T$  for some matrix  $U$ .

Quick check: say  $U \in \mathbb{R}^{r \times p}$  and  $M = UU^T$ .

- 1  $M$  is square.
- 2  $M$  is symmetric.



# Checking if a matrix is PSD

Useful fact: a matrix  $M$  is PSD iff it can be written in the form  $UU^T$  for some matrix  $U$ .

Quick check: say  $U \in \mathbb{R}^{r \times p}$  and  $M = UU^T$ .

- ①  $M$  is square.
- ②  $M$  is symmetric.
- ③ Pick any  $z \in \mathbb{R}^r$ . Then

$$\begin{aligned} z^T M z &= z^T U U^T z = (z^T U)(U^T z) \\ &= (U^T z)^T (U^T z) = \|U^T z\|^2 \geq 0. \end{aligned}$$

# Checking if a matrix is PSD

Useful fact: a matrix  $M$  is PSD iff it can be written in the form  $UU^T$  for some matrix  $U$ .

Quick check: say  $U \in \mathbb{R}^{r \times p}$  and  $M = UU^T$ .

- ①  $M$  is square.
- ②  $M$  is symmetric.
- ③ Pick any  $z \in \mathbb{R}^r$ . Then

$$\begin{aligned} z^T M z &= z^T U U^T z = (z^T U)(U^T z) \\ &= (U^T z)^T (U^T z) = \|U^T z\|^2 \geq 0. \end{aligned}$$

Another useful fact: any covariance matrix is PSD. (Same argument, along with linearity of expectation.)

# Eigenvalues and eigenvectors

- ① Any matrix  $M$  defines a linear transformation  $x \mapsto Mx$ .

# Eigenvalues and eigenvectors

- 1 Any matrix  $M$  defines a linear transformation  $x \mapsto Mx$ .
- 2 We'd like to understand the nature of these transformations. The easiest case is when  $M$  is **diagonal**:

$$\underbrace{\begin{pmatrix} 2 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 10 \end{pmatrix}}_M \underbrace{\begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}}_x = \underbrace{\begin{pmatrix} 2x_1 \\ -x_2 \\ 10x_3 \end{pmatrix}}_{Mx}$$

In this case,  $M$  simply scales each coordinate separately.

# Eigenvalues and eigenvectors

- 1 Any matrix  $M$  defines a linear transformation  $x \mapsto Mx$ .
- 2 We'd like to understand the nature of these transformations. The easiest case is when  $M$  is **diagonal**:

$$\underbrace{\begin{pmatrix} 2 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 10 \end{pmatrix}}_M \underbrace{\begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}}_x = \underbrace{\begin{pmatrix} 2x_1 \\ -x_2 \\ 10x_3 \end{pmatrix}}_{Mx}$$

In this case,  $M$  simply scales each coordinate separately.

- 3 What about more general matrices that are symmetric but not necessarily diagonal? They also just scale coordinates separately, but in a **different coordinate system**.

# Eigenvalues and eigenvectors

- 1 Any matrix  $M$  defines a linear transformation  $x \mapsto Mx$ .
- 2 We'd like to understand the nature of these transformations. The easiest case is when  $M$  is **diagonal**:

$$\underbrace{\begin{pmatrix} 2 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 10 \end{pmatrix}}_M \underbrace{\begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}}_x = \underbrace{\begin{pmatrix} 2x_1 \\ -x_2 \\ 10x_3 \end{pmatrix}}_{Mx}$$

In this case,  $M$  simply scales each coordinate separately.

- 3 What about more general matrices that are symmetric but not necessarily diagonal? They also just scale coordinates separately, but in a **different coordinate system**.

Let  $M$  be a  $p \times p$  matrix.

We say  $u \in \mathbb{R}^p$  is an **eigenvector** if  $M$  maps  $u$  onto the same direction, that is,

$$Mu = \lambda u$$

for some scaling constant  $\lambda$ . This  $\lambda$  is the **eigenvalue** associated with  $u$ .

# Eigenvalues and eigenvectors: examples

We say  $u$  is an eigenvector of  $M$ , with eigenvalue  $\lambda$ , if  $Mu = \lambda u$ .

# Eigenvalues and eigenvectors: examples

**We say  $u$  is an eigenvector of  $M$ , with eigenvalue  $\lambda$ , if  $Mu = \lambda u$ .**

Question: What are the eigenvectors and eigenvalues of:

$$M = \begin{pmatrix} 2 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 10 \end{pmatrix} ?$$



# Eigenvalues and eigenvectors: examples

**We say  $u$  is an eigenvector of  $M$ , with eigenvalue  $\lambda$ , if  $Mu = \lambda u$ .**

Question: What are the eigenvectors and eigenvalues of:

$$M = \begin{pmatrix} 2 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 10 \end{pmatrix} ?$$

Answer: Eigenvectors  $e_1, e_2, e_3$ , with corresponding eigenvalues  $2, -1, 10$ .

## Eigenvalues and eigenvectors: examples

**We say  $u$  is an eigenvector of  $M$ , with eigenvalue  $\lambda$ , if  $Mu = \lambda u$ .**

Question: What are the eigenvectors and eigenvalues of:

$$M = \begin{pmatrix} 2 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 10 \end{pmatrix} ?$$

Answer: Eigenvectors  $e_1, e_2, e_3$ , with corresponding eigenvalues 2,  $-1$ , 10.

Question: Matrix  $M = \begin{pmatrix} 3 & 1 \\ 1 & 3 \end{pmatrix}$  has eigenvectors

$$u_1 = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \quad u_2 = \frac{1}{\sqrt{2}} \begin{pmatrix} -1 \\ 1 \end{pmatrix}.$$

What are the corresponding eigenvalues?

# Eigenvalues and eigenvectors: examples

**We say  $u$  is an eigenvector of  $M$ , with eigenvalue  $\lambda$ , if  $Mu = \lambda u$ .**

Question: What are the eigenvectors and eigenvalues of:

$$M = \begin{pmatrix} 2 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 10 \end{pmatrix} ?$$

Answer: Eigenvectors  $e_1, e_2, e_3$ , with corresponding eigenvalues  $2, -1, 10$ .

Question: Matrix  $M = \begin{pmatrix} 3 & 1 \\ 1 & 3 \end{pmatrix}$  has eigenvectors

$$u_1 = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \quad u_2 = \frac{1}{\sqrt{2}} \begin{pmatrix} -1 \\ 1 \end{pmatrix}.$$

What are the corresponding eigenvalues?

Answer:  $\lambda_1 = 4$  and  $\lambda_2 = 2$ .

## Eigenvalues and eigenvectors: examples

**We say  $u$  is an eigenvector of  $M$ , with eigenvalue  $\lambda$ , if  $Mu = \lambda u$ .**

Question: What are the eigenvectors and eigenvalues of:

$$M = \begin{pmatrix} 2 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 10 \end{pmatrix} ?$$

Answer: Eigenvectors  $e_1, e_2, e_3$ , with corresponding eigenvalues  $2, -1, 10$ .

Question: Matrix  $M = \begin{pmatrix} 3 & 1 \\ 1 & 3 \end{pmatrix}$  has eigenvectors

$$u_1 = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \quad u_2 = \frac{1}{\sqrt{2}} \begin{pmatrix} -1 \\ 1 \end{pmatrix}.$$

What are the corresponding eigenvalues?

Answer:  $\lambda_1 = 4$  and  $\lambda_2 = 2$ .

In both cases the eigenvectors form an orthonormal basis.

# Eigenvectors of a real symmetric matrix

**Theorem.** Let  $M$  be any real symmetric  $p \times p$  matrix. Then  $M$  has

- $p$  eigenvalues  $\lambda_1, \dots, \lambda_p$
- corresponding eigenvectors  $u_1, \dots, u_p \in \mathbb{R}^p$  that are **orthonormal**:

$$u_i \cdot u_j = \begin{cases} 0 & \text{if } i \neq j \\ 1 & \text{if } i = j \end{cases}$$

# Eigenvectors of a real symmetric matrix

**Theorem.** Let  $M$  be any real symmetric  $p \times p$  matrix. Then  $M$  has

- $p$  eigenvalues  $\lambda_1, \dots, \lambda_p$
- corresponding eigenvectors  $u_1, \dots, u_p \in \mathbb{R}^p$  that are **orthonormal**:

$$u_i \cdot u_j = \begin{cases} 0 & \text{if } i \neq j \\ 1 & \text{if } i = j \end{cases}$$

We can think of  $u_1, \dots, u_p$  as being the axes of the natural coordinate system for understanding  $M$ .

# Eigenvectors of a real symmetric matrix

**Theorem.** Let  $M$  be any real symmetric  $p \times p$  matrix. Then  $M$  has

- $p$  eigenvalues  $\lambda_1, \dots, \lambda_p$
- corresponding eigenvectors  $u_1, \dots, u_p \in \mathbb{R}^p$  that are **orthonormal**:

$$u_i \cdot u_j = \begin{cases} 0 & \text{if } i \neq j \\ 1 & \text{if } i = j \end{cases}$$

We can think of  $u_1, \dots, u_p$  as being the axes of the natural coordinate system for understanding  $M$ .

**Theorem.** Let  $M$  be any real symmetric  $p \times p$  matrix, and let  $\lambda_1, \dots, \lambda_p$  be its eigenvalues. Then:

- $M$  is positive semidefinite iff every  $\lambda_i$  is  $\geq 0$ .
- $M$  is positive definite iff every  $\lambda$  is  $> 0$ .