


Informative projections

DSE 220


Informative projection

Suppose we wanted just one feature for the following data.



Informative projection


Suppose we wanted just one feature for the following data.



- We could pick a single coordinate.

Informative projection


Suppose we wanted just one feature for the following data.



- We could pick a single coordinate.
- Or an arbitrary direction.

Informative projection


Suppose we wanted just one feature for the following data.



- We could pick a single coordinate.
- Or an arbitrary direction.

Informative projection


Suppose we wanted just one feature for the following data.




- We could pick a single coordinate.
- Or an arbitrary direction.

A good choice: the **direction of maximum variance**.


Two types of projection



Projection onto \mathbb{R} :




Projection onto a 1-d line in \mathbb{R}^2 :




Projection: formally

What is the projection of $x \in \mathbb{R}^p$ onto direction $u \in \mathbb{R}^p$ (where $\|u\| = 1$)?



Projection: formally

What is the projection of $x \in \mathbb{R}^p$ onto direction $u \in \mathbb{R}^p$ (where $\|u\| = 1$)?




As a one-dimensional value:

$$x \cdot u = u \cdot x = u^T x = \sum_{i=1}^p u_i x_i.$$

Projection: formally

What is the projection of $x \in \mathbb{R}^p$ onto direction $u \in \mathbb{R}^p$ (where $\|u\| = 1$)?



As a one-dimensional value:

$$x \cdot u = u \cdot x = u^T x = \sum_{i=1}^p u_i x_i.$$

As a p -dimensional vector:

$$(x \cdot u)u = uu^T x$$

“Move $x \cdot u$ units in direction u ”

Quick quiz

What is the projection of $x = \begin{pmatrix} 2 \\ 3 \end{pmatrix}$ onto the following directions?

Give, first, a one-dimensional value and, then, a two-dimensional vector.

- 1 The coordinate direction e_1 ?

Quick quiz

What is the projection of $x = \begin{pmatrix} 2 \\ 3 \end{pmatrix}$ onto the following directions?

Give, first, a one-dimensional value and, then, a two-dimensional vector.

① The coordinate direction e_1 ?

② The direction of $\begin{pmatrix} 1 \\ -1 \end{pmatrix}$?

Projection onto multiple directions

Want to project $x \in \mathbb{R}^p$ into the k -dimensional subspace defined by vectors $u_1, \dots, u_k \in \mathbb{R}^p$.

Projection onto multiple directions

Want to project $x \in \mathbb{R}^p$ into the k -dimensional subspace defined by vectors $u_1, \dots, u_k \in \mathbb{R}^p$.

This is easiest when the u_i 's are **orthonormal**:

- They each have length one.
- They are at right angles to each other: $u_i \cdot u_j = 0$ whenever $i \neq j$

Projection onto multiple directions

Want to project $x \in \mathbb{R}^P$ into the k -dimensional subspace defined by vectors $u_1, \dots, u_k \in \mathbb{R}^P$.

This is easiest when the u_i 's are **orthonormal**:

- They each have length one.
- They are at right angles to each other: $u_i \cdot u_j = 0$ whenever $i \neq j$

Then the projection, as a k -dimensional vector, is

$$(x \cdot u_1, x \cdot u_2, \dots, x \cdot u_k) = \underbrace{\begin{pmatrix} \longleftrightarrow u_1 \longrightarrow \\ \longleftrightarrow u_2 \longrightarrow \\ \vdots \\ \longleftrightarrow u_k \longrightarrow \end{pmatrix}}_{\text{call this } U^T} \begin{pmatrix} \updownarrow \\ x \\ \updownarrow \end{pmatrix}$$

Projection onto multiple directions

Want to project $x \in \mathbb{R}^p$ into the k -dimensional subspace defined by vectors $u_1, \dots, u_k \in \mathbb{R}^p$.

This is easiest when the u_i 's are **orthonormal**:

- They each have length one.
- They are at right angles to each other: $u_i \cdot u_j = 0$ whenever $i \neq j$

Then the projection, as a k -dimensional vector, is

$$(x \cdot u_1, x \cdot u_2, \dots, x \cdot u_k) = \underbrace{\begin{pmatrix} \longleftrightarrow u_1 \longrightarrow \\ \longleftrightarrow u_2 \longrightarrow \\ \vdots \\ \longleftrightarrow u_k \longrightarrow \end{pmatrix}}_{\text{call this } U^T} \begin{pmatrix} \updownarrow \\ x \\ \updownarrow \end{pmatrix}$$

As a p -dimensional vector, the projection is

$$(x \cdot u_1)u_1 + (x \cdot u_2)u_2 + \dots + (x \cdot u_k)u_k = UU^T x.$$

Projection onto multiple directions: example

Suppose data are in \mathbb{R}^4 and we want to project onto the first two coordinates.

Projection onto multiple directions: example

Suppose data are in \mathbb{R}^4 and we want to project onto the first two coordinates.

Take vectors $u_1 = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}$, $u_2 = \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \end{pmatrix}$ (notice: orthonormal)

Projection onto multiple directions: example

Suppose data are in \mathbb{R}^4 and we want to project onto the first two coordinates.

Take vectors $u_1 = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}$, $u_2 = \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \end{pmatrix}$ (notice: orthonormal)

Then write $U^T = \begin{pmatrix} \overleftarrow{\hspace{1.5cm}} u_1 \overrightarrow{\hspace{1.5cm}} \\ \overleftarrow{\hspace{1.5cm}} u_2 \overrightarrow{\hspace{1.5cm}} \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}$

Projection onto multiple directions: example

Suppose data are in \mathbb{R}^4 and we want to project onto the first two coordinates.

Take vectors $u_1 = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}$, $u_2 = \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \end{pmatrix}$ (notice: orthonormal)

Then write $U^T = \begin{pmatrix} \overleftarrow{\hspace{1.5cm}} u_1 \overrightarrow{\hspace{1.5cm}} \\ \overleftarrow{\hspace{1.5cm}} u_2 \overrightarrow{\hspace{1.5cm}} \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}$

The projection of $x \in \mathbb{R}^4$,
as a 2-d vector, is

$$U^T x = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$$

Projection onto multiple directions: example

Suppose data are in \mathbb{R}^4 and we want to project onto the first two coordinates.

Take vectors $u_1 = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}$, $u_2 = \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \end{pmatrix}$ (notice: orthonormal)

Then write $U^T = \begin{pmatrix} \longleftarrow u_1 \longrightarrow \\ \longleftarrow u_2 \longrightarrow \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}$

The projection of $x \in \mathbb{R}^4$,
as a 2-d vector, is

$$U^T x = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$$

The projection of x as a
4-d vector is

$$UU^T x = \begin{pmatrix} x_1 \\ x_2 \\ 0 \\ 0 \end{pmatrix}$$

Projection onto multiple directions: example

Suppose data are in \mathbb{R}^4 and we want to project onto the first two coordinates.

Take vectors $u_1 = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}$, $u_2 = \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \end{pmatrix}$ (notice: orthonormal)

Then write $U^T = \begin{pmatrix} \longleftarrow u_1 \longrightarrow \\ \longleftarrow u_2 \longrightarrow \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}$

The projection of $x \in \mathbb{R}^4$,
as a 2-d vector, is

$$U^T x = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$$

The projection of x as a
4-d vector is

$$UU^T x = \begin{pmatrix} x_1 \\ x_2 \\ 0 \\ 0 \end{pmatrix}$$

But we'll generally project along non-coordinate directions.

The best single direction

Suppose we need to map our data $x \in \mathbb{R}^p$ into just **one** dimension:

$$x \mapsto u \cdot x \quad \text{for some unit direction } u \in \mathbb{R}^p$$

What is the direction u of maximum variance?

The best single direction

Suppose we need to map our data $x \in \mathbb{R}^p$ into just **one** dimension:

$$x \mapsto u \cdot x \quad \text{for some unit direction } u \in \mathbb{R}^p$$

What is the direction u of maximum variance?

Theorem: Let Σ be the $p \times p$ covariance matrix of X . The variance of X in direction u is given by $u^T \Sigma u$.

The best single direction

Suppose we need to map our data $x \in \mathbb{R}^p$ into just **one** dimension:

$$x \mapsto u \cdot x \quad \text{for some unit direction } u \in \mathbb{R}^p$$

What is the direction u of maximum variance?

Theorem: Let Σ be the $p \times p$ covariance matrix of X . The variance of X in direction u is given by $u^T \Sigma u$.

- Suppose the mean of X is $\mu \in \mathbb{R}^p$. The projection $u^T X$ has mean

$$\mathbb{E}(u^T X) = u^T \mathbb{E}X = u^T \mu.$$

The best single direction

Suppose we need to map our data $x \in \mathbb{R}^p$ into just **one** dimension:

$$x \mapsto u \cdot x \quad \text{for some unit direction } u \in \mathbb{R}^p$$

What is the direction u of maximum variance?

Theorem: Let Σ be the $p \times p$ covariance matrix of X . The variance of X in direction u is given by $u^T \Sigma u$.

- Suppose the mean of X is $\mu \in \mathbb{R}^p$. The projection $u^T X$ has mean

$$\mathbb{E}(u^T X) = u^T \mathbb{E}X = u^T \mu.$$

- The variance of $u^T X$ is

$$\begin{aligned} \text{var}(u^T X) &= \mathbb{E}(u^T X - u^T \mu)^2 = \mathbb{E}(u^T (X - \mu)(X - \mu)^T u) \\ &= u^T \mathbb{E}(X - \mu)(X - \mu)^T u = u^T \Sigma u. \end{aligned}$$

The best single direction

Suppose we need to map our data $x \in \mathbb{R}^p$ into just **one** dimension:

$$x \mapsto u \cdot x \quad \text{for some unit direction } u \in \mathbb{R}^p$$

What is the direction u of maximum variance?

Theorem: Let Σ be the $p \times p$ covariance matrix of X . The variance of X in direction u is given by $u^T \Sigma u$.

- Suppose the mean of X is $\mu \in \mathbb{R}^p$. The projection $u^T X$ has mean


$$\mathbb{E}(u^T X) = u^T \mathbb{E}X = u^T \mu.$$

- The variance of $u^T X$ is

$$\begin{aligned} \text{var}(u^T X) &= \mathbb{E}(u^T X - u^T \mu)^2 = \mathbb{E}(u^T (X - \mu)(X - \mu)^T u) \\ &= u^T \mathbb{E}(X - \mu)(X - \mu)^T u = u^T \Sigma u. \end{aligned}$$

Another theorem: $u^T \Sigma u$ is maximized by setting u to the first **eigenvector** of Σ . The maximum value is the corresponding **eigenvalue**.

Best single direction: example



This direction is the **first eigenvector** of the 2×2 covariance matrix of the data.

The best k -dimensional projection

Let Σ be the $p \times p$ covariance matrix of X . Its **eigendecomposition** can be computed in $O(p^3)$ time and consists of:

- real **eigenvalues** $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_p$
- corresponding **eigenvectors** $u_1, \dots, u_p \in \mathbb{R}^p$ that are orthonormal: that is, each u_i has unit length and $u_i \cdot u_j = 0$ whenever $i \neq j$.

The best k -dimensional projection

Let Σ be the $p \times p$ covariance matrix of X . Its **eigendecomposition** can be computed in $O(p^3)$ time and consists of:

- real **eigenvalues** $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_p$
- corresponding **eigenvectors** $u_1, \dots, u_p \in \mathbb{R}^p$ that are orthonormal: that is, each u_i has unit length and $u_i \cdot u_j = 0$ whenever $i \neq j$.

Theorem: Suppose we want to map data $X \in \mathbb{R}^p$ to just k dimensions, while capturing as much of the variance of X as possible. The best choice of projection is:

$$x \mapsto (u_1 \cdot x, u_2 \cdot x, \dots, u_k \cdot x),$$

where u_i are the eigenvectors described above.

The best k -dimensional projection

Let Σ be the $p \times p$ covariance matrix of X . Its **eigendecomposition** can be computed in $O(p^3)$ time and consists of:

- real **eigenvalues** $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_p$
- corresponding **eigenvectors** $u_1, \dots, u_p \in \mathbb{R}^p$ that are orthonormal: that is, each u_i has unit length and $u_i \cdot u_j = 0$ whenever $i \neq j$.

Theorem: Suppose we want to map data $X \in \mathbb{R}^p$ to just k dimensions, while capturing as much of the variance of X as possible. The best choice of projection is:


$$x \mapsto (u_1 \cdot x, u_2 \cdot x, \dots, u_k \cdot x),$$

where u_i are the eigenvectors described above.

Projecting the data in this way is **principal component analysis (PCA)**.

Example: MNIST

Contrast coordinate projections with PCA:



MNIST: image reconstruction



Reconstruct this original image from its PCA projection to k dimensions.

MNIST: image reconstruction



Reconstruct this original image from its PCA projection to k dimensions.

$k = 200$



$k = 150$



$k = 100$



$k = 50$



MNIST: image reconstruction



Reconstruct this original image from its PCA projection to k dimensions.

$k = 200$



$k = 150$



$k = 100$



$k = 50$



Q: What are these reconstructions exactly?

MNIST: image reconstruction



Reconstruct this original image from its PCA projection to k dimensions.

$k = 200$



$k = 150$



$k = 100$



$k = 50$



Q: What are these reconstructions exactly?

A: Image x is reconstructed as $UU^T x$, where U is a $p \times k$ matrix whose columns are the top k eigenvectors of Σ .

Review: eigenvalues and eigenvectors

There are several steps to understanding these.

- 1 Any matrix M defines a function (or **transformation**) $x \mapsto Mx$.
- 2 If M is a $p \times q$ matrix, then this transformation maps vector $x \in \mathbb{R}^q$ to vector $Mx \in \mathbb{R}^p$.
- 3 We call it a **linear transformation** because $M(x + x') = Mx + Mx'$.
- 4 We'd like to understand the nature of these transformations. The easiest case is when M is **diagonal**:

$$\underbrace{\begin{pmatrix} 2 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 10 \end{pmatrix}}_M \underbrace{\begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}}_x = \underbrace{\begin{pmatrix} 2x_1 \\ -x_2 \\ 10x_3 \end{pmatrix}}_{Mx}$$

In this case, M simply scales each coordinate separately.

- 5 What about more general matrices that are symmetric but not necessarily diagonal? They also just scale coordinates separately, but in a **different coordinate system**.

Review: eigenvalues and eigenvectors

Let M be a $p \times p$ matrix.

We say $u \in \mathbb{R}^p$ is an **eigenvector** if M maps u onto the same direction, that is,

$$Mu = \lambda u$$

for some scaling constant λ . This λ is the **eigenvalue** associated with u .

Question: What are the eigenvectors and eigenvalues of:

$$M = \begin{pmatrix} 2 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 10 \end{pmatrix} ?$$

Review: eigenvalues and eigenvectors

Let M be a $p \times p$ matrix.

We say $u \in \mathbb{R}^p$ is an **eigenvector** if M maps u onto the same direction, that is,

$$Mu = \lambda u$$

for some scaling constant λ . This λ is the **eigenvalue** associated with u .

Question: What are the eigenvectors and eigenvalues of:

$$M = \begin{pmatrix} 2 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 10 \end{pmatrix} ?$$

Answer: Eigenvectors e_1, e_2, e_3 , with corresponding eigenvalues $2, -1, 10$.

Review: eigenvalues and eigenvectors

Let M be a $p \times p$ matrix.

We say $u \in \mathbb{R}^p$ is an **eigenvector** if M maps u onto the same direction, that is,

$$Mu = \lambda u$$

for some scaling constant λ . This λ is the **eigenvalue** associated with u .

Question: What are the eigenvectors and eigenvalues of:

$$M = \begin{pmatrix} 2 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 10 \end{pmatrix} ?$$

Answer: Eigenvectors e_1, e_2, e_3 , with corresponding eigenvalues $2, -1, 10$.

Notice that these eigenvectors form an orthonormal basis.

Eigenvectors of a real symmetric matrix

Theorem. Let M be any real symmetric $p \times p$ matrix. Then M has

- p eigenvalues $\lambda_1, \dots, \lambda_p$
- corresponding eigenvectors $u_1, \dots, u_p \in \mathbb{R}^p$ that are orthonormal

We can think of u_1, \dots, u_p as being the axes of the natural coordinate system for understanding M .

Eigenvectors of a real symmetric matrix

Theorem. Let M be any real symmetric $p \times p$ matrix. Then M has

- p eigenvalues $\lambda_1, \dots, \lambda_p$
- corresponding eigenvectors $u_1, \dots, u_p \in \mathbb{R}^p$ that are orthonormal

We can think of u_1, \dots, u_p as being the axes of the natural coordinate system for understanding M .

Example: consider the matrix

$$M = \begin{pmatrix} 3 & 1 \\ 1 & 3 \end{pmatrix}$$

It has eigenvectors

$$u_1 = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \quad u_2 = \frac{1}{\sqrt{2}} \begin{pmatrix} -1 \\ 1 \end{pmatrix}.$$

- Are these eigenvectors orthonormal?
- What are the corresponding eigenvalues?

Spectral decomposition

Theorem. Let M be any real symmetric $p \times p$ matrix. Then M has

- p eigenvalues $\lambda_1, \dots, \lambda_p$
- corresponding eigenvectors $u_1, \dots, u_p \in \mathbb{R}^p$ that are orthonormal

Spectral decomposition

Theorem. Let M be any real symmetric $p \times p$ matrix. Then M has

- p eigenvalues $\lambda_1, \dots, \lambda_p$
- corresponding eigenvectors $u_1, \dots, u_p \in \mathbb{R}^p$ that are orthonormal

Spectral decomposition: Here is another way to write M :

$$M = \underbrace{\begin{pmatrix} \uparrow & \uparrow & & \uparrow \\ u_1 & u_2 & \cdots & u_p \\ \downarrow & \downarrow & & \downarrow \end{pmatrix}}_{U: \text{ columns are eigenvectors}} \underbrace{\begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_p \end{pmatrix}}_{\Lambda: \text{ eigenvalues on diagonal}} \underbrace{\begin{pmatrix} \leftarrow u_1 \rightarrow \\ \leftarrow u_2 \rightarrow \\ \vdots \\ \leftarrow u_p \rightarrow \end{pmatrix}}_{U^T}$$

Spectral decomposition

Theorem. Let M be any real symmetric $p \times p$ matrix. Then M has

- p eigenvalues $\lambda_1, \dots, \lambda_p$
- corresponding eigenvectors $u_1, \dots, u_p \in \mathbb{R}^p$ that are orthonormal

Spectral decomposition: Here is another way to write M :

$$M = \underbrace{\begin{pmatrix} \uparrow & \uparrow & & \uparrow \\ u_1 & u_2 & \cdots & u_p \\ \downarrow & \downarrow & & \downarrow \end{pmatrix}}_{U: \text{ columns are eigenvectors}} \underbrace{\begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_p \end{pmatrix}}_{\Lambda: \text{ eigenvalues on diagonal}} \underbrace{\begin{pmatrix} \leftarrow u_1 \rightarrow \\ \leftarrow u_2 \rightarrow \\ \vdots \\ \leftarrow u_p \rightarrow \end{pmatrix}}_{U^T}$$

Thus $Mx = U\Lambda U^T x$, which can be interpreted as follows:

- U^T rewrites x in the $\{u_i\}$ coordinate system
- Λ is a simple coordinate scaling in that basis
- U then sends the scaled vector back into the usual coordinate basis

Spectral decomposition: example

Apply spectral decomposition to the matrix M we saw earlier:


$$M = \begin{pmatrix} 3 & 1 \\ 1 & 3 \end{pmatrix} = \underbrace{\frac{1}{\sqrt{2}} \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix}}_U \underbrace{\begin{pmatrix} 4 & 0 \\ 0 & 2 \end{pmatrix}}_\Lambda \underbrace{\frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix}}_{U^T}$$

Spectral decomposition: example

Apply spectral decomposition to the matrix M we saw earlier:

$$M = \begin{pmatrix} 3 & 1 \\ 1 & 3 \end{pmatrix} = \underbrace{\frac{1}{\sqrt{2}} \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix}}_U \underbrace{\begin{pmatrix} 4 & 0 \\ 0 & 2 \end{pmatrix}}_\Lambda \underbrace{\frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix}}_{U^T}$$

$$M \begin{pmatrix} 1 \\ 2 \end{pmatrix} = ???$$




Spectral decomposition: example

Apply spectral decomposition to the matrix M we saw earlier:

$$M = \begin{pmatrix} 3 & 1 \\ 1 & 3 \end{pmatrix} = \underbrace{\frac{1}{\sqrt{2}} \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix}}_U \underbrace{\begin{pmatrix} 4 & 0 \\ 0 & 2 \end{pmatrix}}_\Lambda \underbrace{\frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix}}_{U^T}$$

$$M \begin{pmatrix} 1 \\ 2 \end{pmatrix} = U \Lambda U^T \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$




Spectral decomposition: example

Apply spectral decomposition to the matrix M we saw earlier:

$$M = \begin{pmatrix} 3 & 1 \\ 1 & 3 \end{pmatrix} = \underbrace{\frac{1}{\sqrt{2}} \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix}}_U \underbrace{\begin{pmatrix} 4 & 0 \\ 0 & 2 \end{pmatrix}}_\Lambda \underbrace{\frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix}}_{U^T}$$

$$M \begin{pmatrix} 1 \\ 2 \end{pmatrix} = U \Lambda U^T \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$




Spectral decomposition: example

Apply spectral decomposition to the matrix M we saw earlier:

$$M = \begin{pmatrix} 3 & 1 \\ 1 & 3 \end{pmatrix} = \underbrace{\frac{1}{\sqrt{2}} \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix}}_U \underbrace{\begin{pmatrix} 4 & 0 \\ 0 & 2 \end{pmatrix}}_\Lambda \underbrace{\frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix}}_{U^T}$$

$$M \begin{pmatrix} 1 \\ 2 \end{pmatrix} = U \Lambda U^T \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$




Spectral decomposition: example

Apply spectral decomposition to the matrix M we saw earlier:

$$M = \begin{pmatrix} 3 & 1 \\ 1 & 3 \end{pmatrix} = \underbrace{\frac{1}{\sqrt{2}} \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix}}_U \underbrace{\begin{pmatrix} 4 & 0 \\ 0 & 2 \end{pmatrix}}_\Lambda \underbrace{\frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix}}_{U^T}$$

$$\begin{aligned} M \begin{pmatrix} 1 \\ 2 \end{pmatrix} &= U \Lambda U^T \begin{pmatrix} 1 \\ 2 \end{pmatrix} \\ &= U \Lambda \frac{1}{\sqrt{2}} \begin{pmatrix} 3 \\ 1 \end{pmatrix} \end{aligned}$$




Spectral decomposition: example

Apply spectral decomposition to the matrix M we saw earlier:


$$M = \begin{pmatrix} 3 & 1 \\ 1 & 3 \end{pmatrix} = \underbrace{\frac{1}{\sqrt{2}} \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix}}_U \underbrace{\begin{pmatrix} 4 & 0 \\ 0 & 2 \end{pmatrix}}_\Lambda \underbrace{\frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix}}_{U^T}$$

$$\begin{aligned} M \begin{pmatrix} 1 \\ 2 \end{pmatrix} &= U \Lambda U^T \begin{pmatrix} 1 \\ 2 \end{pmatrix} \\ &= U \Lambda \frac{1}{\sqrt{2}} \begin{pmatrix} 3 \\ 1 \end{pmatrix} \\ &= U \frac{1}{\sqrt{2}} \begin{pmatrix} 12 \\ 2 \end{pmatrix} \\ &= \begin{pmatrix} 5 \\ 7 \end{pmatrix} \end{aligned}$$



Principal component analysis: recap


Consider data vectors $X \in \mathbb{R}^p$.



Principal component analysis: recap

Consider data vectors $X \in \mathbb{R}^p$.


- The covariance matrix Σ is a $p \times p$ symmetric matrix.



Principal component analysis: recap

Consider data vectors $X \in \mathbb{R}^p$.


- The covariance matrix Σ is a $p \times p$ symmetric matrix.
- Get eigenvalues $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_p$, eigenvectors u_1, \dots, u_p .



Principal component analysis: recap

Consider data vectors $X \in \mathbb{R}^p$.


- The covariance matrix Σ is a $p \times p$ symmetric matrix.
- Get eigenvalues $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_p$, eigenvectors u_1, \dots, u_p .
- u_1, \dots, u_p is an alternative basis in which to represent the data.
- The variance of X in direction u_i is λ_i .



Principal component analysis: recap

Consider data vectors $X \in \mathbb{R}^p$.


- The covariance matrix Σ is a $p \times p$ symmetric matrix.
- Get eigenvalues $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_p$, eigenvectors u_1, \dots, u_p .
- u_1, \dots, u_p is an alternative basis in which to represent the data.
- The variance of X in direction u_i is λ_i .
- To project to k dimensions while losing as little as possible of the overall variance, use $x \mapsto (x \cdot u_1, \dots, x \cdot u_k)$.



Principal component analysis: recap

Consider data vectors $X \in \mathbb{R}^p$.

- The covariance matrix Σ is a $p \times p$ symmetric matrix.
- Get eigenvalues $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_p$, eigenvectors u_1, \dots, u_p .
- u_1, \dots, u_p is an alternative basis in which to represent the data.
- The variance of X in direction u_i is λ_i .
- To project to k dimensions while losing as little as possible of the overall variance, use $x \mapsto (x \cdot u_1, \dots, x \cdot u_k)$.



What is the covariance of the projected data?

Example: personality assessment

What are the dimensions along which personalities differ?

Example: personality assessment

What are the dimensions along which personalities differ?

- *Lexical hypothesis*: most important personality characteristics have become encoded in natural language.

Example: personality assessment

What are the dimensions along which personalities differ?

- *Lexical hypothesis*: most important personality characteristics have become encoded in natural language.
- Allport and Odbert (1936): sat down with the English dictionary and extracted all terms that could be used to distinguish one person's behavior from another's. Roughly 18000 words, of which 4500 could be described as personality traits.

Example: personality assessment

What are the dimensions along which personalities differ?

- *Lexical hypothesis*: most important personality characteristics have become encoded in natural language.
- Allport and Odbert (1936): sat down with the English dictionary and extracted all terms that could be used to distinguish one person's behavior from another's. Roughly 18000 words, of which 4500 could be described as personality traits.
- Step: group these words into (approximate) synonyms. This is done by manual clustering. e.g. Norman (1967):

Spirit	Jolly, merry, witty, lively, peppy
Talkativeness	Talkative, articulate, verbose, gossipy
Sociability	Companionable, social, outgoing
Spontaneity	Impulsive, carefree, playful, zany
Boisterousness	Mischievous, rowdy, loud, prankish
Adventure	Brave, venturesome, fearless, reckless
Energy	Active, assertive, dominant, energetic
Conceit	Boastful, conceited, egotistical
Vanity	Affected, vain, chic, dapper, jaunty
Indiscretion	Nosey, snoopy, indiscreet, meddling
Sensuality	Sexy, passionate, sensual, flirtatious

Example: personality assessment

What are the dimensions along which personalities differ?

- *Lexical hypothesis*: most important personality characteristics have become encoded in natural language.
- Allport and Odbert (1936): sat down with the English dictionary and extracted all terms that could be used to distinguish one person's behavior from another's. Roughly 18000 words, of which 4500 could be described as personality traits.
- Step: group these words into (approximate) synonyms. This is done by manual clustering. e.g. Norman (1967):

Spirit	Jolly, merry, witty, lively, peppy
Talkativeness	Talkative, articulate, verbose, gossipy
Sociability	Companionable, social, outgoing
Spontaneity	Impulsive, carefree, playful, zany
Boisterousness	Mischievous, rowdy, loud, prankish
Adventure	Brave, venturesome, fearless, reckless
Energy	Active, assertive, dominant, energetic
Conceit	Boastful, conceited, egotistical
Vanity	Affected, vain, chic, dapper, jaunty
Indiscretion	Nosey, snoopy, indiscreet, meddling
Sensuality	Sexy, passionate, sensual, flirtatious

- Data collection: Ask a variety of subjects to what extent each of these words describes them.

Personality assessment: the data

Matrix of data (1 = strongly disagree, 5 = strongly agree)

	shy	merry	tense	boastful	forgiving	quiet
Person 1	4	1	1	2	5	5
Person 2	1	4	4	5	2	1
Person 3	2	4	5	4	2	2
		:				

How to extract important directions?

Personality assessment: the data

Matrix of data (1 = strongly disagree, 5 = strongly agree)

	shy	merry	tense	boastful	forgiving	quiet
Person 1	4	1	1	2	5	5
Person 2	1	4	4	5	2	1
Person 3	2	4	5	4	2	2
		:				

How to extract important directions?

- Treat each column as a data point, find tight clusters
- Treat each row as a data point, apply PCA
- Other ideas: factor analysis, independent component analysis, ...

Personality assessment: the data

Matrix of data (1 = strongly disagree, 5 = strongly agree)

	shy	merry	tense	boastful	forgiving	quiet
Person 1	4	1	1	2	5	5
Person 2	1	4	4	5	2	1
Person 3	2	4	5	4	2	2
		⋮				


How to extract important directions?

- Treat each column as a data point, find tight clusters
- Treat each row as a data point, apply PCA
- Other ideas: factor analysis, independent component analysis, ...

Many of these yield similar results


What does PCA accomplish?

Example: suppose two traits (generosity, trust) are highly correlated, to the point where each person either answers “1” to both or “5” to both.




What does PCA accomplish?

Example: suppose two traits (generosity, trust) are highly correlated, to the point where each person either answers “1” to both or “5” to both.



What does PCA accomplish?

Example: suppose two traits (generosity, trust) are highly correlated, to the point where each person either answers “1” to both or “5” to both.



This single PCA dimension entirely accounts for the two traits.

The “Big Five” taxonomy

[illegible]

The “Big Five” taxonomy

Extraversion		Agreeableness		Conscientiousness		Neuroticism		Openness/Intellect	
Low	High	Low	High	Low	High	Low	High	Low	High
-.83 Quiet	.85 Talkative	-.52 Fault-finding	.87 Sympathetic	-.58 Careless	.80 Organized	-.39 Stable*	.73 Tense	-.74 Commonplace	.76 Wide interests
-.80 Reserved	.83 Assertive	-.48 Cold	.85 Kind	-.53 Disorderly	.80 Thorough	-.35 Calm*	.72 Anxious	-.73 Narrow interests	.76 Imaginative
-.75 Shy	.82 Active	-.45 Unfriendly	.85 Appreciative	-.50 Frivolous	.78 Placid	-.21 Contented*	.72 Nervous	-.67 Simple	.72 Intelligent
-.71 Silent	.82 Energetic	-.45 Quarrelsome	.84 Affectionate	-.49 Irresponsible	.78 Efficient	.14 Unemotional*	.71 Moody	-.55 Shallow	.73 Original
-.67 Withdrawn	.82 Outgoing	-.45 Hard-hearted	.84 Soft-hearted	-.40 Slipshod	.73 Responsible		.71 Worrying	-.47 Unintelligent	.68 Insightful
-.66 Retiring	.80 Outspoken	-.38 Unkind	.82 Warm	-.39 Undependable	.72 Reliable		.68 Touchy		.64 Curious
	.79 Dominant	-.33 Cruel	.81 Generous	-.37 Forgetful	.70 Dependable		.64 Fearful		.59 Sophisticated
	.73 Forceful	-.31 Stern*	.78 Trusting		.68 Conscientious		.63 High-strung		.59 Artistic
	.73 Enthusiastic	-.28 Thankless	.77 Helpful		.66 Precise		.63 Self-pitying		.59 Clever
	.68 Show-off	-.24 Stingy*	.77 Forgiving		.66 Practical		.60 Temperamental		.58 Inventive
	.68 Sociable		.74 Pleasant		.65 Deliberate		.59 Unstable		.56 Sharp-witted
	.64 Spunky		.73 Good-natured		.46 Painstaking		.58 Self-punishing		.55 Ingenious
	.64 Adventurous		.73 Friendly		.26 Cautious*		.54 Despondent		.45 Witty*
	.62 Noisy		.72 Cooperative				.51 Emotional		.45 Resourceful*
	.58 Bossy		.67 Gentle						.37 Wise
			.66 Unselfish						.33 Logical*
			.56 Praising						.29 Civilized*
			.51 Sensitive						.22 Foresighted*
									.21 Polished*
									.20 Dignified*

Many applications, such as online match-making.

Singular value decomposition (SVD)

For **symmetric** matrices, such as covariance matrices, we have seen:

- Results about existence of eigenvalues and eigenvectors
- The fact that the eigenvectors form an alternative basis
- The resulting spectral decomposition, which is used in PCA

But what about arbitrary matrices $M \in \mathbb{R}^{p \times q}$?

Singular value decomposition (SVD)

For **symmetric** matrices, such as covariance matrices, we have seen:

- Results about existence of eigenvalues and eigenvectors
- The fact that the eigenvectors form an alternative basis
- The resulting spectral decomposition, which is used in PCA

But what about arbitrary matrices $M \in \mathbb{R}^{p \times q}$?

Any $p \times q$ matrix (say $p \leq q$) has a **singular value decomposition**:

$$M = \underbrace{\begin{pmatrix} \uparrow & & \uparrow \\ u_1 & \cdots & u_p \\ \downarrow & & \downarrow \end{pmatrix}}_{p \times p \text{ matrix } U} \underbrace{\begin{pmatrix} \sigma_1 & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & \sigma_p \end{pmatrix}}_{p \times p \text{ matrix } \Lambda} \underbrace{\begin{pmatrix} \longleftarrow v_1 \longrightarrow \\ \vdots \\ \longleftarrow v_p \longrightarrow \end{pmatrix}}_{p \times q \text{ matrix } V^T}$$

- u_1, \dots, u_p are orthonormal vectors in \mathbb{R}^p
- v_1, \dots, v_p are orthonormal vectors in \mathbb{R}^q
- $\sigma_1 \geq \sigma_2 \geq \cdots \geq \sigma_p$ are **singular values**

Matrix approximation

We can **factor** any $p \times q$ matrix as $M = UW^T$:

$$\begin{aligned} M &= \begin{pmatrix} \uparrow & & \uparrow \\ u_1 & \cdots & u_p \\ \downarrow & & \downarrow \end{pmatrix} \begin{pmatrix} \sigma_1 & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & \sigma_p \end{pmatrix} \begin{pmatrix} \longleftarrow v_1 \longrightarrow \\ \vdots \\ \longleftarrow v_p \longrightarrow \end{pmatrix} \\ &= \underbrace{\begin{pmatrix} \uparrow & & \uparrow \\ u_1 & \cdots & u_p \\ \downarrow & & \downarrow \end{pmatrix}}_{p \times p \text{ matrix } U} \underbrace{\begin{pmatrix} \longleftarrow \sigma_1 v_1 \longrightarrow \\ \vdots \\ \longleftarrow \sigma_p v_p \longrightarrow \end{pmatrix}}_{p \times q \text{ matrix } W^T} \end{aligned}$$

Matrix approximation

We can **factor** any $p \times q$ matrix as $M = UW^T$:


$$\begin{aligned} M &= \begin{pmatrix} \uparrow & & \uparrow \\ u_1 & \cdots & u_p \\ \downarrow & & \downarrow \end{pmatrix} \begin{pmatrix} \sigma_1 & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & \sigma_p \end{pmatrix} \begin{pmatrix} \longleftarrow v_1 \longrightarrow \\ \vdots \\ \longleftarrow v_p \longrightarrow \end{pmatrix} \\ &= \underbrace{\begin{pmatrix} \uparrow & & \uparrow \\ u_1 & \cdots & u_p \\ \downarrow & & \downarrow \end{pmatrix}}_{p \times p \text{ matrix } U} \underbrace{\begin{pmatrix} \longleftarrow \sigma_1 v_1 \longrightarrow \\ \vdots \\ \longleftarrow \sigma_p v_p \longrightarrow \end{pmatrix}}_{p \times q \text{ matrix } W^T} \end{aligned}$$

A concise approximation to M : just take the first k columns of U and the first k rows of W^T , for $k < p$:

$$\hat{M} = \underbrace{\begin{pmatrix} \uparrow & & \uparrow \\ u_1 & \cdots & u_k \\ \downarrow & & \downarrow \end{pmatrix}}_{p \times k} \underbrace{\begin{pmatrix} \longleftarrow \sigma_1 v_1 \longrightarrow \\ \vdots \\ \longleftarrow \sigma_k v_k \longrightarrow \end{pmatrix}}_{k \times q}$$

Example: topic modeling

Blei (2012):



Latent semantic indexing (LSI)

Given a large corpus of n documents:

- Fix a vocabulary, say of V words.
- Bag-of-words representation for documents: each document becomes a vector of length V , with one coordinate per word.
- The corpus is an $n \times V$ matrix, one row per document.

	<i>cat</i>	<i>dog</i>	<i>house</i>	<i>boat</i>	<i>garden</i>	<i>...</i>
Doc 1	4	1	1	0	2	
Doc 2	0	0	3	1	0	
Doc 3	0	1	3	0	0	
		\vdots				

Latent semantic indexing (LSI)

Given a large corpus of n documents:

- Fix a vocabulary, say of V words.
- Bag-of-words representation for documents: each document becomes a vector of length V , with one coordinate per word.
- The corpus is an $n \times V$ matrix, one row per document.

	cat	dog	house	boat	garden	...
Doc 1	4	1	1	0	2	
Doc 2	0	0	3	1	0	
Doc 3	0	1	3	0	0	
		⋮				

Let's find a concise approximation to this matrix M .

Latent semantic indexing, cont'd

Use SVD to get an approximation to M : for small k ,

$$\underbrace{\begin{pmatrix} \leftarrow \text{doc 1} \rightarrow \\ \leftarrow \text{doc 2} \rightarrow \\ \leftarrow \text{doc 3} \rightarrow \\ \vdots \\ \leftarrow \text{doc } n \rightarrow \end{pmatrix}}_{n \times V \text{ matrix } M} \approx \underbrace{\begin{pmatrix} \leftarrow \theta_1 \rightarrow \\ \leftarrow \theta_2 \rightarrow \\ \leftarrow \theta_3 \rightarrow \\ \vdots \\ \leftarrow \theta_n \rightarrow \end{pmatrix}}_{n \times k \text{ matrix } \Theta} \underbrace{\begin{pmatrix} \leftarrow \psi_1 \rightarrow \\ \vdots \\ \leftarrow \psi_k \rightarrow \end{pmatrix}}_{k \times V \text{ matrix } \Psi}$$

Latent semantic indexing, cont'd

Use SVD to get an approximation to M : for small k ,

$$\underbrace{\begin{pmatrix} \leftarrow \text{doc 1} \rightarrow \\ \leftarrow \text{doc 2} \rightarrow \\ \leftarrow \text{doc 3} \rightarrow \\ \vdots \\ \leftarrow \text{doc } n \rightarrow \end{pmatrix}}_{n \times V \text{ matrix } M} \approx \underbrace{\begin{pmatrix} \leftarrow \theta_1 \rightarrow \\ \leftarrow \theta_2 \rightarrow \\ \leftarrow \theta_3 \rightarrow \\ \vdots \\ \leftarrow \theta_n \rightarrow \end{pmatrix}}_{n \times k \text{ matrix } \Theta} \underbrace{\begin{pmatrix} \leftarrow \psi_1 \rightarrow \\ \vdots \\ \leftarrow \psi_k \rightarrow \end{pmatrix}}_{k \times V \text{ matrix } \Psi}$$

Think of this as a *topic model* with k topics.

- ψ_j is a vector of length V describing topic j : coefficient ψ_{jw} is large if word w appears often in that topic.
- Each document is a combination of topics: θ_{ij} is the weight of topic j in document i .

Latent semantic indexing, cont'd

Use SVD to get an approximation to M : for small k ,

$$\underbrace{\begin{pmatrix} \leftarrow \text{doc 1} \rightarrow \\ \leftarrow \text{doc 2} \rightarrow \\ \leftarrow \text{doc 3} \rightarrow \\ \vdots \\ \leftarrow \text{doc } n \rightarrow \end{pmatrix}}_{n \times V \text{ matrix } M} \approx \underbrace{\begin{pmatrix} \leftarrow \theta_1 \rightarrow \\ \leftarrow \theta_2 \rightarrow \\ \leftarrow \theta_3 \rightarrow \\ \vdots \\ \leftarrow \theta_n \rightarrow \end{pmatrix}}_{n \times k \text{ matrix } \Theta} \underbrace{\begin{pmatrix} \leftarrow \Psi_1 \rightarrow \\ \vdots \\ \leftarrow \Psi_k \rightarrow \end{pmatrix}}_{k \times V \text{ matrix } \Psi}$$

Think of this as a *topic model* with k topics.

- Ψ_j is a vector of length V describing topic j : coefficient Ψ_{jw} is large if word w appears often in that topic.
- Each document is a combination of topics: θ_{ij} is the weight of topic j in document i .

Document i originally represented by i th row of M , a vector in \mathbb{R}^V .
Can instead use $\theta_i \in \mathbb{R}^k$, a more concise “semantic” representation.

The rank of a matrix

Suppose we want to approximate a matrix M by a simpler matrix \hat{M} .
What is a suitable notion of “simple”?

The rank of a matrix

Suppose we want to approximate a matrix M by a simpler matrix \hat{M} .
What is a suitable notion of “simple”?

- Let's say M and \hat{M} are $p \times q$, where $p \leq q$.
- Treat each row of \hat{M} as a data point in \mathbb{R}^q .
- We can think of the data as “simple” if it actually lies in a low-dimensional subspace.
- If the rows lie in k -dimensional subspace, we say that \hat{M} has **rank** k .

The rank of a matrix

Suppose we want to approximate a matrix M by a simpler matrix \hat{M} .
What is a suitable notion of “simple”?

- Let's say M and \hat{M} are $p \times q$, where $p \leq q$.
- Treat each row of \hat{M} as a data point in \mathbb{R}^q .
- We can think of the data as “simple” if it actually lies in a low-dimensional subspace.
- If the rows lie in k -dimensional subspace, we say that \hat{M} has **rank** k .

The **rank** of a matrix is the number of linearly independent rows.

The rank of a matrix

Suppose we want to approximate a matrix M by a simpler matrix \hat{M} .
What is a suitable notion of “simple”?

- Let's say M and \hat{M} are $p \times q$, where $p \leq q$.
- Treat each row of \hat{M} as a data point in \mathbb{R}^q .
- We can think of the data as “simple” if it actually lies in a low-dimensional subspace.
- If the rows lie in k -dimensional subspace, we say that \hat{M} has **rank** k .

The **rank** of a matrix is the number of linearly independent rows.

Low-rank approximation: given $M \in \mathbb{R}^{p \times q}$ and an integer k , find the matrix $\hat{M} \in \mathbb{R}^{p \times q}$ that is the best rank- k approximation to M .

The rank of a matrix

Suppose we want to approximate a matrix M by a simpler matrix \hat{M} .
What is a suitable notion of “simple”?

- Let's say M and \hat{M} are $p \times q$, where $p \leq q$.
- Treat each row of \hat{M} as a data point in \mathbb{R}^q .
- We can think of the data as “simple” if it actually lies in a low-dimensional subspace.
- If the rows lie in k -dimensional subspace, we say that \hat{M} has **rank** k .

The **rank** of a matrix is the number of linearly independent rows.

Low-rank approximation: given $M \in \mathbb{R}^{p \times q}$ and an integer k , find the matrix $\hat{M} \in \mathbb{R}^{p \times q}$ that is the best rank- k approximation to M .

That is, find \hat{M} so that

- \hat{M} has $\text{rank} \leq k$
- The approximation error $\sum_{i,j} (M_{ij} - \hat{M}_{ij})^2$ is minimized.

The rank of a matrix

Suppose we want to approximate a matrix M by a simpler matrix \hat{M} . What is a suitable notion of “simple”?

- Let's say M and \hat{M} are $p \times q$, where $p \leq q$.
- Treat each row of \hat{M} as a data point in \mathbb{R}^q .
- We can think of the data as “simple” if it actually lies in a low-dimensional subspace.
- If the rows lie in k -dimensional subspace, we say that \hat{M} has **rank** k .

The **rank** of a matrix is the number of linearly independent rows.

Low-rank approximation: given $M \in \mathbb{R}^{p \times q}$ and an integer k , find the matrix $\hat{M} \in \mathbb{R}^{p \times q}$ that is the best rank- k approximation to M .

That is, find \hat{M} so that

- \hat{M} has $\text{rank} \leq k$
- The approximation error $\sum_{i,j} (M_{ij} - \hat{M}_{ij})^2$ is minimized.

We can get \hat{M} directly from the singular value decomposition of M .

Low-rank approximation

Recall: Singular value decomposition of $p \times q$ matrix M (with $p \leq q$):

$$M = \begin{pmatrix} \uparrow & & \uparrow \\ u_1 & \cdots & u_p \\ \downarrow & & \downarrow \end{pmatrix} \begin{pmatrix} \sigma_1 & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & \sigma_p \end{pmatrix} \begin{pmatrix} \longleftarrow v_1 \longrightarrow \\ \vdots \\ \longleftarrow v_p \longrightarrow \end{pmatrix}$$

- u_1, \dots, u_p is an orthonormal basis of \mathbb{R}^p
- v_1, \dots, v_q is an orthonormal basis of \mathbb{R}^q
- $\sigma_1 \geq \cdots \geq \sigma_p$ are **singular values**

Low-rank approximation

Recall: Singular value decomposition of $p \times q$ matrix M (with $p \leq q$):

$$M = \begin{pmatrix} \uparrow & & \uparrow \\ u_1 & \cdots & u_p \\ \downarrow & & \downarrow \end{pmatrix} \begin{pmatrix} \sigma_1 & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & \sigma_p \end{pmatrix} \begin{pmatrix} \longleftarrow v_1 \longrightarrow \\ \vdots \\ \longleftarrow v_p \longrightarrow \end{pmatrix}$$

- u_1, \dots, u_p is an orthonormal basis of \mathbb{R}^p
- v_1, \dots, v_q is an orthonormal basis of \mathbb{R}^q
- $\sigma_1 \geq \cdots \geq \sigma_p$ are **singular values**

The **best rank- k approximation** to M , for any $k \leq p$, is then

$$\hat{M} = \underbrace{\begin{pmatrix} \uparrow & & \uparrow \\ u_1 & \cdots & u_k \\ \downarrow & & \downarrow \end{pmatrix}}_{p \times k} \underbrace{\begin{pmatrix} \sigma_1 & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & \sigma_k \end{pmatrix}}_{k \times k} \underbrace{\begin{pmatrix} \longleftarrow v_1 \longrightarrow \\ \vdots \\ \longleftarrow v_k \longrightarrow \end{pmatrix}}_{k \times q}$$

Example: Collaborative filtering

Details and images from Koren, Bell, Volinsky (2009).

Recommender systems: matching customers with products.

- Given: data on prior purchases/interests of users
- Recommend: further products of interest

Prototypical example: Netflix.

Example: Collaborative filtering

Details and images from Koren, Bell, Volinsky (2009).

Recommender systems: matching customers with products.

- Given: data on prior purchases/interests of users
- Recommend: further products of interest

Prototypical example: Netflix.

A successful approach: **collaborative filtering**.

- Model dependencies between different products, and between different users.
- Can give reasonable recommendations to a relatively new user.

Example: Collaborative filtering

Details and images from Koren, Bell, Volinsky (2009).

Recommender systems: matching customers with products.

- Given: data on prior purchases/interests of users
- Recommend: further products of interest

Prototypical example: Netflix.


A successful approach: **collaborative filtering**.

- Model dependencies between different products, and between different users.
- Can give reasonable recommendations to a relatively new user.


Two strategies for collaborative filtering:

- Neighborhood methods
- Latent factor methods

Neighborhood methods



Latent factor methods



The matrix factorization approach

User ratings are assembled in a large matrix M :

	<i>Star Wars</i>	<i>Matrix</i>	<i>Casablanca</i>	<i>Camelot</i>	<i>Godfather</i>	...
User 1	5	5	2	0	0	
User 2	0	0	3	4	5	
User 3	0	0	5	0	0	
		⋮				

- Not rated = 0, otherwise scores 1-5.
- For n users and p movies, this has size $n \times p$.
- Most of the entries are unavailable, and we'd like to predict these.

The matrix factorization approach

User ratings are assembled in a large matrix M :

	Star Wars	Matrix	Casablanca	Camelot	Godfather	...
User 1	5	5	2	0	0	
User 2	0	0	3	4	5	
User 3	0	0	5	0	0	
		⋮				

- Not rated = 0, otherwise scores 1-5.
- For n users and p movies, this has size $n \times p$.
- Most of the entries are unavailable, and we'd like to predict these.

Idea: Find the best low-rank approximation of M , and use it to fill in the missing entries.

User and movie factors

Best rank- k approximation is of the form $M \approx UW^T$:

$$\underbrace{\begin{pmatrix} \leftarrow \text{user 1} \rightarrow \\ \leftarrow \text{user 2} \rightarrow \\ \leftarrow \text{user 3} \rightarrow \\ \vdots \\ \leftarrow \text{user } n \rightarrow \end{pmatrix}}_{n \times p \text{ matrix } M} \approx \underbrace{\begin{pmatrix} \leftarrow u_1 \rightarrow \\ \leftarrow u_2 \rightarrow \\ \leftarrow u_3 \rightarrow \\ \vdots \\ \leftarrow u_n \rightarrow \end{pmatrix}}_{n \times k \text{ matrix } U} \underbrace{\begin{pmatrix} \uparrow w_1 & \uparrow w_2 & \cdots & \uparrow w_p \\ \downarrow & \downarrow & & \downarrow \end{pmatrix}}_{k \times p \text{ matrix } W^T}$$

User and movie factors

Best rank- k approximation is of the form $M \approx UW^T$:

$$\underbrace{\begin{pmatrix} \leftarrow \text{user 1} \rightarrow \\ \leftarrow \text{user 2} \rightarrow \\ \leftarrow \text{user 3} \rightarrow \\ \vdots \\ \leftarrow \text{user } n \rightarrow \end{pmatrix}}_{n \times p \text{ matrix } M} \approx \underbrace{\begin{pmatrix} \leftarrow u_1 \rightarrow \\ \leftarrow u_2 \rightarrow \\ \leftarrow u_3 \rightarrow \\ \vdots \\ \leftarrow u_n \rightarrow \end{pmatrix}}_{n \times k \text{ matrix } U} \underbrace{\begin{pmatrix} \uparrow w_1 & \uparrow w_2 & \cdots & \uparrow w_p \\ \downarrow & \downarrow & & \downarrow \end{pmatrix}}_{k \times p \text{ matrix } W^T}$$

Thus user i 's rating of movie j is approximated as

$$M_{ij} \approx u_i \cdot w_j$$

User and movie factors

Best rank- k approximation is of the form $M \approx UW^T$:

$$\underbrace{\begin{pmatrix} \leftarrow \text{user 1} \rightarrow \\ \leftarrow \text{user 2} \rightarrow \\ \leftarrow \text{user 3} \rightarrow \\ \vdots \\ \leftarrow \text{user } n \rightarrow \end{pmatrix}}_{n \times p \text{ matrix } M} \approx \underbrace{\begin{pmatrix} \leftarrow u_1 \rightarrow \\ \leftarrow u_2 \rightarrow \\ \leftarrow u_3 \rightarrow \\ \vdots \\ \leftarrow u_n \rightarrow \end{pmatrix}}_{n \times k \text{ matrix } U} \underbrace{\begin{pmatrix} \uparrow w_1 & \uparrow w_2 & \cdots & \uparrow w_p \\ \downarrow & \downarrow & & \downarrow \end{pmatrix}}_{k \times p \text{ matrix } W^T}$$

Thus user i 's rating of movie j is approximated as

$$M_{ij} \approx u_i \cdot w_j$$

This “latent” representation embeds users and movies within the same k -dimensional space:

- Represent i th user by $u_i \in \mathbb{R}^k$
- Represent j th movie by $w_j \in \mathbb{R}^k$

Top two Netflix factors

