



[Home](#) | [About Us](#) | [Success Stories](#) | [Farmers Association](#) | [Farmers' Innovation](#) | [Publications](#) | [Contact](#)

Horticultural crops :: Vegetables:: Tomato

Septoria Leaf Spot: *Septoria lycopersici*

Symptoms:

- Small, round to irregular spots with grey centre and dark margin on leaves.
- Spots coalesce and leaves are blighted.
- Complete defoliation.
- Stems and flowers are affected.



Irregular spots on leaves



Leaf spots

Causal Organism – *Septoria lycopersici* Speg.

Mycelium : Hyaline and septate

Fruiting body : Pycnidium: globose, without distinct ostiole

Pathogen: Pycnidiospore : filiform, septate, hyaline

Survival and Mode of Spread

Survival: Infected plant debris

Primary spread: Mycelium or conidia

Secondary spread: Conidia through rain splash or wind

Favourable conditions

- High humidity .

Integrated Disease Management

Cultural:

- Removal and destruction of affected plant parts

Biological :

- Seed treatment :Trichoderma asperellum @ 4g/kg seeds

Chemical:

- Fluxapyroxad 250 g/l + Pyraclostrobin 250 g/l SC - 200-250 ml/ha.

Source of Information:

- Crop production guide – Horticulture 2020.

Acknowledgement:

- Dr.N. Indra, Associate Professor, Department of Plant Pathology, TNAU, Coimbatore.
- Dr.P.Renukadevi, Professor, Department of Plant Pathology, TNAU, Coimbatore.
- Photo courtesy Dr.S.Vinodkumar.

Updated on Jun 2023

[Home](#) | [About Us](#) | [Success Stories](#) | [Farmers Association](#) | [Farmers' Innovation](#) | [Publications](#) | [Disclaimer](#) | [Contact Us](#)

© TNAU 2008 - 2024 All Rights Reserved.