Group Theory

By Abhi Mondal

Source

Group Theory for Physicists with Applications by *Pichai Ramadevi and Varun Dubey*

Keywords: Group theory, Symmetry groups, Cyclic groups, Conjugacy class

Contents

1 Motivation **2**

2 Definition of a Group **2**
2.1 Examples . . . 3
2.2 Subgroup . . . 3

3 Basics of Groups' Properties **3**
3.1 Multiplication Table . . . 3
3.2 Order of a Group . . . 3
3.3 Generator Set . . . 4
3.4 Example of a Non-Abelian Structure . . . 5
3.5 Cosets . . . 5
3.6 Equality of Cosets . . . 5
3.7 Cosets Partition the Group . . . 5
3.8 Size of a Coset . . . 6
3.9 Normal Subgroups . . . 6

4 Factor Groups **6**
4.1 Examples . . . 7
4.2 Order of a Factor Group . . . 7
4.3 Important Remarks on Factor Groups . . . 7

5 Conjugacy Classes **9**

A Basic Notions from Ring Theory **10**
A.1 The Ring $\mathbb{Z}_n$ . . . 10
A.2 Examples. . . . 10
A.3 Connection with Group Theory . . . 10

B Modulo Arithmetic and the Structure of $\mathbb{Z}_n$ **11**
B.1 Congruence Modulo n . . . 11
B.2 Residue Classes . . . 11
B.3 Arithmetic in $\mathbb{Z}_n$ . . . 11
B.4 Examples . . . 11
B.5 Algebraic Properties . . . 11
B.6 Connection with Cyclic Groups . . . 12
B.7 Geometric Interpretation . . . 12

C Algebraic Hierarchies **13**
C.1 Structures with One Operation . . . 13
C.2 Structures with Two Operations . . . 13
C.3 Conceptual Perspective . . . 13

D Special Other Groups **14**
D.1 The Quaternion Group Q . . . 14
D.2 Dihedral Group . . . 15

§1 Motivation

The group theory help us to exploit the symmetry without directly solving the differential equation explicitly. But the physicists are not that lazy! They are more curious to know the nature more intuitively!

- 1-D Harmonic Oscillator $\longrightarrow SO(2)$ symmetry
- Hydrogen atom $\longrightarrow$ Spherical symmetry
- Electrons in atom $\longrightarrow$ Permutation symmetry

Discrete Groups Cyclic group, Permutation group, point group, Irreducible representations, Character tables and applications in solid state physics.
Continuous Groups Space-time symmetry, translational symmetry, rotational symmetry. Examples: $SU(2)$, $SU(3)$ (covering group applications).

$$\begin{array}{cc} \text{Rotation of Square} & \text{Rotation of Circle} \\ \downarrow & \downarrow \\ \text{Discrete} & \text{Continuum} \end{array}$$

We can write for Square by $\{I, R_{\pi/2}, R_{\pi}, R_{3\pi/2}\}$ while for Circle, we need rotation by any angle θ which is **continuous**. The rotation group of a circle is Abelian.

§2 Definition of a Group

A set $G = \{a, b, c, \dots\}$ with a binary operation $\cdot$ is called a group if:

1. Closure: $a \cdot b \in G$
2. Associativity: $(a \cdot b) \cdot c = a \cdot (b \cdot c)$
3. Identity element: $\exists e \in G$ such that $ae = ea = a$
4. Inverse: $\forall a \in G, \exists a^{-1} \in G$ such that $aa^{-1} = a^{-1}a = e$

2.1 Examples

1. $\mathbb{R}$ under addition is an Abelian group.
2. $\mathbb{Z}$ under addition is an Abelian group.
3. $\mathbb{R} - \{0\}$ under multiplication is Abelian.
4. The set of all 2×2 matrices under matrix multiplication is not Abelian in general.

If for any group G with $a, b \in G$

$$a \cdot b = b \cdot a$$

then G is called an **Abelian group**.

2.2 Subgroup

If $H \subseteq G$ and H itself satisfies group properties under the same operation, then H is called a subgroup of G. In notation we can write $H \leq G$.

§3 Basics of Groups' Properties

3.1 Multiplication Table

A multiplication table helps verify closure and understand the structure of a finite group.
Example: Consider the rotation group of a square

$$G = \{e, R_{\pi/2}, R_{\pi}, R_{3\pi/2}\}$$

	e	$R_{\pi/2}$	R_{π}	$R_{3\pi/2}$
e	e	$R_{\pi/2}$	R_{π}	$R_{3\pi/2}$
$R_{\pi/2}$	$R_{\pi/2}$	R_{π}	$R_{3\pi/2}$	e
R_{π}	R_{π}	$R_{3\pi/2}$	e	$R_{\pi/2}$
$R_{3\pi/2}$	$R_{3\pi/2}$	e	$R_{\pi/2}$	R_{π}

3.2 Order of a Group

The number of elements in G is called the **order** of G, denoted by $|G|$. For the above example:

$$|G| = 4$$

A group with finite number of elements is called a **finite group**.

3.3 Generator Set

A subset S of elements of a group G is said to generate G if every element of G can be expressed as a finite product of finite powers of elements (or their inverses) of S in some order. For *finite* S, the elements of the minimal set S that generates a group G are called the **generators** of the group and S itself is called the *generating set*.

A group whose generating set contains a *single element* is said to be a **cyclic group**.

Example 3.1. Cyclic Group C_n with $a^n = e$

$$C_n = \{e, a, a^2, \ldots, a^{n-1}\}$$

The order of each element divides n.

Example 3.2. C_3 with $a^3 = e$

$$C_3 = \{e, a, a^2\}$$

Multiplication table:

	e	a	a^2
e	e	a	a^2
a	a	a^2	e
a^2	a^2	e	a

Example 3.3. The *Klein group* V_4 has elements with generator set $\{a, b\}$:

$$V_4 = \{e, a, b, ab\}$$

with properties:

$$a^2 = b^2 = e, \quad ab = ba$$

It is Abelian group but not cyclic.

Example 3.4. Consider a group of a generator set $\{a, b\}$ with the following relations :

$$a^3 = e, \qquad b^2 = e, \qquad ab = b^2 a$$

The group elements are:

$$G = \{e, a, a^2, b, ab, a^2 b\}$$

This is a *non-Abelian* group of order 6. We denoted this group with $\mathfrak{S}(3)$ and called it **Symmetric Group**.
Multiplication Table:

$\mathfrak{S}(3)$	e	a	b	b^2	ab	ab^2
e	e	a	b	b^2	ab	ab^2
a	a	e	ab	ab^2	b	b^2
b	b	ab^2	b^2	e	a	ab
b^2	b^2	ab	e	b	ab^2	a
ab	ab	b^2	ab^2	a	e	b
ab^2	ab^2	b	a	ab	b^2	e

3.4 Example of a Non-Abelian Structure

Consider a group with relations:

$$a^2 = e, \qquad b^2 = e$$

but

$$ab \neq ba.$$

In such a case, the elements can be:

$$G = \{e, a, b, ab, ba, aba, bab, \dots\}$$

The multiplication is determined by the defining relations.

3.5 Cosets

Let G be a group and $H \leq G$ a subgroup. For any $a \in G$, the *left coset* of H determined by a is

$$aH = \{ah \mid h \in H\}. \tag{1}$$

Thus, we fix a and multiply it with every element of H. Since the identity element $e \in H$, we have

$$a = ae \in aH, \tag{2}$$

Similarly we have *right coset*

3.6 Equality of Cosets

Suppose aH and bH have a common element. Then there exist $h_1, h_2 \in H$ such that

$$ah_1 = bh_2. \tag{3}$$

Multiplying on the left by a^{-1} gives

$$a^{-1}b = h_1 h_2^{-1}. \tag{4}$$

Since H is a subgroup, $h_1 h_2^{-1} \in H$, hence

$$a^{-1}b \in H. \tag{5}$$

Conversely, if $a^{-1}b \in H$, then $b = ah$ for some $h \in H$, and therefore

$$bH = aH. \tag{6}$$

Thus,

$$aH = bH \iff a^{-1}b \in H. \tag{7}$$

Hence two cosets are either identical or disjoint.

3.7 Cosets Partition the Group

Every element $g \in G$ belongs to the coset gH. If $g \in aH$, then $g = ah$ for some $h \in H$. Therefore, the cosets are disjoint and their union equals G:

$$G = \bigcup_{a \in G} aH. \tag{8}$$

So the cosets of H form a partition of G.

3.8 Size of a Coset

Define the map

$$\phi : H \to aH, \qquad \phi(h) = ah. \tag{9}$$

This map is bijective. Hence,

$$|aH| = |H|. \tag{10}$$

Thus every coset has exactly the same number of elements as H.

Lagrange's Theorem

If G is finite and $H \leq G$, then G is a disjoint union of k distinct cosets:

$$G = a_1 H \cup a_2 H \cup \cdots \cup a_k H. \tag{11}$$

Since each coset has $|H|$ elements,

$$|G| = k \cdot |H| = [G : H] \cdot |H|. \tag{12}$$

The symbol $[G : H]$ is called the **index** of H in G. It represents the number of distinct left cosets of H in G. Therefore,

$$|H| \mid |G|. \tag{13}$$

i.e. The size of every subgroup must divide the size of the group.

3.9 Normal Subgroups

A subgroup H is called **normal** if

$$aH = Ha \quad \text{for all } a \in G, \tag{14}$$

or equivalently,

$$aHa^{-1} = H. \tag{15}$$

In this case, the set of cosets forms the *quotient group* G/H.

- If G is abelian then every subgroup of G is normal.
- In case G has no non-trivial normal subgroups then G is said to be a **simple group**. Eg. Groups of prime order.

§4 Factor Groups

Let G be a group and let $N \trianglelefteq G$ be a normal subgroup. The *factor group* (or *quotient group*) of G by N is defined as

$$G/N = \{gN \mid g \in G\}, \tag{16}$$

where each element gN is a *left coset* of N in G. The group operation on G/N is given by

$$(gN)(hN) = (gh)N. \tag{17}$$

The normality of N ensures that this operation is well-defined. Moreover,

$$|G/N| = \frac{|G|}{|N|}. \tag{18}$$

4.1 Examples

Example 1: $\mathbb{Z}/2\mathbb{Z}$

Consider the additive group $G = \mathbb{Z}$ and its subgroup

$$N = 2\mathbb{Z} = \{2k \mid k \in \mathbb{Z}\}. \tag{19}$$

The cosets are

$$0 + 2\mathbb{Z} = \{\dots, -2, 0, 2, \dots\}, \tag{20}$$

$$1 + 2\mathbb{Z} = \{\dots, -1, 1, 3, \dots\}. \tag{21}$$

Thus, there are exactly two distinct cosets. Hence

$$\mathbb{Z}/2\mathbb{Z} \cong \mathbb{Z}_2. \tag{22}$$

Example 2: $\mathbb{Z}_6/\langle 3\rangle$

Let $G = \mathbb{Z}_6 = \{0, 1, 2, 3, 4, 5\}$ under addition modulo 6, and consider the subgroup

$$N = \langle 3\rangle = \{0, 3\}. \tag{23}$$

The distinct cosets are

$$0 + N = \{0, 3\}, \quad 1 + N = \{1, 4\}, \quad 2 + N = \{2, 5\}. \tag{24}$$

Thus,

$$|\mathbb{Z}_6/\langle 3\rangle| = 3, \tag{25}$$

and

$$\mathbb{Z}_6/\langle 3\rangle \cong \mathbb{Z}_3. \tag{26}$$

4.2 Order of a Factor Group

Let G be a finite group and let $N \trianglelefteq G$. Then the order of the factor group G/N is given by

$$|G/N| = \frac{|G|}{|N|}. \tag{27}$$

This follows from **Lagrange's theorem**, since the cosets of N in G partition G into disjoint subsets, each having $|N|$ elements.

4.3 Important Remarks on Factor Groups

Let G be a group and $N \leq G$.

1. **Normality is necessary.** The quotient group G/N is defined only if N is a normal subgroup of G, i.e.

$$gNg^{-1} = N \quad \text{for all } g \in G. \tag{28}$$

   If N is not normal, the product of cosets is not well-defined.

2. **Cosets form a partition.** The elements of G/N are the left cosets of N, and these cosets partition G into disjoint subsets of equal size (by **Lagrange's theorem**).

3. **Order formula.** If G is finite, then
$$|G/N| = \frac{|G|}{|N|}. \tag{29}$$

4. **Abelian property.** If G is abelian, then every subgroup $N \leq G$ is automatically normal. Hence quotients of abelian groups are always defined.

5. **Identity element.** The identity element of G/N is the coset N itself.

6. **Inverse of a coset.** For $gN \in G/N$,
$$(gN)^{-1} = g^{-1}N. \tag{30}$$

7. **Trivial and whole-group cases.**
$$G/\{e\} \cong G, \qquad G/G \cong \{e\}. \tag{31}$$

8. **Abelian quotient from commutator subgroup.** The quotient
$$G/[G,G] \tag{32}$$
is always abelian. Here $[G,G]$ denotes the commutator subgroup.

9. **Homomorphism connection (First Isomorphism Theorem).** If $\varphi : G \to H$ is a group homomorphism, then
$$G/\ker(\varphi) \cong \mathrm{Im}(\varphi). \tag{33}$$

   Define
$$\varphi : \mathbb{Z}_6 \to \mathbb{Z}_3, \qquad \varphi([x]_6) = [x]_3. \tag{34}$$

   Explicitly,

$\mathbb{Z}_6$	$\mathbb{Z}_3$
$0,3$	0
$1,4$	1
$2,5$	2

(35)

   Kernel.
$$\ker(\varphi) = \{0,3\} = \langle 3 \rangle. \tag{36}$$

   Coset Decomposition.
$$\mathbb{Z}_6 = \{0,3\} \cup \{1,4\} \cup \{2,5\}. \tag{37}$$

   These are precisely the cosets of $\ker(\varphi)$.

Quotient Group.

$$\mathbb{Z}_6/\langle 3\rangle = \{\{0,3\},\{1,4\},\{2,5\}\}. \tag{38}$$

Hence,

$$\mathbb{Z}_6/\langle 3\rangle \cong \mathbb{Z}_3. \tag{39}$$

$$\begin{tikzcd} \mathbb{Z}_6 \arrow[r, "\varphi"] \arrow[d, "\pi"'] & \mathbb{Z}_3 \\ \mathbb{Z}_6/\langle 3\rangle \arrow[ur, "\cong"'] & \end{tikzcd} \tag{40}$$

The map $\pi : G \longrightarrow G/N$ defined by

$$\pi(g) = gN \tag{41}$$

is called the natural (or canonical) projection. It is a surjective homomorphism with

$$\ker(\pi) = N. \tag{42}$$

Thus, π identifies elements of G that differ by an element of N, and every homomorphism $\varphi : G \longrightarrow H$ factors through this projection via the quotient $G/\ker(\varphi)$.

§5 Conjugacy Classes

For a fixed element a, A conjugacy class of an element a in a group G is defined as the set of all elements in G that are equal to gag^{-1} for some choice of g in G. Mathematically we can write,

$$C = \{gag^{-1} : g \in G\} \qquad C \Longrightarrow \text{conjugacy class of a} \tag{43}$$

> Each element in an abelian group forms its own distinct conjugacy class.

Conjugation is an **equivalence relation**, meaning it is reflexive (a is conjugate to itself), symmetric (if a is conjugate to b, b is conjugate to a), and transitive (if a is conjugate to b and b to c, then a is conjugate to c)

$$\text{Equivalence relation} \Longrightarrow \text{Mutually exclusive} \tag{44}$$

Example 5.1. Consider $S(3)$. Since $geg^{-1} = e$ for all $g \in S(3)$, the set $\{e\}$ forms a conjugacy class. Now $bab^{-1} = bab^2 = ab^4 = ab$, and $(b^2)a(b^2)^{-1} = b^2ab = b^4a = ba = ab^2$. Thus $\{a, ab, ab^2\}$ is a conjugacy class. Further, $aba^{-1} = aba = b^2a^2 = b^2$, and it follows that $\{b, b^2\}$ is a conjugacy class. Hence we obtain

$$\mathfrak{S}(3) = \{e\} \cup \{a, ab, ab^2\} \cup \{b, b^2\} \tag{45}$$

> In the symmetric group $\mathfrak{S}(n)$ elements are conjugate if and only if they have the same cycle structure. In general, conjugate elements shares same order.

§A Basic Notions from Ring Theory

A *ring* is a set R equipped with two binary operations, addition and multiplication, such that:

1. $(R, +)$ is an abelian group,
2. multiplication is associative,
3. multiplication distributes over addition, i.e.

$$a(b + c) = ab + ac, \qquad (a + b)c = ac + bc. \tag{46}$$

If there exists an element $1 \in R$ satisfying

$$1a = a1 = a \quad \text{for all } a \in R, \tag{47}$$

then R is called a ring with identity.

A.1 The Ring $\mathbb{Z}_n$

For a fixed integer $n \geq 1$, the set of residue classes modulo n,

$$\mathbb{Z}_n = \{[0], [1], \ldots, [n-1]\}, \tag{48}$$

forms a finite commutative ring under

$$[a] + [b] = [a + b], \qquad [a] \cdot [b] = [ab]. \tag{49}$$

A.2 Examples.

Example 1 In $\mathbb{Z}_5$,

$$[2] \cdot [3] = [6] = [1], \tag{50}$$

so every nonzero element has a multiplicative inverse; hence $\mathbb{Z}_5$ is a field.

Example 2 In $\mathbb{Z}_6$,

$$[2] \cdot [3] = [6] = [0], \tag{51}$$

so nonzero elements may multiply to zero. Such elements are called zero divisors.

> red The ring $\mathbb{Z}_n$ is a field if and only if n is prime.

A.3 Connection with Group Theory

The additive structure $(\mathbb{Z}_n, +)$ is a finite cyclic group of order n generated by $[1]$. Thus, if an element r in a group satisfies

$$r^n = e, \tag{52}$$

then

$$\langle r \rangle \cong (\mathbb{Z}_n, +), \tag{53}$$

and exponent arithmetic is naturally computed modulo n.

§B Modulo Arithmetic and the Structure of $\mathbb{Z}_n$

B.1 Congruence Modulo n

Let $n \in \mathbb{N}$ with $n \geq 1$. For $a, b \in \mathbb{Z}$, we define

$$a \equiv b \pmod{n} \tag{54}$$

if and only if $n \mid (a - b)$. Equivalently, $a \equiv b \pmod{n}$ if there exists $k \in \mathbb{Z}$ such that $a = b + kn$. This relation is an equivalence relation on $\mathbb{Z}$.

B.2 Residue Classes

For $a \in \mathbb{Z}$, the residue class of a modulo n is defined by

$$[a] = \{a + kn \mid k \in \mathbb{Z}\}. \tag{55}$$

The set of all residue classes modulo n is denoted by

$$\mathbb{Z}_n = \{[0], [1], \ldots, [n-1]\}. \tag{56}$$

Each integer belongs to exactly one residue class.

B.3 Arithmetic in $\mathbb{Z}_n$

Addition and multiplication in $\mathbb{Z}_n$ are defined by

$$[a] + [b] = [a + b], \tag{57}$$

$$[a] \cdot [b] = [ab]. \tag{58}$$

These operations are well-defined and make $\mathbb{Z}_n$ a commutative ring.

B.4 Examples

Example 1.

$$17 \equiv 5 \pmod{12} \tag{59}$$

since

$$17 - 5 = 12, \tag{60}$$

and 12 is divisible by 12.

Example 2. Arithmetic in $\mathbb{Z}_5$:

$$[3] + [4] = [7] = [2], \tag{61}$$

$$[3] \cdot [4] = [12] = [2]. \tag{62}$$

B.5 Algebraic Properties

If

$$a \equiv b \pmod{n} \quad \text{and} \quad c \equiv d \pmod{n}, \tag{63}$$

then

$$a + c \equiv b + d \pmod{n}, \tag{64}$$

$$ac \equiv bd \pmod{n}. \tag{65}$$

B.6 Connection with Cyclic Groups

The additive group $(\mathbb{Z}_n, +)$ is cyclic of order n, generated by $[1]$:

$$\mathbb{Z}_n = \langle [1] \rangle. \tag{66}$$

If r satisfies $r^n = e$, then

$$\langle r \rangle \cong (\mathbb{Z}_n, +), \tag{67}$$

since

$$r^a r^b = r^{a+b}, \tag{68}$$

with exponents computed modulo n.

B.7 Geometric Interpretation

Modulo arithmetic is cyclic in nature. For instance, in clock arithmetic (modulo 12),

$$10 + 5 \equiv 3 \pmod{12}. \tag{69}$$

Thus, two integers are congruent modulo n precisely when they differ by a multiple of n.

§C Algebraic Hierarchies

Algebraic structures are organised hierarchically according to the number of operations defined on a set and the axioms they satisfy. Each successive level in a hierarchy is obtained by imposing an additional structural property.

C.1 Structures with One Operation

Beginning with a set equipped with a single associative operation, we obtain:

- A **semigroup**: associativity holds.
- A **monoid**: a semigroup with an identity element.
- A **group**: a monoid in which every element has an inverse.

Thus, each step adds exactly one new requirement:

$$\text{Semigroup} \implies \text{Monoid} \implies \text{Group}. \tag{70}$$

C.2 Structures with Two Operations

When two operations, addition and multiplication, are introduced, the hierarchy becomes:

- A **ring**: $(R, +)$ is an abelian group, multiplication is associative, and distributive laws hold.
- An **integral domain**: a commutative ring with identity and no zero divisors.
- A **field**: an integral domain in which every nonzero element has a multiplicative inverse.

Symbolically,

$$\text{Ring} \implies \text{Integral Domain} \implies \text{Field}. \tag{71}$$

Each refinement strengthens the algebraic structure by improving the behaviour of multiplication.

C.3 Conceptual Perspective

The hierarchy reflects increasing structural control:

- Identity provides stability.
- Inverses provide reversibility.
- Absence of zero divisors ensures cancellation.
- Multiplicative inverses allow division.

Understanding these layers clarifies the role of structures such as cyclic groups and the ring $\mathbb{Z}_n$ in group-theoretic contexts, where often only part of the available algebraic structure is required.

§D Special Other Groups

D.1 The Quaternion Group Q

The *Quaternion Group* Q is a non-abelian group of order 8. It provides the group structure of the unit quaternions and plays an important role in both algebra and physics.

Definition

The group is defined as

$$Q = \{e, s, i, j, k, si, sj, sk\}, \tag{72}$$

where e is the identity and $s = -e$. i, j, k are imaginary units, and s is the negative of the identity (often represented as -1)

Relations

The elements satisfy the fundamental identities

$$s^2 = e, \tag{73}$$

$$i^2 = j^2 = k^2 = ijk = s. \tag{74}$$

The cyclic multiplication rules are

$$ij = k, \qquad jk = i, \qquad ki = j. \tag{75}$$

The group is non-abelian. For example,

$$ij = sji \quad (\text{equivalently } ij = -ji). \tag{76}$$

The elements e and s commute with all elements of Q.

Structure of Subgroups

A remarkable property of Q is that *every subgroup is normal*. The non-trivial proper subgroups are $\{e, s\}$, and the three subgroups of order 4 $\{e, s, i, si\}, \{e, s, j, sj\}, \{e, s, k, sk\}$.

Element Orders

The distribution of element orders in Q is:

$$|Q| = 8, \tag{77}$$

$$\text{ord}(s) = 2, \tag{78}$$

$$\text{ord}(i) = \text{ord}(j) = \text{ord}(k) = \text{ord}(si) = \text{ord}(sj) = \text{ord}(sk) = 4. \tag{79}$$

Comparison with the Dihedral Group D_4

Q and D_4 are the only non-abelian groups of order 8. However, they are not isomorphic because their element order distributions differ. Though, they share the same number of conjugacy classes (five) and even the same character table, they are not isomorphic.

Conjugacy Classes

The conjugacy class decomposition of Q is

$$Q = \{e\} \cup \{s\} \cup \{i, si\} \cup \{j, sj\} \cup \{k, sk\}. \tag{80}$$

Since e and s commute with all elements, they each form their own conjugacy class.

Representation via Pauli Matrices

A faithful matrix representation is given by

$$i \mapsto -i\sigma_x, \qquad j \mapsto -i\sigma_y, \qquad k \mapsto -i\sigma_z. \tag{81}$$

Connection to Rotations

The algebra of unit quaternions is closely related to the rotation group $SO(3)$. Unit quaternions provide an efficient description of rotations in three-dimensional space.

Pseudoreal Representations

The two-dimensional irreducible representation of Q is **pseudoreal** (or quaternionic). This means the representation is equivalent to its complex conjugate but cannot be brought into a real form by a similarity transformation

One unique shared property is that the two-dimensional representation of the Quaternion group and the fundamental representation of $SU(2)$ are both pseudoreal (or quaternionic). In the context of $SU(2)$, this is demonstrated by the fact that the Pauli matrices are related to their complex conjugates through the antisymmetric matrix σ_y (often denoted as $i\sigma_2$)

Role in SU(2) and Double Covering

The Quaternion group Q is a finite subgroup of $SU(2)$ (the special unitary group in two dimensions). While $SU(2)$ is a continuous Lie group generated by the Pauli matrices, Q forms a discrete subset consisting of the eight elements $\{\pm I, \pm i\sigma_x, \pm i\sigma_y, \pm i\sigma_z\}$. Thus Q can be realised as a subgroup of $SU(2)$ made up of these specific unitary matrices. The group $SU(2)$ is the double cover of the rotation group $SO(3)$, meaning there exists a 2-to-1 homomorphism $SU(2) \to SO(3)$.

As a consequence, a full rotation of 2π in $SO(3)$ corresponds not to the identity in $SU(2)$, but to the element -1. In quaternion notation this element is s.

This explains why spin-$\frac{1}{2}$ particles acquire a minus sign under a 2π rotation. The quaternion group captures this sign structure algebraically and therefore plays an important role in the representation theory of rotations and in quantum mechanics.

D.2 Dihedral Group

The Dihedral Group (D_n) is the group of symmetries of a regular n-sided polygon in two-dimensional space. It consists of both rotations and reflections, making it a fundamental example in geometric and physical symmetry studies.
Generators : Denoted by r (rotation by $2\pi/n$) and s or σ (reflection).
Relations: $r^n = 1$, $s^2 = 1$, $srs = r^{-1}$.

Conjugacy Classes of D_n

Case 1: $n = 2p + 1$ (Odd)
In this case there are $p + 2$ conjugacy classes. The decomposition of D_{2p+1} into classes is

$$D_{2p+1} = \{e\} \cup \underbrace{\{r, r^{2p}\} \cup \{r^2, r^{2p-1}\} \cup \cdots \cup \{r^p, r^{p+1}\}}_{p \text{ classes}} \cup \{s, sr, sr^2, \ldots, sr^{2p}\}. \tag{82}$$

Here:

- $\{e\}$ forms one class,
- the rotations form p classes,
- all reflections form one single class.

Thus the total number of conjugacy classes is

$$1 + p + 1 = p + 2. \tag{83}$$

Case 2: $n = 2p$ (Even)
In this case there are $p + 3$ conjugacy classes. The decomposition of D_{2p} is

$$D_{2p} = \{e\} \cup \{r^p\} \cup \underbrace{\{r, r^{2p-1}\} \cup \{r^2, r^{2p-2}\} \cup \cdots \cup \{r^{p-1}, r^{p+1}\}}_{p-1 \text{ classes}} \cup \{s, sr^2, \ldots, sr^{2p-2}\} \cup \{sr, sr^3, \ldots, sr^{2p-1}\}. \tag{84}$$

Here:

- $\{e\}$ and $\{r^p\}$ each form a separate class,
- the remaining rotations form $p - 1$ classes as $r^p = r^{-p}$,
- reflections split into two distinct classes.

Thus the total number of conjugacy classes is

$$(p + 1) + 2 = p + 3. \tag{85}$$

Example of D_3 (Symmetries of an Equilateral Triangle, Order 6)

- **Subgroups:**
  - Trivial: $\{e\}$ and D_3.
  - Rotation subgroup: $\{e, r, r^2\}$ (Order 3).
  - Reflection subgroups: $\{e, s\}$, $\{e, sr\}$, $\{e, sr^2\}$ (Each of Order 2).
- **Normal Subgroups:**
  - The trivial subgroups $\{e\}$ and D_3 itself.
  - The cyclic subgroup of rotations $\{e, r, r^2\}$ is the only non-trivial normal subgroup.

- The reflection subgroups are *not normal* because conjugating one reflection by a rotation gives a different reflection.

- **Conjugacy Classes:** D_3 has **three classes**:
  - $\{e\}$
  - The rotations: $\{r, r^2\}$
  - The reflections: $\{s, sr, sr^2\}$

Example: D_4 **(Symmetries of a Square, Order** 8**)**

- **Subgroups:** D_4 has several subgroups, including the rotation group C_4 and various reflection subgroups.

- **Normal Subgroups:** Unlike Q_8, not all subgroups of D_4 are normal. The normal subgroups include the rotation group $\{e, r, r^2, r^3\}$ and the center $\{e, r^2\}$.

- **Conjugacy Classes:** D_4 has **five classes**:
  - $\{e\}$
  - $\{r^2\}$ (The 180° rotation)
  - $\{r, r^3\}$ (The 90° and 270° rotations)
  - $\{\sigma, \sigma r^2\}$ (Reflections across axes through midpoints of sides)
  - $\{\sigma r, \sigma r^3\}$ (Reflections across diagonal axes through vertices)