



**Gramin Krishi Mausam Sewa**  
**Agromet Advisory Bulletin for DIBRUGARH District**  
**Assam Agricultural University**  
 Jorhat, Assam



## **Agromet Advisory Bulletin**

Date : 2026-09-11

Weather Forecast of District DIBRUGARH (Assam) Issued On : 2026-09-11 (Valid Till 08:30 IST of the next 5 days)

| Parameter              | 2026-09-12                            | 2026-09-13                                        | 2026-09-14                                        | 2026-09-15 | 2026-09-16 |
|------------------------|---------------------------------------|---------------------------------------------------|---------------------------------------------------|------------|------------|
| Rainfall(mm)           | 15.3                                  | 9.9                                               | 7.4                                               | 4.0        | 9.0        |
| Tmax(°C)               | 36.7                                  | 36.3                                              | 35.9                                              | 35.7       | 33.9       |
| Tmin(°C)               | 24.4                                  | 24.1                                              | 24.1                                              | 24.4       | 23.8       |
| RH-I(%)                | 87                                    | 84                                                | 83                                                | 82         | 84         |
| RH-II(%)               | 82                                    | 79                                                | 78                                                | 76         | 78         |
| Wind Speed(kmph)       | 2                                     | 2                                                 | 1                                                 | 1          | 3          |
| Wind Direction(Degree) | 59                                    | 68                                                | 90                                                | 56         | 27         |
| Cloud Cover(Octa)      | 5                                     | 4                                                 | 3                                                 | 4          | 5          |
| Warning                | Thunderstorms & Lightning, Squall etc | Heavy Rain; Thunderstorms & Lightning, Squall etc | Heavy Rain; Thunderstorms & Lightning, Squall etc | No Warning | No Warning |

### Forecast Summary:

The weather will remain partly cloudy with the possibilities of getting light rainfall in the coming 5 days. Daily maximum and minimum temperature for the ensuing 5 days is predicted to be around 35.7°C and 24.2°C respectively. Morning and afternoon relative humidity will be around 84% and 79% respectively. Wind will blow mainly from northeasterly direction at an average speed of around 2.0 kmph. Extended Range Weather Forecast: During the period from 13th September, 2026 to 19th September, 2026, the rainfall amount, maximum and minimum temperature will remain normal over Assam-Meghalaya Meteorological Subdivision.

### Weather Warnings (Valid Till 08:30 IST of the next day)

There is a probability of occurrence of heavy rain, thunderstorm and lightning in some isolated areas from 12th to 14th September, 2026

### Likely impacts of weather warnings on Agriculture and associated Agromet advisories

Provide mechanical support to the crops like banana, citrus, papaya etc. and make arrangements to drain out excess water from the fields. Harvest the matured vegetable and fruits and transfer it to a safer location. Transfer the seedlings of fruit crops, flower crops, vegetable crops etc. to a safer location. In case if the seedling cannot be transferred, make provisions for firm temporary sheds (made of bamboo and net or plastic).

### General Advisory:

Taking advantage of the suitable weather condition in the coming 5 days, farmers can continue their farm operations like, land preparation, sowing, transplanting, earthing up, application of fertilizers, spraying of pesticides, etc.

### SMS Advisory:

Burning of dried leaves mixed with dried chilli near the vegetable orchards will control the infestation of borers and suckers up to certain extent.

### Crop Specific Advisory:

| Crop (Stage) | Crop Specific Advisory                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |
|--------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| RICE         | <ul style="list-style-type: none"> <li>• Since there is possibility of getting light rainfall in the coming 5 days, repair the bunds in the paddy fields to conserve rain water. Maintain standing water of 5±2 cm depth in the field.</li> <li>• Considering suitable weather condition in the coming 5 days, go for second top dressing with Urea @ 4.5 kg per bigha in the medium and long duration semi dwarf varieties and 1.5 kg per bigha in medium and long duration tall varieties of sali rice which are at panicle initiation stage. In case of short duration (about 100 days) semi dwarf varieties the dose of urea for top dressing should be 9 kg per bigha.</li> <li>• Since the weather condition has become favourable for the attack of swarming caterpillar in rice crop, farmers can go for traditionally controlling the pest by sweeping coconut ropes, dipped in kerosene oil across the rice crop, causing the caterpillars to fall into water and die, or place bird perches in the field. Periodically release ducks into the fields, which will in turn consume the larvae and reduce the pest infestation. To stop the movement of swarming caterpillars from one field to another, spray the bunds with Fipronil 5% SC @ 1ml/lit of water. In infested crop, spray Lambda-cyhalothrin 5 EC @ 1.5 ml/lit of water or Chlorantraniliprole 18.50% SC @ 1ml per 3 liters of water at 7 days interval during evening hours. Three sprayings are necessary to control the pest.</li> </ul> |

### Horticulture Specific Advisory:

| Horticulture (Stage) | Horticulture Specific Advisory                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
|----------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| CAULIFLOWER          | <ul style="list-style-type: none"> <li>• Continue nursery sowing of mid-season varieties of Cauliflower and Broccoli under shed-nets or poly houses. For sowing, collect recommended varieties of Cauliflower like Improved Japanese, Pusa Synthetic, Pusa Snowball and Main Crop Patna and Broccoli like Green Sprouting Medium, Corsair, Cruiser, Emerald, Corona etc.</li> <li>• Prior to sowing, treat the seeds with fungicides like Carboxin @ 2g/kg seeds.</li> </ul>                                                                                   |
| RADISH               | <ul style="list-style-type: none"> <li>• Continue sowing of seeds of early radish varieties in the main field. The recommended early varieties are Pusa Deshi, Pusa Chetki, Pusa Himani etc. The seeds of Radish can be directly sown either on ridges spaced at distance of 45 cm or in flat beds with a row spacing of 20cm and a recommended plant to plant spacing of 10 cm. During the time of final land preparation 26 quintals of FYM along with 7.35 kg Urea, 42 kg SSP and 22.7 kg MOP per bigha is required to be applied in the fields.</li> </ul> |

### Poultry Specific Advisory:

| Poultry | Poultry Specific Advisory                                                                                                                                                                                                                                                                                                                                                                                                                 |
|---------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| CHICKEN | <ul style="list-style-type: none"> <li>• Increase the number of feeders and drinkers during feeding time to reduce competition among birds. Adding antioxidants such as vitamin E, vitamin C, selenium is shown to be helpful to reduce stress and improve feed consumption while maintaining or improving body weight gain. A high energy diet should be provided during summer because birds lose more energy while panting.</li> </ul> |

### Sericulture Specific Advisory:

| Sericulture | Sericulture Specific Advisory                                                                                                                                                                                                                 |
|-------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Muga        | <ul style="list-style-type: none"> <li>• Prepare the 'jali' (mountage) for 'Bhodia' muga crop before 3-4 days of ripening. Do not prepare the jali with wet leaves. Hang the jali in a well aerated, semi dark and rat proof room.</li> </ul> |

### Likely impacts of weather warnings (General)

☐ Visibility may become poor due to intense spell of rainfall. ☐ Temporary disruption of traffic due to water logging in roads/ uprooting of trees. ☐ Damage to kutchra roads ☐ Possibility of damages to vulnerable structures due to heavy to very heavy rain ☐ Water logging in low lying areas. ☐ Landslides/land slip in some locations likely. ☐ Heavy rainfall may damage the standing crops and vegetables in the maturity stage. ☐ Lightning may injure people and cattle at open place. ☐ Dispersion of soil from the field and hence seed displacement and poor germination of seeds.

### Impact based advisories (General)

Follow any traffic advisories. • Avoid going to areas that face water logging problem often. • Avoid staying in landslide prone areas. • Take shelter during thunderstorm/lightning activities. • Provide mulch at the base of the crop to prevent soil and root damage. • Avoid working in the fields during thunderstorm/lightning period. • Postpone sowing of rice, jute, maize and vegetables; if already sown, avoid water stagnation in the field and cover the seeded area with natural mulching materials like straw, farm residues etc. • Be updated.

**Farmers are advised to download Unified "Mausam" and "Meghdoot" android application on mobile for Weather forecast and weather based Agromet Advisories and "Damini" android application for forecast of Thunderstorm and lightening.**

**Mausam MobileApp link: <https://play.google.com/store/apps/details/>**

**Meghdoot MobileApp link: <https://play.google.com/store/apps/details>**

**Damini MobileApp link : <https://play.google.com/store/apps/details>**