



Gramin Krishi Mausam Sewa
Agromet Advisory Bulletin for SURAT District
Navsari Agricultural University
Navsari, Gujarat



Agromet Advisory Bulletin

Date : 2026-09-11

Weather Forecast of District SURAT (Gujarat) Issued On : 2026-09-11 (Valid Till 08:30 IST of the next 5 days)

Parameter	2026-09-12	2026-09-13	2026-09-14	2026-09-15	2026-09-16
Rainfall(mm)	15.0	68.0	125.0	66.0	30.0
Tmax(°C)	35.0	34.0	34.0	31.0	32.0
Tmin(°C)	26.0	26.0	24.0	24.0	25.0
RH-I(%)	75	75	80	85	85
RH-II(%)	70	70	75	75	85
Wind Speed(kmph)	6	11	12	15	12
Wind Direction(Degree)	243	233	254	230	221
Cloud Cover(Octa)	7	8	8	7	5
Warning	No Warning	No Warning	Heavy Rain; Thunderstorms & Lightning, Squall etc	Heavy Rain; Thunderstorms & Lightning, Squall etc	Heavy Rain; Thunderstorms & Lightning, Squall etc

Forecast Summary:

Moderate to Very Heavy rainfall (totaling around 304 mm) and generally cloudy to cloudy skies are expected over the next few days. Temperatures will range from a maximum of 31-35°C to a minimum of 24-26°C. The maximum relative humidity reaching 75-85% and a minimum of 70-85%. Additionally, south-westerly surface winds blowing at average speeds will be around 6–15 kmph.

Weather Warnings (Valid Till 08:30 IST of the next day)

Yellow Alert for Heavy Rain, Thunderstorms & Lightning, Squall etc. has been issued between September 14 to 16, 2026.

Likely impacts of weather warnings on Agriculture and associated Agromet advisories

Likely Impacts on Agriculture Crop Lodging: Squalls and gusty winds can cause the lodging (bending or uprooting) of tall standing crops such as sugarcane, cotton, and fruit crops like banana. **Chemical Wash-off:** The moderate to heavy rainfall associated with the thunderstorm wash away freshly applied fertilizers and pesticide sprays. **Livestock & Human Safety:** Lightning strikes pose a direct and fatal threat to farm laborers working in open fields and livestock grazing outdoors. **Associated Agromet Advisories Postpone Field Operations:** Strictly postpone all scheduled spraying of pesticides, herbicides, and fertilizer top-dressing to prevent them from washing away in the rain. **Crop Support:** Provide immediate physical support (propping, staking, or tying) to tall and heavily fruiting crops like sugarcane and banana to protect them from squall winds. **Safety Measures:** Farmers and farm laborers must immediately stop fieldwork and seek safe, indoor shelter during thunderstorm and lig

General Advisory:

• Irrigation: With heavy rainfall forecasted over the next 5 days, strictly withhold all irrigation to Kharif crops. • Drainage Management: Immediate priority must be given to field drainage. Deepen and clear all drainage channels (trenches/furrows) to prevent severe waterlogging, which will rapidly cause root anoxia and wilt in standing crops. • Spraying & Fertilizers: Strictly suspend all scheduled pesticide, herbicide, and foliar fertilizer sprays. Applying them now will result in 100% wash-off and economic loss. Wait for clear weather. • Pest/Disease Watch: Continuous dense cloud cover and 95% relative humidity create a highly favorable environment for soil-borne fungal diseases (wilt, root rot) and blight. Monitor crops closely once the rains subside. • Weed/Intercultural Operations: Postpone all manual weeding, hoeing, and intercultural operations. Entering waterlogged fields will severely damage soil structure. Wait until the soil returns to 'Vapsa' (workable moisture) condition.

SMS Advisory:

Heavy rain & thunderstorms are expected. Do NOT irrigate Kharif crops. Stop spraying and ensure clear drainage.

Crop Specific Advisory:

Crop (Stage)	Crop Specific Advisory
SUGARCANE (Grand growth)	• Propping: Prevent lodging by tying tall canes together with dry leaves or using bamboo stakes. • Detrashing: Strip lower dry leaves to improve aeration; reuse them as mulch or for tying canes. • Disease Management: Uproot and burn plants with smut, wilt, or Red Rot. Drench Red Rot-affected soil with Carbendazim 50% WP (2 g/L) during clear weather. • Pest Management: Stop granular shoot borer treatments and manually destroy dried shoots. Spray Azadirachtin 10000 ppm (20 ml/10 L) or 0.5% Neem oil for whitefly/Pyrrilla during clear weather.
COTTON (Flowering)	• Fertilizers: Avoid fertilizer application during long dry spells, applying only under optimum moisture; spray 2% urea during dry spells. • Pheromone Traps: Install traps in the field to monitor and control pests effectively. • Sucking Pests: Spray 500 g neem seed extract or 50 ml neem oil per 10 L of water for initial infestations. For heavy infestations, spray 3 g Flonicamid 50 WG or 10 g Diafenthiuron 50 WP per 10 L of water strictly during clear weather. • Disease Control (Root Rot/Wilt): Drench infected plants with Mancozeb 75 WP (27 g), Copper Oxychloride (40 g), or Carbendazim (10 g) mixed in 10 L of water.
RICE (Tillering)	• Water Management: Maintain 2–3 cm of standing water and secure field bunds to prevent nutrient runoff during rains. • Weed Control: Weed manually or use a cono-weeder to aerate roots; spray Bispyribac-sodium strictly during clear weather. • Blast Disease: Spray Tricyclazole 75 WP (6 g) or Iprobenfos 48 EC (10 ml) per 10 L of water every 15–20 days during clear weather. • Bacterial Leaf Blight: Spray 1 g Streptocycline + 20 g Copper Oxychloride in 400–500 L of water per hectare during rain-free periods.
PIGEON PEA (RED GRAM/ARHAR) (Vegetative)	• Undertake interculturaling, and hand weeding operations at 20, 40 and 60 days after sowing under vapsa condition in sown crops.
CASTOR (Vegetative)	• Undertake the gap filling if required in sown maize crop. • Apply need based light irrigation by ensuring soil moisture in prevailing dry spell condition in sown castor crop.
SOYABEAN (Vegetative)	• Undertake the interculturaling and weeding operations at vapsa condition at 20 and 40 days after sowing in sown crop. • Undertake the interculturaling and weeding operations at 30 days after sowing at vapsa condition even if pre-emergence or post-emergence herbicides have been sprayed for effective and economical weed control in the soybean crop. • Use one light trap per hectare to attract and destroy the moths of armyworm. If a severe infestation of armyworm is observed, spray a mixture of Emamectin Benzoate 1.9 EC (8.5 ml) or Indoxacarb 14.5 SC (7 ml) per 10 liters of water during clear weather.
	• Fertilizer: Apply the top-dressing of Nitrogen (Urea) when the crop is 30–

SORGHUM (JOWAR/GREAT MILLET) (Vegetative)	40 days old, ensuring the wind is calm. • Earthing-Up: Mound soil around the plant stems to provide root support and prevent lodging from strong winds. • Pest Control: Install fishmeal traps in the field to attract and control shoot flies. Spray recommended insecticides for stem borers only during clear weather.
--	---

Horticulture Specific Advisory:

Horticulture (Stage)	Horticulture Specific Advisory
MANGO	• Young Plant Support: Stake young plants firmly to protect them from upcoming heavy rain and strong winds. • Stem Borer Control: Kill stem grubs and plug their holes with mud and Chlorpyrifos 20 EC (25 ml/10 L of water) before rains begin. • Disease Prevention: Apply 10% Bordeaux paste to pruned or cut branches during clear weather to stop fungal infections. • New Plantations: Prepare pits using healthy saplings, 10 kg FYM, 1 kg SSP, 1 kg MOP, and 50 ml Chlorpyrifos per tree for termite management. • Fertilizers Postponed: Strictly suspend all chemical (Urea, SSP, MOP) and biofertilizers to prevent severe nutrient leaching during the rain spell.
BANANA	• Weed Management & Interculturing: Perform interculturing and manual weeding in the planted field strictly when the soil reaches the 'Vapsa' (workable moisture) condition to keep the field weed-free and aerate the roots. • Irrigation Management: Since mostly dry weather is forecasted, monitor the soil moisture closely and apply light irrigation to the newly planted crop as needed.

Live Stock Specific Advisory:

Live Stock	Live Stock Specific Advisory
COW	• Housing & Hygiene: Keep animals in clean, dry sheds. Regularly sprinkle dry slaked lime (Chuna) on the floor to maintain hygiene and prevent diseases like Foot Rot and Mastitis. • Fodder & Nutrition: Strictly avoid feeding old damp or fungus-infested fodder from past rains. Ensure a daily dose of mineral mixture and clean drinking water. • Healthcare: Administer deworming medicines as advised by a veterinarian and complete any pending vaccinations (FMD, HS, and BQ). • Pest Control: Clear any old stagnant water around the shed. Burn dry neem leaves in the evening to control mosquitoes, flies, and ticks.

Poultry Specific Advisory:

Poultry	Poultry Specific Advisory
CHICKEN	• Shed Care: Ensure ventilation; hang curtains against high winds but keep them partially open during the day. • Litter: Stir bedding regularly to prevent ammonia, adding slaked lime powder only on damp spots. • Feed: Store feed bags on elevated wooden pallets to prevent toxic fungal growth. • Health: Provide chlorinated or sanitized water, maintain vaccinations, and consult a vet if birds seem sick.

Likely impacts of weather warnings (General)

Nil

Impact based advisories (General)

Nil

Farmers are advised to download Unified “Mausam” and "Meghdoot" android application on mobile for Weather forecast and weather based Agromet Advisories and "Damini" android application for forecast of Thunderstorm and lightening.

Mausam MobileApp link: <https://play.google.com/store/apps/details/>

Meghdoot MobileApp link: <https://play.google.com/store/apps/details>

Damini MobileApp link : <https://play.google.com/store/apps/details>
