




Agromet Advisory Service Bulletin for the State of Gujarat

(Advisories mentioned in the bulletin are prepared by different AMFUs and DAMUs in Gujarat)

Bulletin No. 70/2026

Issued on 09-09-2026

Agro-Climatic zones of Gujarat



Weather warnings (Met-Subdivision Level)

DAY-1	NIL
DAY-2	NIL
DAY-3	NIL
DAY-4	Heavy rains at isolated places very likely in all the districts of South Gujarat region namely Vadodara, Chhota Udepur, Narmada, Bharuch, Surat, Tapi, Dang, Navsari, Valsad and in Daman, Dadra and Nagar Haveli. Light thunderstorm with lightning and surface wind 30-40 kmph accompanied with light to moderate rain very likely at isolated places in the districts of North Gujarat region namely Mehsana, Sabarkantha, Gandhinagar, Aravalli, Kheda, Ahmedabad, Anand, Panchmahal, Dahod and Mahisagar; in all the districts of South Gujarat region namely Vadodara, Chhota Udepur, Narmada, Bharuch, Surat, Tapi, Dang, Navsari, Valsad and in Daman, Dadra and Nagar Haveli; in the districts of Saurashtra namely Surendranagar, Amreli, Bhavnagar and Botad.

DAY-5 **Heavy rains** at isolated places very likely in the districts of in the districts of Gujarat region namely Ahmedabad, Anand, Panchmahal, Dahod, Vadodara, Chhota Udepur, Narmada, Bharuch, Surat, Tapi, Dang, Navsari, Valsad and in Daman, Dadra and Nagar Haveli; in the districts of Saurashtra namely Rajkot, Porbandar, Junagadh, Amreli, Bhavnagar, Gir Somnath, Botad and in Diu.

Light thunderstorm with lightning and surface wind 30-40 kmph accompanied with light to moderate rain very likely at isolated places in all the districts of North Gujarat region namely Banaskantha, Vav-Tharad, Patan, Mehsana, Sabarkantha, Gandhinagar, Aravalli, Kheda, Ahmedabad, Anand, Panchmahal, Dahod and Mahisagar; in all the districts of South Gujarat region namely Vadodara, Chhota Udepur, Narmada, Bharuch, Surat, Tapi, Dang, Navsari, Valsad and in Daman, Dadra and Nagar Haveli; in all the districts of Saurashtra-Kutch namely Surendranagar, Rajkot, Jamnagar, Porbandar, Junagadh, Amreli, Bhavnagar, Morbi, Dwarka, Gir Somnath, Botad, Kutch and in Diu.

Extended Range forecast at Met-Subdivision Level (09th Sept to 13th Sept 2026)

Meteorological Sub-Division	Rainfall	Max Temp
Gujarat region	Normal	Normal
Saurashtra-Kutch	Normal	Normal

Summary of four days rainfall over the State (05 - 08 Sept 2026) :

Met Sub Division	05-09-2026	06-09-2026	07-09-2026	08-09-2026
Gujarat region	ISOL	SCT / DRY	ISOL	DRY / SCT
Saurashtra-Kutch	ISOL / DRY	ISOL / DRY	ISOL / DRY	DRY / ISOL

Rainfall forecast for next 5 days

Met Sub Division	09-09-2026	10-09-2026	11-09-2026	12-09-2026	13-09-2026
Gujarat region	ISOL / SCT	ISOL / SCT	ISOL / SCT	FWS	FWS
Saurashtra-Kutch	ISOL	ISOL	ISOL	FWS / ISOL	FWS

ISOL: Isolated i.e 1 or 2 places

FWS: Fairly wide spread i.e Rain at many places

WS: Widespread, i.e Rain at most places.

SCT: Scattered i.e Rain at a few places

Dry: No rain

Agromet Advisories issued by different Agromet Field Units (AMFUs) of Gujarat

Gujarat region: Met Subdivision

AMFU Navsari (Dangs, Navsari, Valsad districts):

Navsari:

Crop	Weather based Agro-Advisory
Sugarcane	Vegetative Stage/Grand Growth stage: ➤ After 6, 7 & 8 month of planting remove dry leaves and use it as mulch. ➤ Up root and burn whip smut and wilted plant from field. ➤ Drenching of <i>Carbendazim</i> 50 wp 10gm/10 lit water from prevent spread of both disease (Sugarcane red rot & Wilt) nearby healthy plants during clear weather condition. ➤ Prop the sugarcane canes with bamboos or tie the canes together in a clump with sugarcane leaves to protect against wind and heavy rain. ➤ Putting sugarcane leaves having whitefly puparia in 30 mesh wire cage. The adult whitefly emerge from puparia remain inside cage and parasitoid will come out and parasitize other whiteflies. ➤ Spraying <i>Azadirachtin</i> 10000 ppm @ 20 ml/10 lit. of water for management of sugarcane whitefly. Or ➤ Application of neem oil 0.5% (50 ml + 10 g detergent powder/10 litre water) first when initiation of sugarcane whitefly and second spray at 10 days after first spray for effective management of sugarcane whitefly under clear weather condition.
Kharif Paddy	Flowering stage/Reproductive stage: ➤ Keeps 2-3 cm standing water in field during vegetative/Reproductive growth. ➤ Apply split dose of Nitrogen fertilizer as per recommendation. i.e. 2nd dose @ 20 kg N/ha at 40-45 days after transplanting. ➤ Keep field weed free by hand weeding and interculturing with paddy weeder or by application of recommended post-emergence weedicide upto 45 days after transplanting. ➤ After transplanting in paddy crop, monitor the crop condition and depending on rainfall, spraying of <i>Flubendiamide</i> 20 WG @ 2.5 g or <i>Chlorantraniliprole</i> 18.5 SC @ 3 ml or <i>Cartap hydrochloride</i> 75 SG @ 10 g per 10 litres of water during clear weather conditions for management of stem borer.
Pigeon Pea	➤ Weeding and intercultural operations should be carried out during clear weather condition.
Okra	➤ Special care should be taken that enough water should be supplied during the period of harvesting. ➤ Up root and burn yellow vein mosaic infected okra plant from field during clear weather condition. ➤ Spraying <i>Cyantraniliprole</i> 10.26 OD @ 18 ml or <i>Difenthiuron</i> 50 WP @ 12 g/10 Lit. for whitefly (vector) management during clear weather condition. ➤ Spraying of <i>Propargite</i> 57 EC @ 20 ml/10 Lit. or <i>Fenazaquin</i> 10 EC @ 10 ml./10 Lit. of water for red Spider Mite during clear weather condition.
Brinjal	➤ Transplanting should be carried out when the seedlings are 28-30 days old.
Tomato	➤ Transplanting should be carried out when the seedlings are 25-30 days old.
Chilly	➤ Seedlings of chilli become suitable for transplanting in about 35 to 40 days.
Pointed gourd	➤ Harvesting should be carried out as per maturity.
Little gourd	➤ Harvesting should be carried out as per maturity.
Elephant	➤ Apply fertilizer 50: 00: 37.5 NPK kg/ha. (109 kg Urea and 63 kg MOP per hectare).

foot yam	➤ Take measures for weed control.
Greater yam	➤ Take measures for weed control. ➤ Give proper support or train branches of vines. ➤ Apply fertilizer 40:00:40 NPK kg/ha. (87 kg Urea and 67 kg MOP per hectare).
Tannia	➤ Take measures for weed control. ➤ Harvest the suitable tender leaves of Tannia. ➤ Apply fertilizer 40:00:40 NPK kg/ha. (87 kg Urea and 67 kg MOP per hectare).
Mango	➤ Find grubs of stem borer from gallery and holes on the stem and kill them. Do pouring of <i>Chlorpyrifos</i> 20 EC @ 25 ml/ 10 lit in hole and plug it with mud in clear weather condition. ➤ Procure healthy and disease free planting material from authorized and government nursery for the new plantation. ➤ Application of 10 % Bordeaux paste in cut/pruning branches of fruit tree during clear weather condition. ➤ Prepare pit for plantation of new grafts by filling 10 kg FYM, 1 kg SSP and 1 kg MOP. Give 50 ml/tree <i>Chlorpyrifos</i> to manage termite and ants during clear weather condition. ➤ Provide support to the young plants of mango plants to protect against rain and high wind speed. ➤ Apply recommended dose of fertilizer per tree @ 800 g Urea or 1.8 kg Ammonium Sulphate, 1.0 kg SSP and 1.3 kg MOP along with 100 kg FYM to adult tree. ➤ In addition RDF, give biofertilizers viz., <i>Azotobactor</i> 50 ml, PSB 50 ml, KMB 50 ml and <i>Trichoderma</i> 100 g per adult tree in monsoon. Before or after 15 days of fertilizer application.
Sapota	➤ Provide support to the young plants of Sapota plants to protect against rain and high wind speed. ➤ Procure healthy and disease free planting material from authorized and government nursery for the new plantation. ➤ Prepare pit for plantation of new grafts by filling 10 kg FYM, 1 kg SSP and 1 kg MOP. ➤ Give 50 ml/tree <i>Chlorpyrifos</i> to manage termite and ants during clear weather condition. ➤ Apply second split of 25% of RDF (Ammonium Sulphate -1.3 kg or Urea 0.6 kg, SSP 0.8 kg and MOP 0.3 kg + FYM 25 kg each tree) in clear weather condition.

Dangs:

Crop	Weather based Agro-Advisory
Kharif Paddy	Flowering Stage : ➤ Apply Ghanjivamrut fertilizer in standing crop. ➤ Spray of Jivamrut @ 3 lit./10 lit. of water for better growth. ➤ Keep field weed free by hand weeding and interculturing with paddy weeder upto 45 days after transplanting. ➤ Spray the plantbase botanical pesticides like, Agniastra or Brahmastra @ 0.5 Lit. per 10 lit. of water.
Finger Millet	Flowering Stage : ➤ Apply Ghanjivamrut fertilizer in standing crop. ➤ Spray of Jivamrut @ 3 lit./10 lit. of water for better growth. ➤ Keep field weed free by hand weeding and interculturing with Hand weeder upto 45 days after transplanting. ➤ Spray the plantbase botanical pesticides like, Agniastra or Brahmastra @ 0.5 Lit. per 10 lit. of water.
Pulses	Flowering Stage : ➤ Apply Ghanjivamrut fertilizer in standing crop, if required. ➤ Spray of Jivamrut @ 3 lit./10 lit. of water for better growth.

	➤ Keep field weed free by hand weeding and interculturing with Hand weeder upto 45 days after sowing. ➤ Spray the plantbase botanical pesticides like, Agniastra or Brahmastra @ 0.5 Lit. per 10 lit. of water.
Little Millet	Flowering Stage : ➤ Apply Ghanjivamrut fertilizer in standing crop, if required. ➤ Spray of Jivamrut @ 3 lit./10 lit. of water for better growth. ➤ Keep field weed free by hand weeding and interculturing with Hand weeder upto 45 days after after sowing. ➤ Spray the plantbase botanical pesticides like, Agniastra or Brahmastra @ 0.5 Lit. per 10 lit. of water.
Elephant foot yam	➤ Weeding and intercultural operations should be carried out during clear weather condition.
Greater yam	➤ Weeding and intercultural operations should be carried out during clear weather condition. ➤ Give proper support or train branches of vines.
Tannia	➤ Weeding and intercultural operations should be carried out during clear weather condition.
Mango	➤ Find grubs of stem borer from gallery and holes on the stem and kill them. ➤ Procure healthy and disease free planting material from authorized and government nursery for the new plantation. ➤ Prepare pit for plantation of new grafts, apply 10 kg. of FYM or Ghanjivamrut fertilizer. Give 50-100 ml/tree of Agniastra or Brahmastra to manage termite and ants. ➤ Provide wooden or bamboo supports to young mango plants to protect them against strong winds and heavy rainfall. ➤ Apply FYM in Ghanjivamrut in old orchard and use the biofertilizer viz. azatobactor, PSB and KMB.

Valsad

Crop	Weather based Agro-Advisory
Sugarcane	Vegetative Stage/Grand Growth stage: ➤ After 6, 7 & 8 month of planting remove dry leaves and use it as mulch. ➤ Up root and burn whip smut and wilted plant from field. ➤ Drenching of <i>Carbendazim</i> 50 wp 10gm/10 lit water from prevent spread of both disease (Sugarcane red rot & Wilt) nearby healthy plants during clear weather condition. ➤ Prop the sugarcane canes with bamboos or tie the canes together in a clump with sugarcane leaves to protect against wind and heavy rain. ➤ Putting sugarcane leaves having whitefly puparia in 30 mesh wire cage. The adult whitefly emerge from puparia remain inside cage and parasitoid will come out and parasitize other whiteflies. ➤ Spraying <i>Azadirachtin</i> 10000 ppm @ 20 ml/10 lit. of water for management of sugarcane whitefly. Or ➤ Application of neem oil 0.5% (50 ml + 10 g detergent powder/10 litre water) first when initiation of sugarcane whitefly and second spray at 10 days after first spray for effective management of sugarcane whitefly under clear weather condition.
Kharif Paddy	Flowering stage/Reproductive stage: ➤ Keeps 2-3 cm standing water in field during vegetative/Reproductive growth. ➤ Apply split dose of Nitrogen fertilizer as per recommendation. i.e. 2nd dose @ 20 kg N/ha at 40-45 days after transplanting. ➤ Keep field weed free by hand weeding and interculturing with paddy weeder or by application of recommended post-emergence weedicide upto 45 days after transplanting.

	➤ After transplanting in paddy crop, monitor the crop condition and depending on rainfall, spraying of <i>Flubendiamide</i> 20 WG @ 2.5 g or <i>Chlorantraniliprole</i> 18.5 SC @ 3 ml or <i>Cartap hydrochloride</i> 75 SG @ 10 g per 10 litres of water during clear weather conditions for management of stem borer.
Pigeon Pea	➤ Weeding and intercultural operations should be carried out during clear weather condition.
Okra	➤ Special care should be taken that enough water should be supplied during the period of harvesting. ➤ Up root and burn yellow vein mosaic infected okra plant from field during clear weather condition. ➤ Spraying <i>Cyantraniliprole</i> 10.26 OD @ 18 ml or <i>Difenthiuron</i> 50 WP @ 12 g/10 Lit. for whitefly (vector) management during clear weather condition. ➤ Spraying of <i>Propargite</i> 57 EC @ 20 ml/10 Lit. or <i>Fenazaquin</i> 10 EC @ 10 ml./10 Lit. of water for red Spider Mite during clear weather condition.
Brinjal	➤ Transplanting should be carried out when the seedlings are 28-30 days old.
Tomato	➤ Transplanting should be carried out when the seedlings are 25-30 days old.
Chilly	➤ Seedlings of chilli become suitable for transplanting in about 35 to 40 days.
Pointed gourd	➤ Harvesting should be carried out as per maturity.
Little gourd	➤ Harvesting should be carried out as per maturity.
Elephant foot yam	➤ Apply fertilizer 50: 00: 37.5 NPK kg/ha. (109 kg Urea and 63 kg MOP per hectare). ➤ Take measures for weed control.
Greater yam	➤ Take measures for weed control. ➤ Give proper support or train branches of vines. ➤ Apply fertilizer 40:00:40 NPK kg/ha. (87 kg Urea and 67 kg MOP per hectare).
Tannia	➤ Take measures for weed control. ➤ Harvest the suitable tender leaves of Tannia. ➤ Apply fertilizer 40:00:40 NPK kg/ha. (87 kg Urea and 67 kg MOP per hectare).
Mango	➤ Find grubs of stem borer from gallery and holes on the stem and kill them. Do pouring of <i>Chlorpyrifos</i> 20 EC @ 25 ml/ 10 lit in hole and plug it with mud in clear weather condition. ➤ Procure healthy and disease free planting material from authorized and government nursery for the new plantation. ➤ Application of 10 % Bordeaux paste in cut/pruning branches of fruit tree during clear weather condition. ➤ Prepare pit for plantation of new grafts by filling 10 kg FYM, 1 kg SSP and 1 kg MOP. Give 50 ml/tree <i>Chlorpyrifos</i> to manage termite and ants during clear weather condition. ➤ Provide support to the young plants of mango plants to protect against rain and high wind speed. ➤ Apply recommended dose of fertilizer per tree @ 800 g Urea or 1.8 kg Ammonium Sulphate, 1.0 kg SSP and 1.3 kg MOP along with 100 kg FYM to adult tree. ➤ In addition RDF, give biofertilizers viz., <i>Azotobactor</i> 50 ml, PSB 50 ml, KMB 50 ml and <i>Trichoderma</i> 100 g per adult tree in monsoon. Before or after 15 days of fertilizer application.
Sapota	➤ Provide support to the young plants of Sapota plants to protect against rain and high wind speed. ➤ Procure healthy and disease free planting material from authorized and government nursery for the new plantation. ➤ Prepare pit for plantation of new grafts by filling 10 kg FYM, 1 kg SSP and 1 kg MOP. ➤ Give 50 ml/tree <i>Chlorpyrifos</i> to manage termite and ants during clear weather condition.

- Apply second split of 25% of RDF (Ammonium Sulphate -1.3 kg or Urea 0.6 kg, SSP 0.8 kg and MOP 0.3 kg + FYM 25 kg each tree) in clear weather condition.

AMFU Maktampur:

Bharuch

Main Crops	Growth Stage	Practices	Agro-Advisory
Sugar-cane	Grand Growth		• Irrigation Management: As the sugarcane is in its grand growth stage, its water requirement is peak. Monitor soil moisture closely and provide life-saving irrigation at regular intervals to prevent moisture stress and maintain cane weight. • Propping & Wind Protection: At 8 to 10 months, canes are tall and susceptible to lodging from gusty winds. Prop the crop immediately by tying adjacent canes together in clumps using their own dry leaves or bamboo stakes. • Detrashing (Leaf Stripping): Strip off lower dry leaves to improve field aeration. Reuse these leaves as mulch in the field to conserve soil moisture or use them for tying the canes. • Disease Management (Red Rot & Smut): Immediately uproot and burn any plants showing signs of whip smut, wilt, or Red Rot. Drench the soil around Red Rot-affected clumps with Carbendazim 50% WP (2 g/L of water) during clear weather. • Pest Management (Borers & Whitefly): Since the crop is very tall, granular applications for shoot borer are no longer effective and should be stopped. For top/stalk borers, manually remove and destroy dried shoots. If whitefly or Pyrilla infestations are observed, spray Azadirachtin 10000 ppm (20 ml/10 L) or a 0.5% Neem oil solution (50 ml Neem oil + 10 g detergent/10 L) during clear weather.
Cotton	Squaring/Flowering	Fertilizer • Avoid application of fertilizers during long dry spell condition or it should apply under optimum soil moisture condition. Spray 2 % urea during dry spell in standing cotton crop.	Irrigation • If irrigation facilities are available than apply need based irrigation through drip irrigation/alternatefurrow irrigation method by considering soil moisture condition in standing

			crop.
		Pest	• Pheromone Trap: Install pheromone traps in the field to effectively monitor and control pests. • Sucking Pest Management: To manage initial pest infestations, spray 500 g of neem seed extract or 50 ml of neem oil mixed in 10 liters of water. In case of heavy infestation, spray 3 g of Flonicamid 50 WG or 10 g of Diafenthiuron 50 WP per 10 liters of water strictly during clear weather.
		Diseases	• If root rot and wilt disease are observed in the cotton crop, undertake the drenching of infected plants using fungicides like Mancozeb 75 WP (27 g), Copper Oxychloride (40 g), or Carbendazim (10 g) mixed in 10 liters of water.
Rice	Vegetative/Tillering	Water Level	• Maintain 2–3 cm standing water; secure field bunds with overflow outlets to prevent nutrient wash-off during heavy rains.
		Weeding	• Weed manually or use a cono-weeder between rows to aerate roots; spray Bispyribac-sodium strictly during clear weather windows.
		Pest & Disease	• To control blast disease, spray tricyclazole 75 WP 6 g or Iprobenfos 48 EC 10 ml in 10 liter of water at an interval of 15 to 20 days in clear weather condition. • Spray a mixture of 1 g streptomycin and 20 g copper oxychloride in 400-500 liters of water per hectare during a rain-free period to control bacterial leaf blight.
Pigeon Pea	Vegetative		• Undertake interculturing, and hand weeding operations at 20, 40 and 60 days after sowing under vapsa condition in sown crops.
Castor	Vegetative		• Undertake the gap filling if required in sown maize crop. • Apply need based light irrigation by ensuring soil moisture in prevailing dry spell condition in sown castor crop.

Chilli/ Tomato/ Brinjal	Transplanting	• Undertake interculturing, weeding, gap filling, etc operations at vapsa condition in transplanted field. • In continuous humid and cloudy weather, if thrips incidence is observed undertake spraying of neem seed oil 30 to 50 ml in 10 liters of water. In case of heavy attack, spray of spinosad 45 SC 3 ml or spinetoram 11.7 SC 10 ml in 10 liter of water in clear and calm weather condition in chilli crop. • Due to continuous humid and cloudy weather condition, if sucking pest like white fly and jassid are observed in brinjal crop than undertake spraying of difenthiuron 50 WP 4 g or thiamethoxam 25 WG 4g in 10 liters of water in clear and calm weather condition.
Mango	Old and New orchard	• Young Plant Support: Provide strong physical support (staking) to young mango plants to protect them against the upcoming heavy rain and winds. • Pest Control (Stem Borer): Find grubs of the stem borer from galleries and holes on the stem and kill them. Before the heavy rain begins, pour Chlorpyrifos 20 EC (25 ml/10 liters of water) into the holes and plug them with mud during clear weather. • Disease Prevention: Apply 10% Bordeaux paste to cut/pruned branches immediately during clear weather to prevent fungal infections during the upcoming wet spell. • New Plantations: Procure healthy and disease-free planting material from authorized and government nurseries. Prepare pits by filling 10 kg FYM, 1 kg SSP, and 1 kg MOP, adding 50 ml/tree Chlorpyrifos to manage termites. • Fertilizer Application (Postponed): Strictly suspend the application of the Recommended Dose of Fertilizer (Urea, Ammonium Sulphate, SSP, MOP) and biofertilizers (Azotobacter, PSB, KMB, Trichoderma). Applying them now will result in severe nutrient loss due to leaching and runoff from the expected heavy rains. Wait until the rain spell passes.

Banana	Planting	• Weed Management & Interculturing: Perform interculturing and manual weeding in the planted field strictly when the soil reaches the 'Vapsa' (workable moisture) condition to keep the field weed-free and aerate the roots. • Irrigation Management: Since mostly dry weather is forecasted, monitor the soil moisture closely and apply light irrigation to the newly planted crop as needed.
Animal	General Advisory	
Cow /Buffalo /Goat /Sheep etc...	• Housing & Hygiene: Keep animals in clean, dry sheds. Regularly sprinkle dry slaked lime (Chuna) on the floor to maintain hygiene and prevent diseases like Foot Rot and Mastitis. • Fodder & Nutrition: Strictly avoid feeding old damp or fungus-infested fodder from past rains. Ensure a daily dose of mineral mixture and clean drinking water. • Healthcare: Administer deworming medicines as advised by a veterinarian and complete any pending vaccinations (FMD, HS, and BQ). • Pest Control: Clear any old stagnant water around the shed. Burn dry neem leaves in the evening to control mosquitoes, flies, and ticks.	
Poultry	• Shed Care & Wind Protection: With no rain expected, focus on proper ventilation. Hang curtains to block high winds, but keep them partially open during the day to ensure fresh air circulation. • Dry Litter & Lime: Regularly stir the bedding (litter) to keep it dry and prevent harmful ammonia buildup. Mix in slaked lime powder only if you spot any localized dampness. • Safe Feed: Continue storing feed bags on elevated wooden pallets to protect them from floor moisture and toxic fungal growth. • Clean Water & Health: Provide purified drinking water using sanitizers or chlorine tablets. Maintain regular vaccination schedules and consult a vet immediately if birds look dull or sick.	
Fisheries & Prawn	• Equipment & Oxygen: Continuous cloudy skies (5-7 octa) significantly reduce natural oxygen production. Keep paddle-wheel aerators running to maintain optimal dissolved oxygen levels, and ensure they are securely anchored against 15-18 km/h winds. • Feeding: Provide normal, optimal feeding quantities. Mix Vitamin C into the feed to help aquatic animals manage stress caused by the overcast weather.	

Surat

Main Crops	Growth Stage	Practices	Agro-Advisory
Sugar-cane	Grand Growth		• Irrigation Management: As the sugarcane is in its grand growth stage, its water requirement is peak. Monitor soil moisture closely and provide life-saving irrigation at regular intervals to prevent moisture stress and maintain cane weight. • Propping & Wind Protection: At 8 to 10 months, canes are tall and susceptible to lodging from gusty winds. Prop the crop immediately by tying adjacent canes together in clumps using their own dry leaves or bamboo stakes. • Detrashing (Leaf Stripping): Strip off lower dry leaves to improve field aeration. Reuse these leaves as mulch in the field to conserve soil moisture or use them for tying the canes. • Disease Management (Red Rot & Smut): Immediately uproot and burn any plants showing signs of whip smut, wilt, or Red Rot. Drench the soil around Red Rot-affected clumps with Carbendazim 50% WP (2 g/L of water) during clear weather. • Pest Management (Borers & Whitefly): Since the crop is very tall, granular applications for shoot borer are no longer effective and should be stopped. For top/stalk borers, manually remove and destroy dried shoots. If whitefly or Pyrilla infestations are observed, spray Azadirachtin 10000 ppm (20 ml/10 L) or a 0.5% Neem oil solution (50 ml Neem oil + 10 g detergent/10 L) during clear weather.
Cotton	Vegetative/Squaring/Branching	Fertilizer • Avoid application of fertilizers during long dry spell condition or it should apply under optimum soil moisture condition. Spray 2 % urea during dry spell in standing cotton crop.	• If irrigation facilities are available than apply need based irrigation through drip irrigation/alternatefurrow irrigation method by considering soil moisture condition in standing crop.
		Pest • Pheromone Trap: Install pheromone traps in the field to effectively monitor and control pests. • Sucking Pest Management: To manage initial pest infestations, spray 500 g of neem seed extract or 50 ml of neem oil mixed in 10 liters of water. In case of heavy infestation, spray 3 g of Flonicamid 50 WG or 10 g of Diafenthiuron 50 WP per 10 liters of water strictly during clear weather.	
		Diseases • If root rot and wilt disease are observed in the cotton crop, undertake the drenching of infected plants using fungicides like Mancozeb 75 WP (27 g), Copper	

			Oxychloride (40 g), or Carbendazim (10 g) mixed in 10 liters of water.
Rice	Vegetative	Water Level	Maintain 2–3 cm standing water; secure field bunds with overflow outlets to prevent nutrient wash-off during heavy rains.
		Weeding	Weed manually or use a cono-weeder between rows to aerate roots; spray Bispyribac-sodium strictly during clear weather windows.
		Pest & Disease	- Monitor the transplanted paddy crop and manage stem borer by spraying Flubendiamide 20 WG (2.5 g), Chlorantraniliprole 18.5 SC (3 ml), or Cartap hydrochloride 75 SG (10 g) in 10 liters of water during clear weather. - Spray a mixture of 1 g streptomycin and 20 g copper oxychloride in 400-500 liters of water per hectare during a rain-free period to control bacterial leaf blight.
Pigeon Pea	Vegetative		Undertake interculturing, and hand weeding operations at 20, 40 and 60 days after sowing under vapsa condition in sown crops.
Castor	Vegetative		- Undertake the gap filling if required in sown maize crop. - Apply need based light irrigation by ensuring soil moisture in prevailing dry spell condition in sown castor crop.
Soybean	Vegetative		- Undertake the interculturing and weeding operations at vapsa condition at 20 and 40 days after sowing in sown crop. - Undertake the interculturing and weeding operations at 30 days after sowing at vapsa condition even if pre-emergence or post-emergence herbicides have been sprayed for effective and economical weed control in the soybean crop. - Use one light trap per hectare to attract and destroy the moths of armyworm. If a severe infestation of armyworm is observed, spray a mixture of Emamectin Benzoate 1.9 EC (8.5 ml) or Indoxacarb 14.5 SC (7 ml) per 10 liters of water during clear weather.
Chilli/ Tomato/ Brinjal	Transplanting		- Undertake interculturing, weeding, gap filling, etc operations at vapsa condition in transplanted field. - In continuous humid and cloudy weather, if thrips incidence is observed undertake spraying of neem seed oil 30 to 50 ml in 10 liters of water. In case of heavy attack, spray of spinosad 45 SC 3 ml or spinetoram 11.7 SC 10 ml in 10 liter of water in clear and calm weather condition in chilli crop. - Due to continuous humid and cloudy weather condition, if sucking pest like white fly and jassid are observed in brinjal crop then undertake spraying of difenthiuron 50 WP 4 g or thiamethoxam 25 WG 4g in 10 liters of water in clear and calm weather condition.

Kharif Sorghum	Vegetative	• Fertilizer: Apply the top-dressing of Nitrogen (Urea) when the crop is 30–40 days old, ensuring the wind is calm. • Earthing-Up: Mound soil around the plant stems to provide root support and prevent lodging from strong winds. • Pest Control: Install fishmeal traps in the field to attract and control shoot flies. Spray recommended insecticides for stem borers only during clear weather.
Kharif Maize	Vegetative	• Apply need based light irrigation by considering weather and soil moisture condition in standing crop. • Destroy the adults of fall armyworm by installing one light trap per hectare. Install 5 pheromone traps per hectare to attract male moth of this pest. Spray Neem seed powder 500 gm (5% Extract) or Neembased insecticide 20 ml (1 EC) to 40 ml (0.15 EC) in 10 liter of water in whole plant properly in clear weather condition. In heavy incidence, spray chlorantraniliprole 18.5 SC 4 ml or spinetoram 11.7 SC 10ml in 10 liter of water to whole plant properly during clear weather condition. • Apply need based light irrigation by considering weather and soil moisture condition in standing crop.
Mango	Old and New orchard	• Young Plant Support: Provide strong physical support (staking) to young mango plants to protect them against the upcoming heavy rain and winds. • Pest Control (Stem Borer): Find grubs of the stem borer from galleries and holes on the stem and kill them. Before the heavy rain begins, pour Chlorpyrifos 20 EC (25 ml/10 liters of water) into the holes and plug them with mud during clear weather. • Disease Prevention: Apply 10% Bordeaux paste to cut/pruned branches immediately during clear weather to prevent fungal infections during the upcoming wet spell. • New Plantations: Procure healthy and disease-free planting material from authorized and government nurseries. Prepare pits by filling 10 kg FYM, 1 kg SSP, and 1 kg MOP, adding 50 ml/tree Chlorpyrifos to manage termites. • Fertilizer Application (Postponed): Strictly suspend the application of the Recommended Dose of Fertilizer (Urea, Ammonium Sulphate, SSP, MOP) and biofertilizers (Azotobacter, PSB, KMB, Trichoderma). Applying them now will result in severe nutrient loss due to leaching and runoff from the expected heavy rains. Wait until the rain spell passes.

Banana	Planting		• Weed Management & Interculturing: Perform interculturing and manual weeding in the planted field strictly when the soil reaches the 'Vapsa' (workable moisture) condition to keep the field weed-free and aerate the roots. • Irrigation Management: Since mostly dry weather is forecasted, monitor the soil moisture closely and apply light irrigation to the newly planted crop as needed.
Animal	General Advisory		
Cow /Buffalo /Goat /Sheep etc...			• Housing & Hygiene: Keep animals in clean, dry sheds. Regularly sprinkle dry slaked lime (Chuna) on the floor to maintain hygiene and prevent diseases like Foot Rot and Mastitis. • Fodder & Nutrition: Strictly avoid feeding old damp or fungus-infested fodder from past rains. Ensure a daily dose of mineral mixture and clean drinking water. • Healthcare: Administer deworming medicines as advised by a veterinarian and complete any pending vaccinations (FMD, HS, and BQ). • Pest Control: Clear any old stagnant water around the shed. Burn dry neem leaves in the evening to control mosquitoes, flies, and ticks.
Poultry			• Shed Care & Wind Protection: With no rain expected, focus on proper ventilation. Hang curtains to block high winds, but keep them partially open during the day to ensure fresh air circulation. • Dry Litter & Lime: Regularly stir the bedding (litter) to keep it dry and prevent harmful ammonia buildup. Mix in slaked lime powder only if you spot any localized dampness. • Safe Feed: Continue storing feed bags on elevated wooden pallets to protect them from floor moisture and toxic fungal growth. • Clean Water & Health: Provide purified drinking water using sanitizers or chlorine tablets. Maintain regular vaccination schedules and consult a vet immediately if birds look dull or sick.
Fisheries & Prawn			• Equipment & Oxygen: Continuous cloudy skies (5-7 octa) significantly reduce natural oxygen production. Keep paddle-wheel aerators running to maintain optimal dissolved oxygen levels, and ensure they are securely anchored against 15-18 km/h winds. • Feeding: Provide normal, optimal feeding quantities. Mix Vitamin C into the feed to help aquatic animals manage stress caused by the overcast weather.

Narmada

Main Crops	Growth Stage	Practices	Agro-Advisory
Sugar-cane	Grand Growth		• Irrigation Management: As the sugarcane is in its grand growth stage, its water requirement is peak. Monitor soil moisture closely and provide life-saving irrigation at regular intervals to prevent moisture stress and maintain cane weight. • Propping & Wind Protection: At 8 to 10 months, canes are tall and susceptible to lodging from gusty winds. Prop the crop immediately by tying adjacent

			canes together in clumps using their own dry leaves or bamboo stakes. • Detrashing (Leaf Stripping): Strip off lower dry leaves to improve field aeration. Reuse these leaves as mulch in the field to conserve soil moisture or use them for tying the canes. • Disease Management (Red Rot & Smut): Immediately uproot and burn any plants showing signs of whip smut, wilt, or Red Rot. Drench the soil around Red Rot-affected clumps with Carbendazim 50% WP (2 g/L of water) during clear weather. • Pest Management (Borers & Whitefly): Since the crop is very tall, granular applications for shoot borer are no longer effective and should be stopped. For top/stalk borers, manually remove and destroy dried shoots. If whitefly or Pyrilla infestations are observed, spray Azadirachtin 10000 ppm (20 ml/10 L) or a 0.5% Neem oil solution (50 ml Neem oil + 10 g detergent/10 L) during clear weather.
Cotton	Vegetative/Squaring/Branching	Fertilizer	• Avoid application of fertilizers during long dry spell condition or it should apply under optimum soil moisture condition. Spray 2 % urea during dry spell in standing cotton crop.
		Irrigation	• If irrigation facilities are available than apply need based irrigation through drip irrigation/alternate furrow irrigation method by considering soil moisture condition in standing crop.
		Pest	• Pheromone Trap: Install pheromone traps in the field to effectively monitor and control pests. • Sucking Pest Management: To manage initial pest infestations, spray 500 g of neem seed extract or 50 ml of neem oil mixed in 10 liters of water. In case of heavy infestation, spray 3 g of Flonicamid 50 WG or 10 g of Diafenthiuron 50 WP per 10 liters of water strictly during clear weather.
		Diseases	• If root rot and wilt disease are observed in the cotton crop, undertake the drenching of infected plants using fungicides like Mancozeb 75 WP (27 g), Copper Oxychloride (40 g), or Carbendazim (10 g) mixed in 10 liters of water.
Rice	Vegetative	Water Level	• Maintain 2–3 cm standing water; secure field bunds with overflow outlets to prevent nutrient wash-off during heavy rains.
		Weeding	• Weed manually or use a cono-weeder between rows to aerate roots; spray Bispyribac-sodium strictly during clear weather windows.
		Pest & Disease	• Monitor the transplanted paddy crop and manage stem borer by spraying Flubendiamide 20 WG (2.5 g), Chlorantraniliprole 18.5 SC (3 ml), or Cartap hydrochloride 75 SG (10 g) in 10 liters of water during clear weather.

			Spray a mixture of 1 g streptocycline and 20 g copper oxychloride in 400-500 liters of water per hectare during a rain-free period to control bacterial leaf blight.
Pigeon Pea	Vegetative		Undertake interculturing, and hand weeding operations at 20, 40 and 60 days after sowing under vapsa condition in sown crops.
Castor	Vegetative		- Undertake the gap filling if required in sown maize crop. - Apply need based light irrigation by ensuring soil moisture in prevailing dry spell condition in sown castor crop.
Chilli/ Tomato/ Brinjal	Transplanting		- Undertake interculturing, weeding, gap filling, etc operations at vapsa condition in transplanted field. - In continuous humid and cloudy weather, if thrips incidence is observed undertake spraying of neem seed oil 30 to 50 ml in 10 liters of water. In case of heavy attack, spray of spinosad 45 SC 3 ml or spinetoram 11.7 SC 10 ml in 10 liter of water in clear and calm weather condition in chilli crop. - Due to continuous humid and cloudy weather condition, if sucking pest like white fly and jassid are - observed in brinjal crop then undertake spraying of difenthiuron 50 WP 4 g or thiamethoxam 25 WG 4g in 10 liters of water in clear and calm weather condition.
Kharif Sorghum	Vegetative		Weeding & Interculturing: Perform interculturing and manual weeding strictly when the soil reaches the 'Vapsa'(workable moisture) condition. Fertilizer: Apply the top-dressing of Nitrogen (Urea) when the crop is 30–40 days old, ensuring the wind is calm. Earthing-Up: Mound soil around the plant stems to provide root support and prevent lodging from strong winds. Pest Control: Install fishmeal traps in the field to attract and control shoot flies. Spray recommended insecticides for stem borers only during clear weather.
Kharif Maize	Vegetative		- Apply need based light irrigation by considering weather and soil moisture condition in standing crop. - Destroy the adults of fall armyworm by installing one light trap per hectare. Install 5 pheromone traps per hectare to attract male moth of this pest. Spray Neem seed powder 500 gm (5% Extract) or Neem based insecticide 20 ml (1 EC) to 40 ml (0.15 EC) in 10 liter of water in whole plant properly in clear weather condition. In heavy incidence, spray chlorantraniliprole 18.5 SC 4 ml or spinetoram 11.7 SC 10 ml in 10 liter of water to whole plant properly during clear weather condition. - Apply need based light irrigation by considering weather and soil moisture condition in standing crop.

Mango	Old and New orchard		• Young Plant Support: Provide strong physical support (staking) to young mango plants to protect them against the upcoming heavy rain and winds. • Pest Control (Stem Borer): Find grubs of the stem borer from galleries and holes on the stem and kill them. Before the heavy rain begins, pour Chlorpyrifos 20 EC (25 ml/10 liters of water) into the holes and plug them with mud during clear weather. • Disease Prevention: Apply 10% Bordeaux paste to cut/pruned branches immediately during clear weather to prevent fungal infections during the upcoming wet spell. • New Plantations: Procure healthy and disease-free planting material from authorized and government nurseries. Prepare pits by filling 10 kg FYM, 1 kg SSP, and 1 kg MOP, adding 50 ml/tree Chlorpyrifos to manage termites. • Fertilizer Application (Postponed): Strictly suspend the application of the Recommended Dose of Fertilizer (Urea, Ammonium Sulphate, SSP, MOP) and biofertilizers (Azotobacter, PSB, KMB, Trichoderma). Applying them now will result in severe nutrient loss due to leaching and runoff from the expected heavy rains. Wait until the rain spell passes.
Banana	Planting		• Weed Management & Interculturing: Perform interculturing and manual weeding in the planted field strictly when the soil reaches the 'Vapsa' (workable moisture) condition to keep the field weed-free and aerate the roots. • Irrigation Management: Since mostly dry weather is forecasted, monitor the soil moisture closely and apply light irrigation to the newly planted crop as needed.
Animal	General Advisory		
Cow /Buffalo /Goat /Sheep etc...	• Housing & Hygiene: Keep animals in clean, dry sheds. Regularly sprinkle dry slaked lime (Chuna) on the floor to maintain hygiene and prevent diseases like Foot Rot and Mastitis. • Fodder & Nutrition: Strictly avoid feeding old damp or fungus-infested fodder from past rains. Ensure a daily dose of mineral mixture and clean drinking water. • Healthcare: Administer deworming medicines as advised by a veterinarian and complete any pending vaccinations (FMD, HS, and BQ). • Pest Control: Clear any old stagnant water around the shed. Burn dry neem leaves in the evening to control mosquitoes, flies, and ticks.		
Poultry	• Shed Care & Wind Protection: With no rain expected, focus on proper ventilation. Hang curtains to block high winds, but keep them partially open during the day to ensure fresh air circulation. • Dry Litter & Lime: Regularly stir the bedding (litter) to keep it dry and prevent harmful ammonia buildup. Mix in slaked lime powder only if you spot any localized dampness. • Safe Feed: Continue storing feed bags on elevated wooden pallets to protect them from floor moisture and toxic fungal growth. • Clean Water & Health: Provide purified drinking water using sanitizers		

or chlorine tablets. Maintain regular vaccination schedules and consult a vet immediately if birds look dull or sick.

Tapi

Main Crops	Growth Stage	Practices	Agro-Advisory
Sugar-cane	Grand Growth		• Irrigation Management: As the sugarcane is in its grand growth stage, its water requirement is peak. Monitor soil moisture closely and provide life-saving irrigation at regular intervals to prevent moisture stress and maintain cane weight. • Propping & Wind Protection: At 8 to 10 months, canes are tall and susceptible to lodging from gusty winds. Prop the crop immediately by tying adjacent canes together in clumps using their own dry leaves or bamboo stakes. • Detrashing (Leaf Stripping): Strip off lower dry leaves to improve field aeration. Reuse these leaves as mulch in the field to conserve soil moisture or use them for tying the canes. • Disease Management (Red Rot & Smut): Immediately uproot and burn any plants showing signs of whip smut, wilt, or Red Rot. Drench the soil around Red Rot-affected clumps with Carbendazim 50% WP (2 g/L of water) during clear weather. • Pest Management (Borers & Whitefly): Since the crop is very tall, granular applications for shoot borer are no longer effective and should be stopped. For top/stalk borers, manually remove and destroy dried shoots. If whitefly or Pyrilla infestations are observed, spray Azadirachtin 10000 ppm (20 ml/10 L) or a 0.5% Neem oil solution (50 ml Neem oil + 10 g detergent/10 L) during clear weather.
Cotton	Vegetative/Squaring/Branching	Fertilizer Irrigation Pest	• Avoid application of fertilizers during long dry spell condition or it should apply under optimum soil moisture condition. Spray 2 % urea during dry spell in standing cotton crop. • If irrigation facilities are available than apply need based irrigation through drip irrigation/alternatefurrow irrigation method by considering soil moisture condition in standing crop. • Pheromone Trap: Install pheromone traps in the field to effectively monitor and control pests. • Sucking Pest Management: To manage initial pest infestations, spray 500 g of neem seed extract or 50 ml of neem oil mixed in 10 liters of water. In case of heavy infestation, spray 3 g of Flonicamid 50 WG or 10 g of Diafenthiuron 50 WP per 10 liters of water strictly during clear weather.

		Diseases	If root rot and wilt disease are observed in the cotton crop, undertake the drenching of infected plants using fungicides like Mancozeb 75 WP (27 g), Copper Oxychloride (40 g), or Carbendazim (10 g) mixed in 10 liters of water.
Rice	Vegetative	Water Level	Maintain 2–3 cm standing water; secure field bunds with overflow outlets to prevent nutrient wash-off during heavy rains.
		Weeding	Weed manually or use a cono-weeder between rows to aerate roots; spray Bispyribac-sodium strictly during clear weather windows.
		Pest & Disease	- Monitor the transplanted paddy crop and manage stem borer by spraying Flubendiamide 20 WG (2.5 g), Chlorantraniliprole 18.5 SC (3 ml), or Cartap hydrochloride 75 SG (10 g) in 10 liters of water during clear weather. - Spray a mixture of 1 g streptomycin and 20 g copper oxychloride in 400-500 liters of water per hectare during a rain-free period to control bacterial leaf blight.
Pigeon Pea	Vegetative		Undertake interculturing, and hand weeding operations at 20, 40 and 60 days after sowing under vapsa condition in sown crops.
Castor	Vegetative		- Undertake the gap filling if required in sown maize crop. - Apply need based light irrigation by ensuring soil moisture in prevailing dry spell condition in sown castor crop.
Chilli/ Tomato/ Brinjal	Transplanting		- Undertake interculturing, weeding, gap filling, etc operations at vapsa condition in transplanted field. - In continuous humid and cloudy weather, if thrips incidence is observed undertake spraying of neem seed oil 30 to 50 ml in 10 liters of water. In case of heavy attack, spray of spinosad 45 SC 3 ml or spinetoram 11.7 SC 10 ml in 10 liter of water in clear and calm weather condition in chilli crop. - Due to continuous humid and cloudy weather condition, if sucking pest like white fly and jassid are observed in brinjal crop then undertake spraying of difenthiuron 50 WP 4 g or thiamethoxam 25 WG 4g in 10 liters of water in clear and calm weather condition.
Kharif Sorghum	Vegetative		Weeding & Interculturing: Perform interculturing and manual weeding strictly when the soil reaches the 'Vapsa'(workable moisture) condition. Fertilizer: Apply the top-dressing of Nitrogen (Urea) when the crop is 30–40 days old, ensuring the wind is calm. Earthing-Up: Mound soil around the plant stems to provide root support and prevent lodging from strong winds. Pest Control: Install fishmeal traps in the field to attract and control shoot flies. Spray recommended insecticides for stem borers only during clear weather.

Kharif Maize	Vegetative	• Apply need based light irrigation by considering weather and soil moisture condition in standing crop. • Destroy the adults of fall armyworm by installing one light trap per hectare. Install 5 pheromone traps per hectare to attract male moth of this pest. Spray Neem seed powder 500 gm (5% Extract) or Neem based insecticide 20 ml (1 EC) to 40 ml (0.15 EC) in 10 liter of water in whole plant properly in clear weather condition. In heavy incidence, spray chlorantraniliprole 18.5 SC 4 ml or spinetoram 11.7 SC 10 ml in 10 liter of water to whole plant properly during clear weather condition. • Apply need based light irrigation by considering weather and soil moisture condition in standing crop.
Mango	Old and New orchard	• Young Plant Support: Provide strong physical support (staking) to young mango plants to protect them against the upcoming heavy rain and winds. • Pest Control (Stem Borer): Find grubs of the stem borer from galleries and holes on the stem and kill them. Before the heavy rain begins, pour Chlorpyrifos 20 EC (25 ml/10 liters of water) into the holes and plug them with mud during clear weather. • Disease Prevention: Apply 10% Bordeaux paste to cut/pruned branches immediately during clear weather to prevent fungal infections during the upcoming wet spell. • New Plantations: Procure healthy and disease-free planting material from authorized and government nurseries. Prepare pits by filling 10 kg FYM, 1 kg SSP, and 1 kg MOP, adding 50 ml/tree Chlorpyrifos to manage termites. • Fertilizer Application (Postponed): Strictly suspend the application of the Recommended Dose of Fertilizer (Urea, Ammonium Sulphate, SSP, MOP) and biofertilizers (Azotobacter, PSB, KMB, Trichoderma). Applying them now will result in severe nutrient loss due to leaching and runoff from the expected heavy rains. Wait until the rain spell passes.
Animal	General Advisory	
Cow /Buffalo /Goat /Sheep etc...	• Housing & Hygiene: Keep animals in clean, dry sheds. Regularly sprinkle dry slaked lime (Chuna) on the floor to maintain hygiene and prevent diseases like Foot Rot and Mastitis. • Fodder & Nutrition: Strictly avoid feeding old damp or fungus-infested fodder from past rains. Ensure a daily dose of mineral mixture and clean drinking water. • Healthcare: Administer deworming medicines as advised by a veterinarian and complete any pending vaccinations (FMD, HS, and BQ). • Pest Control: Clear any old stagnant water around the shed. Burn dry neem leaves in the evening to control mosquitoes, flies, and ticks.	
Poultry	• Shed Care & Wind Protection: With no rain expected, focus on proper ventilation. Hang curtains to block high winds, but keep them partially open during the day to ensure fresh air circulation. • Dry Litter & Lime: Regularly stir the bedding (litter) to keep it dry and	

	prevent harmful ammonia buildup. Mix in slaked lime powder only if you spot any localized dampness. • Safe Feed: Continue storing feed bags on elevated wooden pallets to protect them from floor moisture and toxic fungal growth. • Clean Water & Health: Provide purified drinking water using sanitizers or chlorine tablets. Maintain regular vaccination schedules and consult a vet immediately if birds look dull or sick.
--	--

AMFU Anand:

Anand

Main Crops	Stage	Disease/ Pest/ Management	Advisory
BANANA	Vegetative		• Undertake the interculturing and weeding at vapsa condition in planted crop. • Apply need based light irrigation by considering weather and soil moisture condition in standing crop. • Undertake earthing up up to 15-20 cm height around root zone and frequently remove suckers around mother plant.
KHARIF RICE	Flowering / Grain filling		• Maintain 5-7 cm stagnant water in rice field. • Spray mixture of streptocycline 1 g +copper oxychloride 20 g fungicides in 400-500 liter of water per hectare for control of bacterial leaf blight disease during rain free period. • To control blast disease, spray tricyclazole 75 WP 6 g or Iprobenfos 48 EC 10 ml in 10 liter of water at an interval of 15 to 20 days in clear weather condition.
COTTON	Flowering / Boll formation		• If irrigation facilities are available than apply need based irrigation through drip irrigation/alternate furrow irrigation method by considering soil moisture condition in standing crop. • Avoid application of fertilizers during long dry spell condition or it should apply under optimum soil moisture condition. Spray 2 % urea during dry spell in standing cotton crop. • In initial stage of incidence of sucking pest, spray neem seed extract 500 g or Neem oil 50 ml in 10 liters of water. In heavy incidence, spray flonicamid 50 WG 3 g or difenthiuron 50 WP 10 g in 10 liters of water in clear and calm weather condition. • If root rot and wilt disease is observed in cotton crop than undertake the drenching of fungicides like mancozeb 75 WP 27 g or copper oxychloride 40 g or carbendazim 10 g in 10 liter of water to infected plants and also apply urea or ammonium sulfate at 4 to 5 days after drenching of above fungicide.
KHARIF PEARL MILLET	Flowering		• Undertake the interculturing and weeding operations at 20 and 40 days after sowing under vapsa condition in sown crop. • Apply nitrogen 40 kg/ha as a top dressing after weeding and thinning under optimum soil moisture level at one month after sowing. • To control blast disease, carry out two spray of carbendazim 50 WP 10

			g in 10 liters water at an interval of 15-20 days in clear sky condition. • Apply need based light irrigation by considering weather and soil moisture condition in standing crop.
FODDER CROPS	Vegetative/ Flowering		• Apply need based light irrigation by considering weather and soil moisture condition in standing fodder crops. • Cutting of fodder sorghum should be done at 60 to 65 days (50 % flowering stage) after sowing in single cut varieties. • First cutting of fodder sorghum should be done at 45 to 45 days and next cutting at subsequent 30 days in multi cut varieties.
BIDI TOBACCO	Transplanting		• Undertake transplanting of ready seedlings of bidi tobacco at spacing of 90 cm x 75 cm or 105 cm x 90 cm from third week of August to third week of September under assured irrigation facilities.
GREEN GRAM/ BLACK GRAM	Flowering / Pod formation		• Mosaic of leaf curl disease is transmitted due to white fly so to control it, spray imidacloprid 17.8 SC 3 ml or acetamiprid 20 SP 2 g or azadirachtin 40 ml in 10 liter of water in clear sky condition. • If Aphid incidence is observed in green gram or black gram crop than undertake the spraying of imidacloprid 70 WS 7.5 g or thiamethoxam 70 WS 2.8 g in 10 liter of water in clear sky condition. • Apply need based light irrigation by considering weather and soil moisture condition in standing crops.
PIGEON PEA	Vegetative/ Flowering		• Undertake interculturing, and hand weeding operations at 40 and 60 days after sowing under vapsa condition in sown crops. • Apply need based light irrigation by considering weather and soil moisture condition in standing crops.
CHILLI/ TOMATO/ BRINJAL	Vegetative/ Flowering		• Undertake interculturing, weeding, gap filling, etc operations at vapsa condition in transplanted field. • In continuous humid and cloudy weather, if thrips incidence is observed undertake spraying of neem seed oil 30 to 50 ml in 10 liters of water. In case of heavy attack, spray of spinosad 45 SC 3 ml or spinetoram 11.7 SC 10 ml in 10 liter of water in clear and calm weather condition in chilli crop. • In tomato crop, at initial incidence of Pinworm, install 40 yellow sticky trap per hectare. In heavy incidence, spray with Chlorantraniliprole 18.5 SC 3 ml or Flubendiamide 480 SC 3 ml in 10 liter of water in clear weather condition. • Due to continuous humid and cloudy weather condition, if sucking pest like white fly and jassid are observed in brinjal crop than undertake spraying of difenthiuron 50 WP 4 g or thiamethoxam 25 WG 4 g in 10 liters of water in clear and calm weather condition. • Apply need based light irrigation by considering weather and soil moisture condition in standing crops.
OKRA	Vegetative/ Flowering		• If yellow vein mosaic disease is observed in okra crop, than undertake the spraying of insecticides like tolfeprydaz 15 EC 20 ml or fenpropathrin 30 EC 3.4 ml in 10 liters of water in clear and calm weather condition. • Apply need based light irrigation by considering weather and soil moisture condition in standing crops.
LIVESTOCK			• Spray phenyl in the animal shed to avoid flies and mosquitos. For control of tick and other parasite, spray chemicals as per prescription of nearest animal clinic.

CK	- Give mineral mixture balance food to animals. Apply feed silage and hay material along with concentrate feed. Use chaff cutter for fodder. - Keep the fodder at safe place to avoid spoilage from rainfall.
-----------	--

Chhota Udepur

Main Crops	Stage	Disease/ Pest/ Management	Advisory
BANANA	Vegetative		- Undertake the interculturing and weeding at vapsa condition in planted crop. - Apply need based light irrigation by considering weather and soil moisture condition in standing crop. - Undertake earthing up up to 15-20 cm height around root zone and frequently remove suckers around mother plant.
COTTON	Flowering/ Boll formation		- If irrigation facilities are available than apply need based irrigation through drip irrigation/alternate furrow irrigation method by considering soil moisture condition in standing crop. - Avoid application of fertilizers during long dry spell condition or it should apply under optimum soil moisture condition. Spray 2 % urea during dry spell in standing cotton crop. - In initial stage of incidence of sucking pest, spray neem seed extract 500 g or Neem oil 50 ml in 10 liters of water. In heavy incidence, spray flonicamid 50 WG 3 g or difenthiuron 50 WP 10 g in 10 liters of water in clear and calm weather condition. - If root rot and wilt disease is observed in cotton crop than undertake the drenching of fungicides like mancozeb 75 WP 27 g or copper oxychloride 40 g or carbendazim 10 g in 10 liter of water to infected plants and also apply urea or ammonium sulfate at 4 to 5 days after drenching of above fungicide.
KHARIF RICE	Flowering/ Grain filling		- Maintain 5-7 cm stagnant water in rice field. - Spray mixture of streptocycline 1 g + copper oxychloride 20 g fungicides in 400-500 liter of water per hectare for control of bacterial leaf blight disease during rain free period. - To control blast disease, spray tricyclazole 75 WP 6 g or Iprobenfos 48 EC 10 ml in 10 liter of water at an interval of 15 to 20 days in clear weather condition.
GREEN GRAM/ BLACK GRAM	Flowering/ Pod formation		- Mosaic of leaf curl disease is transmitted due to white fly so to control it, spray imidacloprid 17.8 SC 3 ml or acetamiprid 20 SP 2 g or azadirachtin 40 ml in 10 liter of water in clear sky condition. - If Aphid incidence is observed in green gram or black gram crop than undertake the spraying of imidacloprid 70 WS 7.5 g or thiamethoxam 70 WS 2.8 g in 10 liter of water in clear sky condition. - Apply need based light irrigation by considering weather and soil moisture condition in standing crops.
PIGEON PEA	Vegetative /		Undertake interculturing, and hand weeding operations at 40 and 60 days after sowing under vapsa condition in sown crops.

	Flowering		• Apply need based light irrigation by considering weather and soil moisture condition in standing crops.
SOYBEAN	Pod formation / Grain filling		• Use one light trap per hectare to attract and destroy the moths of armyworm. If a severe infestation of armyworm is observed, spray a mixture of Emamectin Benzoate 1.9 EC (8.5 ml) or Indoxacarb 14.5 SC (7 ml) per 10 liters of water during clear weather. • Apply need based light irrigation by considering weather and soil moisture condition in standing crop.
KHARIF MAIZE	Flowering/ Tasseling		• Apply need based light irrigation by considering weather and soil moisture condition in standing crop. • Destroy the adults of fall armyworm by installing one light trap per hectare. Install 5 pheromone traps per hectare to attract male moth of this pest. Spray Neem seed powder 500 gm (5% Extract) or Neem based insecticide 20 ml (1 EC) to 40 ml (0.15 EC) in 10 liter of water in whole plant properly in clear weather condition. In heavy incidence, spray chlorantraniliprole 18.5 SC 4 ml or spinetoram 11.7 SC 10 ml in 10 liter of water to whole plant properly during clear weather condition. • Apply need based light irrigation by considering weather and soil moisture condition in standing crop.
CHILLI/ TOMATO/ BRINJAL	Vegetative / Flowering		• Undertake interculturing, weeding, gap filling, etc operations at vapsa condition in transplanted field. • In continuous humid and cloudy weather, if thrips incidence is observed undertake spraying of neem seed oil 30 to 50 ml in 10 liters of water. In case of heavy attack, spray of spinosad 45 SC 3 ml or spinetoram 11.7 SC 10 ml in 10 liter of water in clear and calm weather condition in chilli crop. • In tomato crop, at initial incidence of Pinworm, install 40 yellow sticky trap per hectare. In heavy incidence, spray with Chlorantraniliprole 18.5 SC 3 ml or Flubendiamide 480 SC 3 ml in 10 liter of water in clear weather condition. • Due to continuous humid and cloudy weather condition, if sucking pest like white fly and jassid are observed in brinjal crop then undertake spraying of difenthiuron 50 WP 4 g or thiamethoxam 25 WG 4 g in 10 liters of water in clear and calm weather condition. • Apply need based light irrigation by considering weather and soil moisture condition in standing crops.
CASTOR	Vegetative		• Undertake interculturing, weeding, gap filling, etc operations at vapsa condition in sown crop. • Apply need based light irrigation by ensuring soil moisture in prevailing dry spell condition in sown castor crop.
LIVESTOCK	• Spray phenyl in the animal shed to avoid flies and mosquitos. For control of tick and other parasite, spray chemicals as per prescription of nearest animal clinic. • Give mineral mixture balance food to animals. Apply feed silage and hay material along with concentrate feed. Use chaff cutter for fodder. Keep the fodder at safe place to avoid spoilage from rainfall.		

Dahod

Main Crops	Stage	Disease/ Pest/ Management	Advisory
COTTON	Flowering/ Boll formation		• If irrigation facilities are available than apply need based irrigation through drip irrigation/alternate furrow irrigation method by considering soil moisture condition in standing crop. • Avoid application of fertilizers during long dry spell condition or it should apply under optimum soil moisture condition. Spray 2 % urea during dry spell in standing cotton crop. • In initial stage of incidence of sucking pest, spray neem seed extract 500 g or Neem oil 50 ml in 10 liters of water. In heavy incidence, spray flonicamid 50 WG 3 g or difenthiuron 50 WP 10 g in 10 liters of water in clear and calm weather condition. • If root rot and wilt disease is observed in cotton crop than undertake the drenching of fungicides like mancozeb 75 WP 27 g or copper oxychloride 40 g or carbendazim 10 g in 10 liter of water to infected plants and also apply urea or ammonium sulfate at 4 to 5 days after drenching of above fungicide.
KHARIF MAIZE	Flowering/ Tasseling		• Apply need based light irrigation by considering weather and soil moisture condition in standing crop. • Destroy the adults of fall armyworm by installing one light trap per hectare. Install 5 pheromone traps per hectare to attract male moth of this pest. Spray Neem seed powder 500 gm (5% Extract) or Neem based insecticide 20 ml (1 EC) to 40 ml (0.15 EC) in 10 liter of water in whole plant properly in clear weather condition. In heavy incidence, spray chlorantraniliprole 18.5 SC 4 ml or spinetoram 11.7 SC 10 ml in 10 liter of water to whole plant properly during clear weather condition. • Apply need based light irrigation by considering weather and soil moisture condition in standing crop.
KHARIF RICE	Flowering/ Grain filling		• Maintain 5-7 cm stagnant water in rice field. • Spray mixture of streptocycline 1 g + copper oxychloride 20 g fungicides in 400-500 liter of water per hectare for control of bacterial leaf blight disease during rain free period. • To control blast disease, spray tricyclazole 75 WP 6 g or Iprobenfos 48 EC 10 ml in 10 liter of water at an interval of 15 to 20 days in clear weather condition.
GREEN GRAM/ BLACK GRAM	Flowering/ Pod formation		• Mosaic of leaf curl disease is transmitted due to white fly so to control it, spray imidacloprid 17.8 SC 3 ml or acetamiprid 20 SP 2 g or azadirachtin 40 ml in 10 liter of water in clear sky condition. • If Aphid incidence is observed in green gram or black gram crop than undertake the spraying of imidacloprid 70 WS 7.5 g or thiamethoxam 70 WS 2.8 g in 10 liter of water in clear sky condition. • Apply need based light irrigation by considering weather and soil moisture condition in standing crops.

PIGEON PEA	Vegetative / Flowering		- Undertake interculturing, and hand weeding operations at 40 and 60 days after sowing under vapsa condition in sown crops. - Apply need based light irrigation by considering weather and soil moisture condition in standing crops.
SOYBEAN	Pod formation / Grain filling		- Use one light trap per hectare to attract and destroy the moths of armyworm. If a severe infestation of armyworm is observed, spray a mixture of Emamectin Benzoate 1.9 EC (8.5 ml) or Indoxacarb 14.5 SC (7 ml) per 10 liters of water during clear weather. - Apply need based light irrigation by considering weather and soil moisture condition in standing crop.
CHILLI/ TOMATO/ BRINJAL	Vegetative / Flowering		- Undertake interculturing, weeding, gap filling, etc operations at vapsa condition in transplanted field. - In continuous humid and cloudy weather, if thrips incidence is observed undertake spraying of neem seed oil 30 to 50 ml in 10 liters of water. In case of heavy attack, spray of spinosad 45 SC 3 ml or spinetoram 11.7 SC 10 ml in 10 liter of water in clear and calm weather condition in chilli crop. - In tomato crop, at initial incidence of Pinworm, install 40 yellow sticky trap per hectare. In heavy incidence, spray with Chlorantraniliprole 18.5 SC 3 ml or Flubendiamide 480 SC 3 ml in 10 liter of water in clear weather condition. - Due to continuous humid and cloudy weather condition, if sucking pest like white fly and jassid are observed in brinjal crop then undertake spraying of difenthiuron 50 WP 4 g or thiamethoxam 25 WG 4 g in 10 liters of water in clear and calm weather condition. - Apply need based light irrigation by considering weather and soil moisture condition in standing crops.
CASTOR	Vegetative		- Undertake interculturing, weeding, gap filling, etc operations at vapsa condition in sown crop. - Apply need based light irrigation by ensuring soil moisture in prevailing dry spell condition in sown castor crop.
LIVESTOCK			- Spray phenyl in the animal shed to avoid flies and mosquitos. For control of tick and other parasite, spray chemicals as per prescription of nearest animal clinic. - Give mineral mixture balance food to animals. Apply feed silage and hay material along with concentrate feed. Use chaff cutter for fodder. - Keep the fodder at safe place to avoid spoilage from rainfall.

Kheda

Main Crops	Stage	Disease/ Pest/ Management	Advisory
BANANA	Vegetative		- Undertake the interculturing and weeding at vapsa condition in planted crop. - Apply need based light irrigation by considering weather and soil moisture condition in standing crop. - Undertake earthing up up to 15-20 cm height around root zone and

			frequently remove suckers around mother plant.
COTTON	Flowering/ Boll formation		• If irrigation facilities are available than apply need based irrigation through drip irrigation/alternate furrow irrigation method by considering soil moisture condition in standing crop. • Avoid application of fertilizers during long dry spell condition or it should apply under optimum soil moisture condition. Spray 2 % urea during dry spell in standing cotton crop. • In initial stage of incidence of sucking pest, spray neem seed extract 500 g or Neem oil 50 ml in 10 liters of water. In heavy incidence, spray flonicamid 50 WG 3 g or difenthiuron 50 WP 10 g in 10 liters of water in clear and calm weather condition. • If root rot and wilt disease is observed in cotton crop than undertake the drenching of fungicides like mancozeb 75 WP 27 g or copper oxychloride 40 g or carbendazim 10 g in 10 liter of water to infected plants and also apply urea or ammonium sulfate at 4 to 5 days after drenching of above fungicide.
KHARIF RICE	Flowering/ Grain filling		• Maintain 5-7 cm stagnant water in rice field. • Spray mixture of streptocycline 1 g +copper oxychloride 20 g fungicides in 400-500 liter of water per hectare for control of bacterial leaf blight disease during rain free period. • To control blast disease, spray tricyclazole 75 WP 6 g or Iprobenfos 48 EC 10 ml in 10 liter of water at an interval of 15 to 20 days in clear weather condition.
KHARIF PEARL MILLET	Flowering		• Undertake the interculturing and weeding operations at 20 and 40 days after sowing under vapsa condition in sown crop. • Apply nitrogen 40 kg/ha as a top dressing after weeding and thinning under optimum soil moisture level at one month after sowing. • To control blast disease, carry out two spray of carbendazim 50 WP 10 g in 10 liters water at an interval of 15-20 days in clear sky condition. • Apply need based light irrigation by considering weather and soil moisture condition in standing crop.
FODDER CROPS	Vegetative / Flowering		• Apply need based light irrigation by considering weather and soil moisture condition in standing fodder crops. • Cutting of fodder sorghum should be done at 60 to 65 days (50 % flowering stage) after sowing in single cut varieties. • First cutting of fodder sorghum should be done at 45 to 45 days and next cutting at subsequent 30 days in multi cut varieties.
BIDI TOBACCO	Nursery/ Transplant ing		• Undertake weeding operation in nuresry. • To control worms or ants, apply chlorpyrifos @ 20 ml/15 liters of water at 10 days after sowing under clear and calm weather condition in the in the nursery. • Undertake transplanting of ready seedlings of bidi tobacco at spacing of 90 cm x 75 cm or 105 cm x 90 cm from third week of August to third week of September under assured irrigation facilities.
GREEN GRAM/ BLACK GRAM	Flowering/ Pod formation		• Mosaic of leaf curl disease is transmitted due to white fly so to control it, spray imidacloprid 17.8 SC 3 ml or acetamiprid 20 SP 2 g or azadirachtin 40 ml in 10 liter of water in clear sky condition. • If Aphid incidence is observed in green gram or black gram crop than undertake the spraying of imidacloprid 70 WS 7.5 g or thiamethoxam 70 WS 2.8 g in 10 liter of water in clear sky condition.

			• Apply need based light irrigation by considering weather and soil moisture condition in standing crops.
PIGEON PEA	Vegetative / Flowering		• Undertake interculturing, and hand weeding operations at 40 and 60 days after sowing under vapsa condition in sown crops. • Apply need based light irrigation by considering weather and soil moisture condition in standing crops.
SOYBEAN	Pod formation / Grain filling		• Use one light trap per hectare to attract and destroy the moths of armyworm. If a severe infestation of armyworm is observed, spray a mixture of Emamectin Benzoate 1.9 EC (8.5 ml) or Indoxacarb 14.5 SC (7 ml) per 10 liters of water during clear weather. • Apply need based light irrigation by considering weather and soil moisture condition in standing crop.
CHILLI/ TOMATO/ BRINJAL	Vegetative / Flowering		• Undertake interculturing, weeding, gap filling, etc operations at vapsa condition in transplanted field. • In continuous humid and cloudy weather, if thrips incidence is observed undertake spraying of neem seed oil 30 to 50 ml in 10 liters of water. In case of heavy attack, spray of spinosad 45 SC 3 ml or spinetoram 11.7 SC 10 ml in 10 liter of water in clear and calm weather condition in chilli crop. • In tomato crop, at initial incidence of Pinworm, install 40 yellow sticky trap per hectare. In heavy incidence, spray with Chlorantraniliprole 18.5 SC 3 ml or Flubendiamide 480 SC 3 ml in 10 liter of water in clear weather condition. • Due to continuous humid and cloudy weather condition, if sucking pest like white fly and jassid are observed in brinjal crop than undertake spraying of difenthiuron 50 WP 4 g or thiamethoxam 25 WG 4 g in 10 liters of water in clear and calm weather condition. • Apply need based light irrigation by considering weather and soil moisture condition in standing crops.
CASTOR	Vegetative		• Undertake interculturing, weeding, gap filling, etc operations at vapsa condition in sown crop. • Apply need based light irrigation by ensuring soil moisture in prevailing dry spell condition in sown castor crop.
OKRA	Vegetative / Flowering		• If yellow vein mosaic disease is observed in okra crop, than undertake the spraying of insecticides like tolfeprad 15 EC 20 ml or fenpropathrin 30 EC 3.4 ml in 10 liters of water in clear and calm weather condition. • Apply need based light irrigation by considering weather and soil moisture condition in standing crops.
LIVESTOCK			• Spray phenyl in the animal shed to avoid flies and mosquitos. For control of tick and other parasite, spray chemicals as per prescription of nearest animal clinic. • Give mineral mixture balance food to animals. Apply feed silage and hay material along with concentrate feed. Use chaff cutter for fodder. • Keep the fodder at safe place to avoid spoilage from rainfall.

Mahisagar

Main Crops	Stage	Disease/ Pest/ Management	Advisory
PAPAYA	Vegetative / Flowering		- Use bordeaux mixture paste (100:3:3) on stem for control of damping off disease. - Undertake the interculturing and hand weeding in vapsacondition papaya crop.
COTTON	Flowering/ Boll formation		- If irrigation facilities are available than apply need based irrigation through drip irrigation/alternate furrow irrigation method by considering soil moisture condition in standing crop. - Avoid application of fertilizers during long dry spell condition or it should apply under optimum soil moisture condition. Spray 2 % urea during dry spell in standing cotton crop. - In initial stage of incidence of sucking pest, spray neem seed extract 500 g or Neem oil 50 ml in 10 liters of water. In heavy incidence, spray flonicamid 50 WG 3 g or difenthiuron 50 WP 10 g in 10 liters of water in clear and calm weather condition. - If root rot and wilt disease is observed in cotton crop than undertake the drenching of fungicides like mancozeb 75 WP 27 g or copper oxychloride 40 g or carbendazim 10 g in 10 liter of water to infected plants and also apply urea or ammonium sulfate at 4 to 5 days after drenching of above fungicide.
KHARIF RICE	Flowering/ Grain filling		- Maintain 5-7 cm stagnant water in rice field. - Spray mixture of streptocycline 1 g +copper oxychloride 20 g fungicides in 400-500 liter of water per hectare for control of bacterial leaf blight disease during rain free period. - To control blast disease, spray tricyclazole 75 WP 6 g or Iprobenfos 48 EC 10 ml in 10 liter of water at an interval of 15 to 20 days in clear weather condition.
GROUDN UT	Flowering to Pegging		- Spray chlorothalonil 75 WP 27 g or Tebuconazole 10 ml/10 liter of water in clear and calm weather condition to prevent leaf spot infestation in groundnut. - Apply need based light irrigation by considering weather and soil moisture condition in standing crops.
KHARIF MAIZE	Flowering/ Tasseling		- Apply need based light irrigation by considering weather and soil moisture condition in standing crop. - Destroy the adults of fall armyworm by installing one light trap per hectare. Install 5 pheromone traps per hectare to attract male moth of this pest. Spray Neem seed powder 500 gm (5% Extract) or Neem based insecticide 20 ml (1 EC) to 40 ml (0.15 EC) in 10 liter of water in whole plant properly in clear weather condition. In heavy incidence, spray chlorantraniliprole 18.5 SC 4 ml or spinetoram 11.7 SC 10 ml in 10 liter of water to whole plant properly during clear weather condition. - Apply need based light irrigation by considering weather and soil moisture condition in standing crop.

SOYBEAN	Pod formation/ Grain filling		• Use one light trap per hectare to attract and destroy the moths of armyworm. If a severe infestation of armyworm is observed, spray a mixture of Emamectin Benzoate 1.9 EC (8.5 ml) or Indoxacarb 14.5 SC (7 ml) per 10 liters of water during clear weather. • Apply need based light irrigation by considering weather and soil moisture condition in standing crop.
KHARIF PEARL MILLET	Flowering		• To control blast disease, carry out two spray of carbendazim 50 WP 10 g in 10 liters water at an interval of 15-20 days in clear sky condition. • Apply need based light irrigation by considering weather and soil moisture condition in standing crops.
FODDER CROPS	Vegetative / Flowering		• Apply need based light irrigation by considering weather and soil moisture condition in standing fodder crops. • Cutting of fodder sorghum should be done at 60 to 65 days (50 % flowering stage) after sowing in single cut varieties. • First cutting of fodder sorghum should be done at 45 to 45 days and next cutting at subsequent 30 days in multi cut varieties.
BIDI TOBACCO	Nursery/ Transplanting		• Undertake weeding operation in nuresry. • To control worms or ants, apply chlorpyrifos @ 20 ml/15 liters of water at 10 days after sowing under clear and calm weather condition in the in the nursery. • Undertake transplanting of ready seedlings of bidi tobacco at spacing of 90 cm x 75 cm or 105 cm x 90 cm from third week of August to third week of September under assured irrigation facilities.
CHILLI/ TOMATO / BRINJAL	Vegetative / Flowering		• Undertake interculturing, weeding, gap filling, etc operations at vapsa condition in transplanted field. • In continuous humid and cloudy weather, if thrips incidence is observed undertake spraying of neem seed oil 30 to 50 ml in 10 liters of water. In case of heavy attack, spray of spinosad 45 SC 3 ml or spinetoram 11.7 SC 10 ml in 10 liter of water in clear and calm weather condition in chilli crop. • In tomato crop, at initial incidence of Pinworm, install 40 yellow sticky trap per hectare. In heavy incidence, spray with Chlorantraniliprole 18.5 SC 3 ml or Flubendiamide 480 SC 3 ml in 10 liter of water in clear weather condition. • Due to continuous humid and cloudy weather condition, if sucking pest like white fly and jassid are observed in brinjal crop then undertake spraying of difenthiuron 50 WP 4 g or thiamethoxam 25 WG 4 g in 10 liters of water in clear and calm weather condition. • Apply need based light irrigation by considering weather and soil moisture condition in standing crops.
PIGEON PEA	Vegetative / Flowering		• Undertake interculturing, and hand weeding operations at 40 and 60 days after sowing under vapsa condition in sown crops. • Apply need based light irrigation by considering weather and soil moisture condition in standing crops.
CHILLI/ TOMATO / BRINJAL	Vegetative		• Undertake interculturing, weeding, gap filling, etc operations at vapsa condition in transplanted field. • In continuous humid and cloudy weather, if thrips incidence is observed undertake spraying of neem seed oil 30 to 50 ml in 10 liters of water. In case of heavy attack, spray of spinosad 45 SC 3 ml or

			spinetoram 11.7 SC 10 ml in 10 liter of water in clear and calm weather condition in chilli crop. • Due to continuous humid and cloudy weather condition, if sucking pest like white fly and jassid are observed in brinjal crop than undertake spraying of difenthiuron 50 WP 4 g or thiamethoxam 25 WG 4 g in 10 liters of water in clear and calm weather condition. • Apply need based light irrigation by considering weather and soil moisture condition in standing crops.
CASTOR	Vegetative		• Undertake interculturing, weeding, gap filling, etc operations at vapsa condition in sown crop. • Apply need based light irrigation by ensuring soil moisture in prevailing dry spell condition in sown castor crop.
LIVESTOCK			• Spray phenyl in the animal shed to avoid flies and mosquitos. For control of tick and other parasite, spray chemicals as per prescription of nearest animal clinic. • Give mineral mixture balance food to animals. Apply feed silage and hay material along with concentrate feed. Use chaff cutter for fodder. • Keep the fodder at safe place to avoid spoilage from rainfall.

Panchmahal

Main Crops	Stage	Disease/ Pest/ Management	Advisory
COTTON	Flowering/ Boll formation		• If irrigation facilities are available than apply need based irrigation through drip irrigation/alternate furrow irrigation method by considering soil moisture condition in standing crop. • Avoid application of fertilizers during long dry spell condition or it should apply under optimum soil moisture condition. Spray 2 % urea during dry spell in standing cotton crop. • In initial stage of incidence of sucking pest, spray neem seed extract 500 g or Neem oil 50 ml in 10 liters of water. In heavy incidence, spray flonicamid 50 WG 3 g or difenthiuron 50 WP 10 g in 10 liters of water in clear and calm weather condition. • If root rot and wilt disease is observed in cotton crop than undertake the drenching of fungicides like mancozeb 75 WP 27 g or copper oxychloride 40 g or carbendazim 10 g in 10 liter of water to infected plants and also apply urea or ammonium sulfate at 4 to 5 days after drenching of above fungicide.
KHARIF RICE	Flowering/ Grain filling		• Maintain 5-7 cm stagnant water in rice field. • Spray mixture of streptocycline 1 g + copper oxychloride 20 g fungicides in 400-500 liter of water per hectare for control of bacterial leaf blight disease during rain free period. • To control blast disease, spray tricyclazole 75 WP 6 g or Iprobenfos 48 EC 10 ml in 10 liter of water at an interval of 15 to 20 days in clear weather condition.
KHARIF MAIZE	Flowering/ Tasseling		• Apply need based light irrigation by considering weather and soil moisture condition in standing crop. • Destroy the adults of fall armyworm by installing one light trap per

			hectare. Install 5 pheromone traps per hectare to attract male moth of this pest. Spray Neem seed powder 500 gm (5% Extract) or Neem based insecticide 20 ml (1 EC) to 40 ml (0.15 EC) in 10 liter of water in whole plant properly in clear weather condition. In heavy incidence, spray chlorantraniliprole 18.5 SC 4 ml or spinetoram 11.7 SC 10 ml in 10 liter of water to whole plant properly during clear weather condition. • Apply need based light irrigation by considering weather and soil moisture condition in standing crop.
SOYBEAN	Pod formation/ Grain filling		• Use one light trap per hectare to attract and destroy the moths of armyworm. If a severe infestation of armyworm is observed, spray a mixture of Emamectin Benzoate 1.9 EC (8.5 ml) or Indoxacarb 14.5 SC (7 ml) per 10 liters of water during clear weather. • Apply need based light irrigation by considering weather and soil moisture condition in standing crop.
KHARIF PEARL MILLET	Flowering		• To control blast disease, carry out two spray of carbendazim 50 WP 10 g in 10 liters water at an interval of 15-20 days in clear sky condition. • Apply need based light irrigation by considering weather and soil moisture condition in standing crop.
FODDER CROPS	Vegetative / Flowering		• Apply need based light irrigation by considering weather and soil moisture condition in standing fodder crops. • Cutting of fodder sorghum should be done at 60 to 65 days (50 % flowering stage) after sowing in single cut varieties. • First cutting of fodder sorghum should be done at 45 to 45 days and next cutting at subsequent 30 days in multi cut varieties.
CHILLI/TOMATO / BRINJAL	Vegetative / Flowering		• Undertake interculturing, weeding, gap filling, etc operations at vapsa condition in transplanted field. • In continuous humid and cloudy weather, if thrips incidence is observed undertake spraying of neem seed oil 30 to 50 ml in 10 liters of water. In case of heavy attack, spray of spinosad 45 SC 3 ml or spinetoram 11.7 SC 10 ml in 10 liter of water in clear and calm weather condition in chilli crop. • In tomato crop, at initial incidence of Pinworm, install 40 yellow sticky trap per hectare. In heavy incidence, spray with Chlorantraniliprole 18.5 SC 3 ml or Flubendiamide 480 SC 3 ml in 10 liter of water in clear weather condition. • Due to continuous humid and cloudy weather condition, if sucking pest like white fly and jassid are observed in brinjal crop then undertake spraying of difenthiuron 50 WP 4 g or thiamethoxam 25 WG 4 g in 10 liters of water in clear and calm weather condition. • Apply need based light irrigation by considering weather and soil moisture condition in standing crops.
PIGEON PEA	Vegetative / Flowering		• Undertake interculturing, and hand weeding operations at 40 and 60 days after sowing under vapsa condition in sown crops. • Apply need based light irrigation by considering weather and soil moisture condition in standing crops.
CASTOR	Vegetative		• Undertake interculturing, weeding, gap filling, etc operations at vapsa condition in sown crop. • Apply need based light irrigation by ensuring soil moisture in prevailing dry spell condition in sown castor crop.

LIVESTOCK	- Spray phenyl in the animal shed to avoid flies and mosquitos. For control of tick and other parasite, spray chemicals as per prescription of nearest animal clinic. - Give mineral mixture balance food to animals. Apply feed silage and hay material along with concentrate feed. Use chaff cutter for fodder. - Keep the fodder at safe place to avoid spoilage from rainfall.
------------------	---

Vadodara

Main Crops	Stage	Disease/ Pest/ Management	Advisory
BANANA	Vegetative		- Undertake the interculturing and weeding at vapsa condition in planted crop. - Apply need based light irrigation by considering weather and soil moisture condition in standing crop. - Undertake earthing up up to 15-20 cm height around root zone and frequently remove suckers around mother plant.
BANANA	Vegetative		- Undertake the interculturing and weeding at vapsa condition in planted crop. - Apply need based light irrigation by considering weather and soil moisture condition in standing crop. - Undertake earthing up up to 15-20 cm height around root zone and frequently remove suckers around mother plant.
KHARIF RICE	Flowering / Grain filling		- Maintain 5-7 cm stagnant water in rice field. - Spray mixture of streptocycline 1 g +copper oxychloride 20 g fungicides in 400-500 liter of water per hectare for control of bacterial leaf blight disease during rain free period. - To control blast disease, spray tricyclazole 75 WP 6 g or Iprobenfos 48 EC 10 ml in 10 liter of water at an interval of 15 to 20 days in clear weather condition.
GREEN GRAM/ BLACK GRAM	Flowering / Pod formation		- Mosaic of leaf curl disease is transmitted due to white fly so to control it, spray imidacloprid 17.8 SC 3 ml or acetamiprid 20 SP 2 g or azadirachtin 40 ml in 10 liter of water in clear sky condition. - If Aphid incidence is observed in green gram or black gram crop then undertake the spraying of imidacloprid 70 WS 7.5 g or thiamethoxam 70 WS 2.8 g in 10 liter of water in clear sky condition. - Apply need based light irrigation by considering weather and soil moisture condition in standing crops.
PIGEON PEA	Vegetative / Flowering		- Undertake interculturing, and hand weeding operations at 40 and 60 days after sowing under vapsa condition in sown crops. - Apply need based light irrigation by considering weather and soil moisture condition in standing crops.
SOYBEAN	Pod formation / Grain filling		- Use one light trap per hectare to attract and destroy the moths of armyworm. If a severe infestation of armyworm is observed, spray a mixture of Emamectin Benzoate 1.9 EC (8.5 ml) or Indoxacarb 14.5 SC (7 ml) per 10 liters of water during clear weather. - Apply need based light irrigation by considering weather and soil moisture condition in standing crop.

KHARIF MAIZE	Flowering / Tasseling		• Apply need based light irrigation by considering weather and soil moisture condition in standing crop. • Destroy the adults of fall armyworm by installing one light trap per hectare. Install 5 pheromone traps per hectare to attract male moth of this pest. Spray Neem seed powder 500 gm (5% Extract) or Neem based insecticide 20 ml (1 EC) to 40 ml (0.15 EC) in 10 liter of water in whole plant properly in clear weather condition. In heavy incidence, spray chlorantraniliprole 18.5 SC 4 ml or spinetoram 11.7 SC 10 ml in 10 liter of water to whole plant properly during clear weather condition. • Apply need based light irrigation by considering weather and soil moisture condition in standing crop.
CHILLI/ TOMATO / BRINJAL	Vegetative/ Flowering		• Undertake interculturing, weeding, gap filling, etc operations at vapsa condition in transplanted field. • In continuous humid and cloudy weather, if thrips incidence is observed undertake spraying of neem seed oil 30 to 50 ml in 10 liters of water. In case of heavy attack, spray of spinosad 45 SC 3 ml or spinetoram 11.7 SC 10 ml in 10 liter of water in clear and calm weather condition in chilli crop. • In tomato crop, at initial incidence of Pinworm, install 40 yellow sticky trap per hectare. In heavy incidence, spray with Chlorantraniliprole 18.5 SC 3 ml or Flubendiamide 480 SC 3 ml in 10 liter of water in clear weather condition. • Due to continuous humid and cloudy weather condition, if sucking pest like white fly and jassid are observed in brinjal crop then undertake spraying of difenthiuron 50 WP 4 g or thiamethoxam 25 WG 4 g in 10 liters of water in clear and calm weather condition. • Apply need based light irrigation by considering weather and soil moisture condition in standing crops.
CASTOR	Vegetative		• Undertake interculturing, weeding, gap filling, etc operations at vapsa condition in sown crop. • Apply need based light irrigation by ensuring soil moisture in prevailing dry spell condition in sown castor crop.
LIVESTOCK			• Spray phenyl in the animal shed to avoid flies and mosquitos. For control of tick and other parasite, spray chemicals as per prescription of nearest animal clinic. • Give mineral mixture balance food to animals. Apply feed silage and hay material along with concentrate feed. Use chaff cutter for fodder. • Keep the fodder at safe place to avoid spoilage from rainfall.

AMFU Dantiwada:**ARAVALLI**

Agricultural Advisory			
Major Crops	Stage	Disease/Pest Management	Advisory
Castor	Growth	Semi-looper and Castor hairy caterpillar	➤ If there has been no rainfall in the region, apply irrigation to the standing crop. ➤ Apply Malathion 50 EC @ 60 ml/15 lit or Emamectin benzoate 5 wg @ 7.5 g/15 lit or Chlorpyrifos 50 ec + cypermethrin 5 ec @ 15 ml/15 lit for control of Castor hairy caterpillar and Semi-looper clear sky conditions.
Hybrid/ B.T. cotton	Vegetative /Branching stage	Pinkball worm/ Leaf Spot/Root rot	➤ If there has been no rainfall in the region, apply irrigation to the standing crop. ➤ Avoid applying fertilizers during prolonged dry spells; apply only when soil moisture is adequate during dry spells, spray a 2% urea solution directly onto standing cotton crops. ➤ Install pink bollworm pheromone traps. ➤ For the control of pink bollworm, spray 10ml deltamethrin OR trizophos in 10 litre of water OR spray 100 gm Beauveria bassiana powder and 20 ml prophenophos mix in 10 liter of water. ➤ At the initiation of the root rot disease drench copper Oxychloride (50% WP) 0.2 % solution (40g/ 10 litre water) around stem of the plants in rainless period.
Groundnut (Kharif)	Vegetative stage /Flowering	Yellowing of Leaves/ Weed/ Aphids	➤ If there has been no rainfall in the region, apply irrigation to the standing crop. ➤ Apply Ammonium sulphate fertilizer 30 kg/ha for correction of yellowing. ➤ To prevent leaf spot disease infestation spray Tebuconazole 18.3 % SC 10 in 10 liters of water as precautionary measure under clear sky. ➤ Spray ferrous sulfate 100 gram & citric acid 10 gram/10 litre of water to cure yellowing of groundnut crop. ➤ Monitor groundnut crop for infestation of aphids. If infestation found above ETL, then spray Imidacloprid 17.8 % EC 5 ml in 10 liter of water.
		White Grub/Rust, Helicoverpa and Spodoptera	➤ To spraying white grubs harbouring host plant with chlorpyrifos 20 EC 0.04% (20 ml/10 l water). ➤ To Collection and destruction of White grub adults from host plant around the field.

			➤ Two sprays of pongamia oil (30 ml/10 l of water) OR ponneem (30 ml/10 l of water) at 15 days interval starting from pest infestation for effective and economical management of defoliators (Helicoverpa and Spodoptera). ➤ For the control of rust, spray 5 ml Tridemorph in 10 litres of water.
Pearl millet (Kharif)	Vegetative stage	Interculturing/ Weeding	➤ Apply irrigation to the standing crop as per the crop requirement. ➤ Apply supplementary dose of 87 kg/ha urea to pearl millet at tillering stage under clear sky.
		Shootfly	➤ Two sprays of Endosulfan (0.07%) at 10 days and 20 days after germination for shootfly control.
Sesame	Vegetative stage	Leaf blight/ Stem rot/ Leaf webber	➤ If there has been no rainfall in the region, apply irrigation to the standing crop. ➤ If infestation of leaf Webber observe above ETL, spray Cartep Hydrochloride 15 gram in 10 litres of water OR Neem Seed Kernel Extract (NSKE) 3% (300g /10 l water). ➤ To control leaf blight and stem rot, spray Copper oxychloride 50 % WP 40 g/10 lit or Mancozeb 75 % WP 25 ml/10 lit. ➤ Apply basal dose of Nitrogen @ 25 kg/ha and phosphorus @ 40 kg/ha.
Maize	Vegetative stage	Interculturing/ Weeding	➤ Apply need based light irrigation by considering weather and soil moisture condition in standing crops.
Vegetable crops	Flowering /Fruits/ Vegetative stage	Interculturing/ Weeding	➤ If there has been no rainfall in the region, apply irrigation to the standing crop. ➤ Ensure weed-free cultivation of vegetable crops, and regularly send graded produce to the market. ➤ It is also advised to use shed nets, which helps in reducing vector borne diseases and produce healthy seedlings. ➤ To prevent soil borne disease drenching of nursery bed should be done with Mencozeb 75 % WP @ 2.5 gm/ litre of water after 15 DAS. ➤ Spray Quinalphos 20 ml in 10 liters of water for control of leaf miner.
		Thrips/ White fly	➤ For control of thrips, spray Profenofos 20 ml with 500 ml of 5 % neem seed extract or neem seed oil 50 ml in 10 litres of water. In case of severe attack, spray Spinosad or Spinetoram 3 ml per 10 liter of water. ➤ To minimize population of whitefly, Spray 500 ml

			of 5 % neem seed extract or neem oil 50 ml with Difenthiuron 50 WP @ 16 gm or Trizophos 40 EC 25 ml / 10 lit of water.
Cattle	➤ Feed animals with a mixture of green grass + hay + minerals + dry feed. ➤ Ensure adequate clean drinking water availability at all times and increase watering frequency during dry spell. ➤ Avoid overcrowding in sheds; ensure proper ventilation. ➤ Udder of milking animals must be properly clean with zinc oxide or boric powder. ➤ Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Deworming, T.B. and J.D. ➤ Keep fodder in safe place to avoid spoilage due to rainfall. ➤ Vaccination: During the monsoon season, it is recommended to vaccinate cattle against Haemorrhagic Septicaemia.		

BANASKANTHA

Agricultural Advisory			
Major Crops	Stage	Disease/Pest Management	Advisory
Castor	Growth	Semi-looper and Castor hairy caterpillar	➤ If there has been no rainfall in the region, apply irrigation to the standing crop. ➤ Apply Malathion 50 EC @ 60 ml/15 lit or Emamectin benzoate 5 wg @ 7.5 g/15 lit or Chlorpyrifos 50 ec + cypermethrin 5 ec @ 15 ml/15 lit for control of Castor hairy caterpillar and Semi-looper clear sky conditions.
Hybrid/ B.T. cotton	Vegetative /Branching stage	Pinkball worm/ Leaf Spot/Root rot	➤ If there has been no rainfall in the region, apply irrigation to the standing crop. ➤ Avoid applying fertilizers during prolonged dry spells; apply only when soil moisture is adequate during dry spells, spray a 2% urea solution directly onto standing cotton crops. ➤ Install pink bollworm pheromone traps. ➤ For the control of pink bollworm, spray 10ml deltamethrin OR trizophos in 10 litre of water OR spray 100 gm Beauveria bassiana powder and 20 ml prophenophos mix in 10 liter of water. ➤ At the initiation of the root rot disease drench copper Oxychloride (50% WP) 0.2 % solution (40g/ 10 litre water) around stem of the plants in rainless period.

Groundnut (Kharif)	Vegetative stage /Flowering	Yellowing of Leaves/ Weed/ Aphids	➤ If there has been no rainfall in the region, apply irrigation to the standing crop. ➤ Apply Ammonium sulphate fertilizer 30 kg/ha for correction of yellowing. ➤ To prevent leaf spot disease infestation spray Tebuconazole 18.3 % SC 10 in 10 liters of water as precautionary measure under clear sky. ➤ Spray ferrous sulfate 100 gram & citric acid 10 gram/10 litre of water to cure yellowing of groundnut crop. ➤ Monitor groundnut crop for infestation of aphids. If infestation found above ETL, then spray Imidacloprid 17.8 % EC 5 ml in 10 liter of water.
		White Grub/Rust, Helicoverpa and Spodoptera	➤ To spraying white grubs harbouring host plant with chlorpyrifos 20 EC 0.04% (20 ml/10 l water). ➤ To Collection and destruction of White grub adults from host plant around the field. ➤ Two sprays of pongamia oil (30 ml/10 l of water) OR ponneem (30 ml/10 l of water) at 15 days interval starting from pest infestation for effective and economical management of defoliators (Helicoverpa and Spodoptera). ➤ For the control of rust, spray 5 ml Tridemorph in 10 litres of water.
Pearl millet (Kharif)	Vegetative stage	Interculturing/ Weeding	➤ Apply irrigation to the standing crop as per the crop requirement. ➤ Apply supplementary dose of 87 kg/ha urea to pearl millet at tillering stage under clear sky.
		Shootfly	➤ Two sprays of Endosulfan (0.07%) at 10 days and 20 days after germination for shootfly control.
Sesame	Vegetative stage	Leaf blight/ Stem rot/ Leaf webber	➤ If there has been no rainfall in the region, apply irrigation to the standing crop. ➤ If infestation of leaf Webber observe above ETL, spray Cartep Hydrochloride 15 gram in 10 litres of water OR Neem Seed Kernel Extract (NSKE) 3% (300g /10 l water). ➤ To control leaf blight and stem rot, spray Copper oxychloride 50 % WP 40 g/10 lit or Mancozeb 75 % WP 25 ml/10 lit. ➤ Apply basal dose of Nitrogen @ 25 kg/ha and phosphorus @ 40 kg/ha.
Maize	Vegetative stage	Interculturing/ Weeding	➤ Apply need based light irrigation by considering weather and soil moisture condition in standing crops.

Vegetable crops	Flowering /Fruits/ Vegetative stage	Interculturing/ Weeding	➤ If there has been no rainfall in the region, apply irrigation to the standing crop. ➤ Ensure weed-free cultivation of vegetable crops, and regularly send graded produce to the market. ➤ It is also advised to use shed nets, which helps in reducing vector borne diseases and produce healthy seedlings. ➤ To prevent soil borne disease drenching of nursery bed should be done with Mencozeb 75 % WP @ 2.5 gm/ litre of water after 15 DAS. ➤ Spray Quinalphos 20 ml in 10 liters of water for control of leaf miner.
		Thrips/ White fly	➤ For control of thrips, spray Profenofos 20 ml with 500 ml of 5 % neem seed extract or neem seed oil 50 ml in 10 litres of water. In case of severe attack, spray Spinosad or Spinetoram 3 ml per 10 liter of water. ➤ To minimize population of whitefly, Spray 500 ml of 5 % neem seed extract or neem oil 50 ml with Difenthiuron 50 WP @ 16 gm or Trizophos 40 EC 25 ml / 10 lit of water.
Cattle	➤ Feed animals with a mixture of green grass + hay + minerals + dry feed. ➤ Ensure adequate clean drinking water availability at all times and increase watering frequency during dry spell. ➤ Avoid overcrowding in sheds; ensure proper ventilation. ➤ Udder of milking animals must be properly clean with zinc oxide or boric powder. ➤ Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Deworming, T.B. and J.D. ➤ Keep fodder in safe place to avoid spoilage due to rainfall. ➤ Vaccination: During the monsoon season, it is recommended to vaccinate cattle against Haemorrhagic Septicaemia.		

GANDHINAGAR

Agricultural Advisory			
Major Crops	Stage	Disease/Pest Management	Advisory
Castor	Growth	Semi-looper and Castor hairy caterpillar	➤ If there has been no rainfall in the region, apply irrigation to the standing crop. ➤ Apply Malathion 50 EC @ 60 ml/15 lit or Emamectin benzoate 5 wg @ 7.5 g/15 lit or Chlorpyrifos 50 ec + cypermethrin 5 ec @ 15 ml/15 lit for control of Castor hairy caterpillar and Semi-looper clear sky conditions.

Hybrid/ B.T. cotton	Vegetative /Branching stage	Pinkball worm/ Leaf Spot/Root rot	➤ If there has been no rainfall in the region, apply irrigation to the standing crop. ➤ Avoid applying fertilizers during prolonged dry spells; apply only when soil moisture is adequate during dry spells, spray a 2% urea solution directly onto standing cotton crops. ➤ Install pink bollworm pheromone traps. ➤ For the control of pink bollworm, spray 10ml deltamethrin OR trizophos in 10 litre of water OR spray 100 gm Beauveria bassiana powder and 20 ml prophenophos mix in 10 liter of water. ➤ At the initiation of the root rot disease drench copper Oxychloride (50% WP) 0.2 % solution (40g/ 10 litre water) around stem of the plants in rainless period.
Groundnut (Kharif)	Vegetative stage /Flowering	Yellowing of Leaves/ Weed/ Aphids	➤ If there has been no rainfall in the region, apply irrigation to the standing crop. ➤ Apply Ammonium sulphate fertilizer 30 kg/ha for correction of yellowing. ➤ To prevent leaf spot disease infestation spray Tebuconazole 18.3 % SC 10 in 10 liters of water as precautionary measure under clear sky. ➤ Spray ferrous sulfate 100 gram & citric acid 10 gram/10 litre of water to cure yellowing of groundnut crop. ➤ Monitor groundnut crop for infestation of aphids. If infestation found above ETL, then spray Imidacloprid 17.8 % EC 5 ml in 10 liter of water.
		White Grub/Rust, Helicoverpa and Spodoptera	➤ To spraying white grubs harbouring host plant with chlorpyrifos 20 EC 0.04% (20 ml/10 l water). ➤ To Collection and destruction of White grub adults from host plant around the field. ➤ Two sprays of pongamia oil (30 ml/10 l of water) OR ponneem (30 ml/10 l of water) at 15 days interval starting from pest infestation for effective and economical management of defoliators (Helicoverpa and Spodoptera). ➤ For the control of rust, spray 5 ml Tridemorph in 10 litres of water.
Pearl millet (Kharif)	Vegetative stage	Interculturing/ Weeding	➤ Apply irrigation to the standing crop as per the crop requirement. ➤ Apply supplementary dose of 87 kg/ha urea to pearl millet at tillering stage under clear sky.
		Shootfly	➤ Two sprays of Endosulfan (0.07%) at 10 days and 20 days after germination for shootfly control.

Sesame	Vegetative stage	Leaf blight/ Stem rot/ Leaf webber	➤ If there has been no rainfall in the region, apply irrigation to the standing crop. ➤ If infestation of leaf Webber observe above ETL, spray Cartep Hydrochloride 15 gram in 10 litres of water OR Neem Seed Kernel Extract (NSKE) 3% (300g /10 l water). ➤ To control leaf blight and stem rot, spray Copper oxychloride 50 % WP 40 g/10 lit or Mancozeb 75 % WP 25 ml/10 lit. ➤ Apply basal dose of Nitrogen @ 25 kg/ha and phosphorus @ 40 kg/ha.
Maize	Vegetative stage	Interculturing/ Weeding	➤ Apply need based light irrigation by considering weather and soil moisture condition in standing crops.
Vegetable crops	Flowering /Fruits/ Vegetative stage	Interculturing/ Weeding	➤ If there has been no rainfall in the region, apply irrigation to the standing crop. ➤ Ensure weed-free cultivation of vegetable crops, and regularly send graded produce to the market. ➤ It is also advised to use shed nets, which helps in reducing vector borne diseases and produce healthy seedlings. ➤ To prevent soil borne disease drenching of nursery bed should be done with Mencozeb 75 % WP @ 2.5 gm/ litre of water after 15 DAS. ➤ Spray Quinalphos 20 ml in 10 liters of water for control of leaf miner.
		Thrips/ White fly	➤ For control of thrips, spray Profenofos 20 ml with 500 ml of 5 % neem seed extract or neem seed oil 50 ml in 10 litres of water. In case of severe attack, spray Spinosad or Spinetoram 3 ml per 10 liter of water. ➤ To minimize population of whitefly, Spray 500 ml of 5 % neem seed extract or neem oil 50 ml with Difenthiuron 50 WP @ 16 gm or Trizophos 40 EC 25 ml / 10 lit of water.
Cattle	➤ Feed animals with a mixture of green grass + hay + minerals + dry feed. ➤ Ensure adequate clean drinking water availability at all times and increase watering frequency during dry spell. ➤ Avoid overcrowding in sheds; ensure proper ventilation. ➤ Udder of milking animals must be properly clean with zinc oxide or boric powder. ➤ Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Deworming, T.B. and J.D. ➤ Keep fodder in safe place to avoid spoilage due to rainfall. ➤ Vaccination: During the monsoon season, it is recommended to vaccinate cattle against Haemorrhagic Septicaemia.		

MEHSANA

Agricultural Advisory			
Major Crops	Stage	Disease/Pest Management	Advisory
Castor	Growth	Semi-looper and Castor hairy caterpillar	➤ If there has been no rainfall in the region, apply irrigation to the standing crop. ➤ Apply Malathion 50 EC @ 60 ml/15 lit or Emamectin benzoate 5 wg @ 7.5 g/15 lit or Chlorpyrifos 50 ec + cypermethrin 5 ec @ 15 ml/15 lit for control of Castor hairy caterpillar and Semi-looper clear sky conditions.
Hybrid/ B.T. cotton	Vegetative /Branching stage	Pinkball worm/ Leaf Spot/Root rot	➤ If there has been no rainfall in the region, apply irrigation to the standing crop. ➤ Avoid applying fertilizers during prolonged dry spells; apply only when soil moisture is adequate during dry spells, spray a 2% urea solution directly onto standing cotton crops. ➤ Install pink bollworm pheromone traps. ➤ For the control of pink bollworm, spray 10ml deltamethrin OR trizophos in 10 litre of water OR spray 100 gm Beauveria bassiana powder and 20 ml prophenophos mix in 10 liter of water. ➤ At the initiation of the root rot disease drench copper Oxchloride (50% WP) 0.2 % solution (40g/ 10 litre water) around stem of the plants in rainless period.
Groundnut (Kharif)	Vegetative stage /Flowering	Yellowing of Leaves/ Weed/ Aphids	➤ If there has been no rainfall in the region, apply irrigation to the standing crop. ➤ Apply Ammonium sulphate fertilizer 30 kg/ha for correction of yellowing. ➤ To prevent leaf spot disease infestation spray Tebuconazole 18.3 % SC 10 in 10 liters of water as precautionary measure under clear sky. ➤ Spray ferrous sulfate 100 gram & citric acid 10 gram/10 litre of water to cure yellowing of groundnut crop. ➤ Monitor groundnut crop for infestation of aphids. If infestation found above ETL, then spray Imidacloprid 17.8 % EC 5 ml in 10 liter of water.
		White Grub/Rust, Helicoverpa and Spodoptera	➤ To spraying white grubs harbouring host plant with chlorpyrifos 20 EC 0.04% (20 ml/10 l water). ➤ To Collection and destruction of White grub adults from host plant around the field. ➤ Two sprays of pongamia oil (30 ml/10 l of water)

			OR ponneem (30 ml/10 l of water) at 15 days interval starting from pest infestation for effective and economical management of defoliators (Helicoverpa and Spodoptera). ➤ For the control of rust, spray 5 ml Tridemorph in 10 litres of water.
Pearl millet (Kharif)	Vegetative stage	Interculturing/ Weeding	➤ Apply irrigation to the standing crop as per the crop requirement. ➤ Apply supplementary dose of 87 kg/ha urea to pearl millet at tillering stage under clear sky.
		Shootfly	➤ Two sprays of Endosulfan (0.07%) at 10 days and 20 days after germination for shootfly control.
Sesame	Vegetative stage	Leaf blight/ Stem rot/ Leaf webber	➤ If there has been no rainfall in the region, apply irrigation to the standing crop. ➤ If infestation of leaf Webber observe above ETL, spray Cartep Hydrochloride 15 gram in 10 litres of water OR Neem Seed Kernel Extract (NSKE) 3% (300g /10 l water). ➤ To control leaf blight and stem rot, spray Copper oxychloride 50 % WP 40 g/10 lit or Mancozeb 75 % WP 25 ml/10 lit. ➤ Apply basal dose of Nitrogen @ 25 kg/ha and phosphorus @ 40 kg/ha.
Maize	Vegetative stage	Interculturing/ Weeding	➤ Apply need based light irrigation by considering weather and soil moisture condition in standing crops.
Vegetable crops	Flowering /Fruits/ Vegetative stage	Interculturing/ Weeding	➤ If there has been no rainfall in the region, apply irrigation to the standing crop. ➤ Ensure weed-free cultivation of vegetable crops, and regularly send graded produce to the market. ➤ It is also advised to use shed nets, which helps in reducing vector borne diseases and produce healthy seedlings. ➤ To prevent soil borne disease drenching of nursery bed should be done with Mencozeb 75 % WP @ 2.5 gm/ litre of water after 15 DAS. ➤ Spray Quinalphos 20 ml in 10 liters of water for control of leaf miner.
		Thrips/ White fly	➤ For control of thrips, spray Profenofos 20 ml with 500 ml of 5 % neem seed extract or neem seed oil 50 ml in 10 litres of water. In case of severe attack, spray Spinosad or Spinetoram 3 ml per 10 liter of water. ➤ To minimize population of whitefly, Spray 500 ml of 5 % neem seed extract or neem oil 50 ml with

			Difenthiuron 50 WP @ 16 gm or Trizophos 40 EC 25 ml / 10 lit of water.
Cattle	➤ Feed animals with a mixture of green grass + hay + minerals + dry feed. ➤ Ensure adequate clean drinking water availability at all times and increase watering frequency during dry spell. ➤ Avoid overcrowding in sheds; ensure proper ventilation. ➤ Udder of milking animals must be properly clean with zinc oxide or boric powder. ➤ Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Deworming, T.B. and J.D. ➤ Keep fodder in safe place to avoid spoilage due to rainfall. ➤ Vaccination: During the monsoon season, it is recommended to vaccinate cattle against Haemorrhagic Septicaemia.		

PATAN

Agricultural Advisory			
Major Crops	Stage	Disease/Pest Management	Advisory
Castor	Growth	Semi-looper and Castor hairy caterpillar	➤ If there has been no rainfall in the region, apply irrigation to the standing crop. ➤ Apply Malathion 50 EC @ 60 ml/15 lit or Emamectin benzoate 5 wg @ 7.5 g/15 lit or Chlorpyrifos 50 ec + cypermethrin 5 ec @ 15 ml/15 lit for control of Castor hairy caterpillar and Semi-looper clear sky conditions.
Hybrid/ B.T. cotton	Vegetative /Branching stage	Pinkball worm/ Leaf Spot/Root rot	➤ If there has been no rainfall in the region, apply irrigation to the standing crop. ➤ Avoid applying fertilizers during prolonged dry spells; apply only when soil moisture is adequate during dry spells, spray a 2% urea solution directly onto standing cotton crops. ➤ Install pink bollworm pheromone traps. ➤ For the control of pink bollworm, spray 10ml deltamethrin OR trizophos in 10 litre of water OR spray 100 gm Beauveria bassiana powder and 20 ml prophenophos mix in 10 liter of water. ➤ At the initiation of the root rot disease drench copper Oxychloride (50% WP) 0.2 % solution (40g/ 10 litre water) around stem of the plants in rainless period.

Groundnut (Kharif)	Vegetative stage /Flowering	Yellowing of Leaves/ Weed/ Aphids	➤ If there has been no rainfall in the region, apply irrigation to the standing crop. ➤ Apply Ammonium sulphate fertilizer 30 kg/ha for correction of yellowing. ➤ To prevent leaf spot disease infestation spray Tebuconazole 18.3 % SC 10 in 10 liters of water as precautionary measure under clear sky. ➤ Spray ferrous sulfate 100 gram & citric acid 10 gram/10 litre of water to cure yellowing of groundnut crop. ➤ Monitor groundnut crop for infestation of aphids. If infestation found above ETL, then spray Imidacloprid 17.8 % EC 5 ml in 10 liter of water.
		White Grub/Rust, Helicoverpa and Spodoptera	➤ To spraying white grubs harbouring host plant with chlorpyrifos 20 EC 0.04% (20 ml/10 l water). ➤ To Collection and destruction of White grub adults from host plant around the field. ➤ Two sprays of pongamia oil (30 ml/10 l of water) OR ponneem (30 ml/10 l of water) at 15 days interval starting from pest infestation for effective and economical management of defoliators (Helicoverpa and Spodoptera). ➤ For the control of rust, spray 5 ml Tridemorph in 10 litres of water.
Pearl millet (Kharif)	Vegetative stage	Interculturing/ Weeding	➤ Apply irrigation to the standing crop as per the crop requirement. ➤ Apply supplementary dose of 87 kg/ha urea to pearl millet at tillering stage under clear sky.
		Shootfly	➤ Two sprays of Endosulfan (0.07%) at 10 days and 20 days after germination for shootfly control.
Sesame	Vegetative stage	Leaf blight/ Stem rot/ Leaf webber	➤ If there has been no rainfall in the region, apply irrigation to the standing crop. ➤ If infestation of leaf Webber observe above ETL, spray Cartep Hydrochloride 15 gram in 10 litres of water OR Neem Seed Kernel Extract (NSKE) 3% (300g /10 l water). ➤ To control leaf blight and stem rot, spray Copper oxychloride 50 % WP 40 g/10 lit or Mancozeb 75 % WP 25 ml/10 lit. ➤ Apply basal dose of Nitrogen @ 25 kg/ha and phosphorus @ 40 kg/ha.
Maize	Vegetative stage	Interculturing/ Weeding	➤ Apply need based light irrigation by considering weather and soil moisture condition in standing crops.

Vegetable crops	Flowering /Fruits/ Vegetative stage	Interculturing/ Weeding	➤ If there has been no rainfall in the region, apply irrigation to the standing crop. ➤ Ensure weed-free cultivation of vegetable crops, and regularly send graded produce to the market. ➤ It is also advised to use shed nets, which helps in reducing vector borne diseases and produce healthy seedlings. ➤ To prevent soil borne disease drenching of nursery bed should be done with Mencozeb 75 % WP @ 2.5 gm/ litre of water after 15 DAS. ➤ Spray Quinalphos 20 ml in 10 liters of water for control of leaf miner.
		Thrips/ White fly	➤ For control of thrips, spray Profenofos 20 ml with 500 ml of 5 % neem seed extract or neem seed oil 50 ml in 10 litres of water. In case of severe attack, spray Spinosad or Spinetoram 3 ml per 10 liter of water. ➤ To minimize population of whitefly, Spray 500 ml of 5 % neem seed extract or neem oil 50 ml with Difenthiuron 50 WP @ 16 gm or Trizophos 40 EC 25 ml / 10 lit of water.
Cattle	➤ Feed animals with a mixture of green grass + hay + minerals + dry feed. ➤ Ensure adequate clean drinking water availability at all times and increase watering frequency during dry spell. ➤ Avoid overcrowding in sheds; ensure proper ventilation. ➤ Udder of milking animals must be properly clean with zinc oxide or boric powder. ➤ Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Deworming, T.B. and J.D. ➤ Keep fodder in safe place to avoid spoilage due to rainfall. ➤ Vaccination: During the monsoon season, it is recommended to vaccinate cattle against Haemorrhagic Septicaemia.		

SABARKANTHA

Agricultural Advisory			
Major Crops	Stage	Disease/Pest Management	Advisory
Castor	Growth	Semi-looper and Castor hairy caterpillar	➤ If there has been no rainfall in the region, apply irrigation to the standing crop. ➤ Apply Malathion 50 EC @ 60 ml/15 lit or Emamectin benzoate 5 wg @ 7.5 g/15 lit or Chlorpyrifos 50 ec + cypermethrin 5 ec @ 15 ml/15 lit for control of Castor hairy caterpillar and Semi-looper clear sky conditions.

Hybrid/ B.T. cotton	Vegetative /Branching stage	Pinkball worm/ Leaf Spot/Root rot	➤ If there has been no rainfall in the region, apply irrigation to the standing crop. ➤ Avoid applying fertilizers during prolonged dry spells; apply only when soil moisture is adequate during dry spells, spray a 2% urea solution directly onto standing cotton crops. ➤ Install pink bollworm pheromone traps. ➤ For the control of pink bollworm, spray 10ml deltamethrin OR trizophos in 10 litre of water OR spray 100 gm Beauveria bassiana powder and 20 ml prophenophos mix in 10 liter of water. ➤ At the initiation of the root rot disease drench copper Oxychloride (50% WP) 0.2 % solution (40g/ 10 litre water) around stem of the plants in rainless period.
Groundnut (Kharif)	Vegetative stage /Flowering	Yellowing of Leaves/ Weed/ Aphids	➤ If there has been no rainfall in the region, apply irrigation to the standing crop. ➤ Apply Ammonium sulphate fertilizer 30 kg/ha for correction of yellowing. ➤ To prevent leaf spot disease infestation spray Tebuconazole 18.3 % SC 10 in 10 liters of water as precautionary measure under clear sky. ➤ Spray ferrous sulfate 100 gram & citric acid 10 gram/10 litre of water to cure yellowing of groundnut crop. ➤ Monitor groundnut crop for infestation of aphids. If infestation found above ETL, then spray Imidacloprid 17.8 % EC 5 ml in 10 liter of water.
		White Grub/Rust, Helicoverpa and Spodoptera	➤ To spraying white grubs harbouring host plant with chlorpyrifos 20 EC 0.04% (20 ml/10 l water). ➤ To Collection and destruction of White grub adults from host plant around the field. ➤ Two sprays of pongamia oil (30 ml/10 l of water) OR ponneem (30 ml/10 l of water) at 15 days interval starting from pest infestation for effective and economical management of defoliators (Helicoverpa and Spodoptera). ➤ For the control of rust, spray 5 ml Tridemorph in 10 litres of water.
Pearl millet (Kharif)	Vegetative stage	Interculturing/ Weeding	➤ Apply irrigation to the standing crop as per the crop requirement. ➤ Apply supplementary dose of 87 kg/ha urea to pearl millet at tillering stage under clear sky.
		Shootfly	➤ Two sprays of Endosulfan (0.07%) at 10 days and 20 days after germination for shootfly control.

Sesame	Vegetative stage	Leaf blight/ Stem rot/ Leaf webber	➤ If there has been no rainfall in the region, apply irrigation to the standing crop. ➤ If infestation of leaf Webber observe above ETL, spray Cartep Hydrochloride 15 gram in 10 litres of water OR Neem Seed Kernel Extract (NSKE) 3% (300g /10 l water). ➤ To control leaf blight and stem rot, spray Copper oxychloride 50 % WP 40 g/10 lit or Mancozeb 75 % WP 25 ml/10 lit. ➤ Apply basal dose of Nitrogen @ 25 kg/ha and phosphorus @ 40 kg/ha.
Maize	Vegetative stage	Interculturing/ Weeding	➤ Apply need based light irrigation by considering weather and soil moisture condition in standing crops.
Vegetable crops	Flowering /Fruits/ Vegetative stage	Interculturing/ Weeding	➤ If there has been no rainfall in the region, apply irrigation to the standing crop. ➤ Ensure weed-free cultivation of vegetable crops, and regularly send graded produce to the market. ➤ It is also advised to use shed nets, which helps in reducing vector borne diseases and produce healthy seedlings. ➤ To prevent soil borne disease drenching of nursery bed should be done with Mencozeb 75 % WP @ 2.5 gm/ litre of water after 15 DAS. ➤ Spray Quinalphos 20 ml in 10 liters of water for control of leaf miner.
		Thrips/ White fly	➤ For control of thrips, spray Profenofos 20 ml with 500 ml of 5 % neem seed extract or neem seed oil 50 ml in 10 litres of water. In case of severe attack, spray Spinosad or Spinetoram 3 ml per 10 liter of water. ➤ To minimize population of whitefly, Spray 500 ml of 5 % neem seed extract or neem oil 50 ml with Difenthiuron 50 WP @ 16 gm or Trizophos 40 EC 25 ml / 10 lit of water.
Cattle	➤ Feed animals with a mixture of green grass + hay + minerals + dry feed. ➤ Ensure adequate clean drinking water availability at all times and increase watering frequency during dry spell. ➤ Avoid overcrowding in sheds; ensure proper ventilation. ➤ Udder of milking animals must be properly clean with zinc oxide or boric powder. ➤ Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Deworming, T.B. and J.D. ➤ Keep fodder in safe place to avoid spoilage due to rainfall. ➤ Vaccination: During the monsoon season, it is recommended to vaccinate cattle against Haemorrhagic Septicaemia.		

Saurashtra-Kutch Met Subdivision:

AMFU Bhachau:

Bhachau

General advice	Likely impacts of weather warnings on Agriculture and associated Agromet advisories : NIL General Advisory ❖ <u>Under moisture stress condition following activity should be taken to save crop:</u> ▪ Carry out interculturing and weeding as per need in all kharif crops. ▪ To protect crop against sucking pests like jassid, thrips and aphid spray imidachlopride 30.5 % SC 4 ml/10 litre water. ▪ Do not apply any kind of fertilizer under moisture stress condition. ▪ Spray Kaoline 4 % solution i.e 400 gm kaolin in 10 litre of water. ▪ Straw, groundnut shell and agricultural waste mulching in wide space crops is advisable under moisture stress condition. ▪ Due to dry spell apply life saving irrigation through drip/sprinkler under moisture stress condition. ▪ Carry out earthing up in cotton. ▪ Under present weather condition caterpillar infestation observe in groundnut, soybean and pulses, so use light trap to control the infestation. Sms:Due to dry spell apply life saving irrigation through drip/sprinkler under moisture stress condition.		
Crop	Stage	Pest/diseases	Advisory
Kharif Ground nut	White grub, Early leaf Spot, Aphid, Jassid, Thrips, leaf miner, Armyworm, Red Hairy Caterpillar	Flowering/ Pegging	• Due to deficiency of Fe in soil chlorosis observed in ground nut crop. To control spray Ferrus Sulphate @ 100 gm with 10 gm citric acid in 10 lit. Water, for 1 ha makes 500 lit. Solution • Apply trichoderma spp. @ 2.5 kg with Castor cake @ 500 kg/ha. If there is more infestation apply carbendazim 0.05% @ 15 g/pump as a drenching during dry spell. Avoid inter culturing operation in crop during pegging stage. • To spraying white grubs harbouring host trees with Chlorpyrifos 20 EC 0.04% (20 ml/10 l water). • To Collection and destruction of White grub adults from host trees around the field. • For the control of sucking pest (aphid, thrips, jassid), spray 3 ml Phosphamidon or 15 ml Dimethoate or 10 ml Methyl-O- Dameton or 4 ml Imidacloprid or 5 ml Fipronil or 2 g Acetamiprid or 4 g Thiomethoxam in 10 liters of water. If ladybird beetle is observed with aphid then postpone the spraying operation. • Spray of Difenconazole 25% @ 10g/10lit. of water or Propiconazole 25% EC @ 6-7

			g/10lit.ofwaterorHexaconazole5%EC@ 30 ml/10 lit. or tebuconazole 250 EC @ 10 ml/10lit.ofwaterafterappearanceofearly leaf spot disease. • For the control of rust, Spray Tridemorf 80% @ 5ml or Mancozeb 75% 25g or Chlorothalonil 75% @ 25 to 30g dissolved in 10 liters. • Release egg parasite of <i>Telenomus remus</i> @ 50000/ ha, against <i>S. litura</i>. / Release of <i>Trichogramma chilonis</i> (50000/ha) against leaf miner and other defoliators.
Black gram	Pod development & maturity		• Carry out inter culturing and weeding under condition. Pluck mature pods in time.
soybean	Aphid, Jassid,	Vegetative growth	➤ For the control of caterpillar, spray 25 ml chlorpyrifos 20% EC or 10 ml methomyl 40% SP or 16 ml trizophos 40% EC in 10 litre of water. or use of integrated pest management like a pheromone trap, <i>Verticillium lecanii</i> fungi etc. Spray quinalphos 25 EC, oxydemeton methyl 25 EC, or dimethoate 30 EC @ 2ml/lit for control of Aphid, Jassid and Thrips
Pearl millet	Tillering	-	➤ Carry out weeding and inter culturing operations at wapsa condition in sown crop. ➤ Apply split dose of nitrogen as a top dressing after weeding under optimum soil moisture level at one month after sowing.
Cotton	Squaring & flowering	Jassids, aphid & White fly Pink Bollworm	• Under present weather condition thrips and jassids infestation observed in cotton, at above ETL spray acephate 75% SP 15gm + Fipronil 5 SC 15 ml with 1500 ppm neem oil 40 ml / 10 litre of water under clear weather. • Under present weather condition pink bollworm infestation observed in cotton. For the monitoring of pink bollworm infestation in cotton crop farmers are advised to put pheromone traps 5-6/ha. If infestation found, pick and bury the infested flowers/bolls in soil for control of pink bollworm infestation in cotton. Also make spray of profenofos 50 EC 20ml in 10 litres of water. • Carry out earthing up, interculturing under weeding operation to remove weed and conserve soil moisture. • Farmers are advised to spray 3 % KNO₃ solution at flowering, boll initiation and 50% boll formation to Bt. Cotton crop. • Foliar spraying of multi-micronutrient formulation Grade IV (Fe-Mn-Zn-CuB, 4.0-1.0-6.0-0.5-0.5 %) @ 1% at 60, 75 and 90 DAS in Bt. cotton crop is advised.


Green gram	Pod development & maturity	-	Pluck mature pods in time.
Castor	Growth	-	- Carry out weeding in time. - For the control of semilooper and prodenia in castor spray emamectin benzoate 5 gm/10 litre water.
Sesame	Flowering/capsule	Leaf Webber  Mites Phytophthora blight	- If infestation of leaf Webber observe above ETL, spray Cartep Hydrochloride 15 gram in 10 litres of water OR Neem Seed Kernel Extract (NSKE) 3% (300g /10 l water). - Carry out interculturing and weeding under condition. - Spray Mencozeb 75 % WP for prevention of infestation of Alternaria/fungal leaf spot disease in sesame. - Mite infestation observed in sesame, so monitor the crop. If population is found above ETL, then spraying of Propargite 57% @ 1.5-2ml/litre of water is advised. - Spray Ridomil 20 gm /10 litre of water for prevention of infestation of Alternaria/fungal leaf spot and Powdery mildew diseases in sesame.
Chili	Flowering & fruiting	Thrips  Mites Viral leaf curling	- For control of thrips, spray Profenofos 20 ml with 500 ml of 5 % neem seed extract or neem seed oil in 10 litres of water. In case of severe attack, spray Spinosad or Spinetoram 3 ml per 10 liter of water. - If infestation of mite observed in chilli, then spray Ethion 50 EC 15 ml or Propargite 57 % EC @ 20 ml/ 10 litre of water under clear weather. - Under weather conditions, rouging of viral infested plants should be done in chilli. Infected plants should be uprooted and buried in the soil. After rouging, spraying of Imidacloprid @ 3 ml per 10 litre of water is advised to control the vector.
Lemon	Fruit development	Citrus canker Leaf miner	- Spray copper oxychloride 40 gm or streptocyclie 1.5 gm in 10 litres of water for the control of citrus canker in lemon under clear weather. - Spray profenofos 20 ml in 10 litres of water for control of leaf miner.
Cattle	-	-	➤ Spray phenyl in the animal shed to avoid flies and mosquitoes. Give stored fodder with mineral mixture. Keep animal in shed during rain.

AMFU Targhadia:

**RAJKOT, MORBI, SURENDRANAGAR,DEVBHUMI DWARKA
AMRELI, JAMNAGAR DISTRICTS.**

Weather Warning :NIL			
Likely impacts of weather warnings on Agriculture and associated Agromet advisories : NIL			
General Advisory for ❖ <u>Under moisture stress condition following activity should be taken to save crop:</u> - Carry out interculturing and weeding as per need in all kharif crops. - To protect crop against sucking pests like jassid, thrips and aphid spray imidachlopride 30.5 % SC 4 ml/10 litre water. - Do not apply any kind of fertilizer under moisture stress condition. - Spray Kaoline 4 % solution i.e 400 gm kaolin in 10 litre of water. - Straw, groundnut shell and agricultural waste mulching in wide space crops is advisable under moisture stress condition. - Due to dry spell apply lifesaving irrigation through drip/sprinkler under moisture stress condition. - Carry out earthing up in cotton. - Under present weather condition caterpillar infestation observe in groundnut, soybean and pulses, so use light trap to control the infestation.			
❖ Weather outlook for subsequent week: As per extended range forecast Humid and partly cloudy to generally cloudy and likely light to moderate rainfall weather during 13 to 18 September-2026.			
• Sms: Due to dry spell apply life saving irrigation through drip/sprinkler under moisture stress condition.			
Crops	Stage	Insect/Pest/ Variety	Advisory
Kharif groundnut	Flowering , pegging	Aphid Weed Stem rot/ White fungus Leaf spot  White grub 	- Monitor groundnut crop for infestation of aphids. If infestation found above ETL, then spray Imidacloprid 17.8 % EC 5 ml in 10 litre of water. - Carry out interculturing and weeding under vapsa condition for better growth. - Apply 2.5 kg/ha Trichoderma fungus with well decompose FYM OR castor cake @ 500 kg/ha as preventive measures against Sclerotium rolfsii stem rot. - Spray Hexaconazole or Propiconazole 10 ml/10 lit of to prevent leaf spot & rust infestation in groundnut under clear weather. - Do drenching of Chlorpyrifos 20 % EC or

		Heliothis & Prodenia	Quinalphos 25 % EC 4 liters/ha or mix Chlorpyrifos 20 % EC or Quinalphos 25 % EC 4 litres with 100 kg sand and broadcast in 1 hectare area and irrigate if there is no rainfall. • DO spray of Beauveria bassiana@ 40 gm/10 litre of water for the control of Heliothis & Prodenia infestation in groundnut crop under clear weather.
Cotton	Squaring & flowering	Jassids, aphid & White fly Pink Bollworm	• Under present weather condition thrips and jassids infestation observed in cotton, at above ETL spray acephate 75 % SP 15 gm + Fipronil 5 SC 15 ml with 1500 ppm neem oil 40 ml /10 litre of water under clear weather. • Under present weather condition pink bollworm infestation observed in cotton. For the monitoring of pink bollworm infestation in cotton crop farmers are advised to put pheromone traps 5-6/ha. If infestation found, pick and bury the infested flowers/bolls in soil for control of pink bollworm infestation in cotton. Also make spray of profenofos 50 EC 20 ml in 10 liters of water. • Carry out earthing up, interculturing underweeding operation to remove weed and conserve soil moisture. • Farmers are advised to spray 3 % KNO₃ solution at flowering, boll initiation and 50% boll formation to Bt. Cotton crop. • Foliar spraying of multi-micronutrient formulation Grade IV (Fe-Mn-Zn-CuB, 4.0-1.0-6.0-0.5-0.5 %) @ 1% at 60, 75 and 90 DAS in Bt. cotton crop is advised.
Sesame	Flowering/ capsule	Leaf Webber  Mites Phytophthora blight	• If infestation of leaf Webber observe above ETL, spray Cartep Hydrochloride 15 gram in 10 litres of water OR Neem Seed Kernel Extract (NSKE) 3% (300g /10 l water). • Carry out interculturing and weeding under condition. • Spray Mencozeb 75 % WP for prevention of infestation of Alternaria/fungal leaf spot disease in sesame. • Mite infestation observed in seame, so monitor the crop. If population is found aboe ETL, then spraying of Propargite 57 % @ 1.5-2 ml/litre of water is advised. • Spray Ridomil 20 gm /10 litre of water for prevention of infestation of Alternaria/fungal leaf spot and Powdery mildew diseases in sesame.

Pearl millet	Heading / Flowering	-	Carry out weeding in time.
Green gram	Pod development & maturity	-	Pluck mature pods in time.
Black gram	Pod development & maturity	-	Carry out inter culturing and weeding under condition.
Cow pea	Pod development & maturity	-	Pluck mature pods in time.
Pigeon pea	Growth	-	Carry out inter culturing and weeding under condition.
Castor	Growth	-	- Carry out weeding in time. - For the control of semilooper and prodenia in castor spray emamectin benzoate 5 gm/10 litre water.
Fodder Sorghum	Growing	Shoot fly 	Continuously monitor the sorghum crop for shoot fly infestation, if infestation observe above ETL spray imidacloprid 17.8 SL 5 ml/10 litre of water.
Fodder Maize	Growing	Fall army worm	Monitor the Maize crop for fall army worm infestation, if infestation observe above ETL spray Emamectin benzoate 5 % SG 5 gm/10 litre of water.
Ladyfinger	Flowering/ fruiting	White flies Mites	- Under clear weather spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with with acetamiprid 20 % SP@ 5.0 gm/10 lit of water for white flies control in okra. - Constant monitoring of ladyfinger crop against attack of mite is advised. If population is found more, then spraying of Propargite @ 2 ml/litre of water is advised. - After harvesting of mature okra, application of urea @ 20 kg/ acre should be done. - Apply light irrigation
Ridge gourd	Flowering/ fruiting	White fly Mite Downy mildew	- In all broad leafy vegetables whitefly infestation observed above ETL. Spray 500 ml of 5 % neem seed extract or neem seed oil 50 ml with acetamiprid 20 % SP@ 5.0 gm or imidacloprid 30.5 SC 4 ml/10 lit of water for whiteflies control. - If mite population is found more, then spraying of Ethion @ 1.5-2 ml/litre of water for the control of Mites in cucurbitaceous . - For the control of downy mildew spray mencozeb 40g/pump. - Apply light irrigation.
Bottle gourd			
Sponge guard			
Bitter guard			
Cluster bean			
Cucumber			
Brinjal	Flowering/ fruiting	White fly	To minimize population of whitefly, Spray 500 ml of 5 % neem seed

		Shoot & Fruit borer	extract or neem oil 50 ml with acetamiprid 20% SP@ 5.0 gm or Difenthiuron 50 WP @ 16 gm or Trizophos 40 EC 25 ml / 10 lit of water. - In brinjal, monitoring of borers by using pheromone traps 4-6 per acre is advised. For control of shoot and fruit borer in brinjal, infested fruits and shoots should be collected and buried into the soil. If fruits and shoot borer population is above ETL then spraying of chlorantraniliprole 18.5 % SC 3.0 ml / 10 litre of water is advised. - Apply light irrigation.
Chili	Flowering & fruiting	Thrips  Mites Viral leaf curling	- For control of thrips, spray Profenofos 20 ml with 500 ml of 5 % neem seed extract or neem seed oil 50 ml in 10 litres of water. In case of severe attack, spray Spinosad or Spinetoram 3 ml per 10 liter of water. - If infestation of mite observed in chilli, then spray Ethion 50 EC 15 ml or Propargite 57 % EC @ 20 ml/ 10 litre of water under clear weather. - Under weather conditions, rouging of viral infested plants should be done in chilli. Infected plants should be uprooted and buried in the soil. After rouging, spraying of Imidacloprid @ 3 ml per 10 litre of water is advised to control the vector.
Lemon	Fruit development	Citrus canker Leaf minor	- Spray copper oxychloride 40 gm or streptocyclie 1.5 gm in 10 liters of water for the control of citrys canker in lemon under clear weather. - Spray profenofos 20 ml in 10 liters of water for control of leaf minor.
Livestock & poultry 	Keep animal in shed during rainfall hours. Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick. Deworming, T.B. and J.D. Testing should be done. Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. For milch animals, regularly follow schedule of 1 kg feed + 50 g mineral mixture per 2 liters of milk yield. Feed animals with a mixture of green grass + hay + minerals + dry feed. Ensure regarding vaccination of animals for different diseases in rainy season. Spray phenyl in the animal shed to avoid fly's and mosquitoes. Udder of milking animals must be properly clean with zinc oxide or boric powder. Also give deworming tablet to younger animals. For control of ticks and mites spray Deltamethrin or Amitral 2 ml/liter of water. Keep the fodder in safe place to prevent spoilage due to rain.		
Note	Kindly download - MAUSAM App for location specific forecast & warning. - MEGHDOOT App for Agromet advisory. - DAMINI App for lighting warning.		

AMFU Junagadh:

GIR SOMNATH

Crop	Variety/ Pest/Insect	Stage	Agromet Advisories
Kharif Groundnut	White grub, Early leaf Spot, Aphid, Jassid, Thrips, leaf miner, Armyworm, Red Hairy Caterpillar	Flowering / Pegging	➤ If irrigation facility is available, then apply irrigation to crop at critical growth stages. ➤ Interculturing and hand weeding should be carried out. ➤ To spraying white grubs harbouring host trees with chlorpyrifos 20 EC 0.04% (20 ml/10 l water). ➤ To Collection and destruction of White grub adults from host trees around the field. ➤ For the control of sucking pest (aphid, thrips jassid), spray 3 ml Phosphamidon OR 15 ml Dimethoate OR 10 ml Methyl-O-Dameton OR 4 ml Imidacloprid OR 5 ml Fipronil OR 2 g Acetamiprid OR 4 g Thiomethoxam in 10 liters of water. If ladybird beetle is observed with aphid then postpone the spraying operation. ➤ Release egg parasite of Telenomus remus @ 50000/ ha, against S. litura. / Release of Trichogramma chilonis (50000/ha) against leaf miner and other defoliators. ➤ In Ground nut, spray 0.1% FeSO₄ @ 100 gram with 10 gram citric acid solution to correct yellowing.
Cotton	Stem borer, Aphid, Leafhopper and Mealy bug	Vegetative / Branching	➤ If irrigation facility is available, then apply irrigation to crop at critical growth stages. ➤ Interculturing and hand weeding should be carried out. ➤ Post Emergence application of Pyriproxyfen sodium 6% + Quizalofop ethyl 4% MEC herbicide @ 2-2.5 ml/ litre OR Quizalofop ethyl 5 EC (0.01%) @ 50g a. i./ha (2 ml/lit of water) for control of weed. ➤ If Stem borer infestation seen then give soil application of Carbofuran 3G @ 30 kg/ha and earthing up to support crop, drain excess water and management of stem borer. ➤ Apply top dressing of nitrogenous fertilizer in cotton to support healthy vegetative growth

			stage.
Soybean	Aphid, Jassid, Thrips, leaf miner, Spodoptera litura, Helicoverpa armigera	Vegetative	➤ If irrigation facility is available, then apply irrigation to crop at critical growth stages. ➤ Interculturing and hand weeding should be carried out. ➤ For the control of caterpillar, spray 25 ml chloropyariphos 20% EC OR 10 ml methomile 40% SP OR 16 ml trizophos 40% EC in 10 litre of water. OR use of integrated pest management like a pheromone trap, Verticillium lecanii fungi etc. ➤ To spray quinalphos 25 EC, oxydemeton methyl 25 EC, or dimethoate 30 EC @ 2ml /lit for control of Aphid, Jassid and Thrips.
Pigeon pea	Gujarat Pigeon pea – 101 & 103, BDN – 2 & 711	Germination / Vegetative	➤ If irrigation facility is available, then apply irrigation to crop at critical growth stages. ➤ Arrange for extensive drainage facilities to remove excess rain water from standing crop fields. ➤ Interculturing and hand weeding should be carried out in during vapsa condition.
Vegetable crops	-	Sowing/ Flowering/ Fruiting	➤ Install 50% to 75% green shade nets to protect nurseries, young plants, and vegetable crops from direct, intense sunlight. ➤ To prevent soil borne disease drenching of nursery bed should be done with Mencozeb 75 % WP @ 2.5 gm/ litre of water after 15 DAS. ➤ For white fly and leaf minor control in vegetables spray neem seed extract 5% or 40 ml neem oil with thiamethoxam 25 % WG 5 gm or imidacloprid 30.5 % SC 5 ml in 10 liters of water. ➤ Postpone harvesting of vegetables during forecast period.
Livestock	-	-	➤ Spray phenyl in the animal shed to avoid flies and mosquitoes. ➤ Feed 20 to 50 gram mineral mixture to each milch animal according their body weight to maintain health and to avoid the problems

			like milk fever and proleptoses. ➤ Use chaff cutter for fodder. ➤ For the control of tick and other parasites, spray deltemethrin or aemitz 2 ml in 1 litre of water. ➤ Udder of milking animals must be properly clean with zinc oxide or boric powder.
--	--	--	---

JUNAGADH

Crop	Variety/ Pest/Insect	Stage	Agromet Advisories
Kharif Groundnut	White grub, Early leaf Spot, Aphid, Jassid, Thrips, leaf miner, Armyworm, Red Hairy Caterpillar	Flowering / Pegging	➤ If irrigation facility is available, then apply irrigation to crop at critical growth stages. ➤ Interculturing and hand weeding should be carried out. ➤ To spraying white grubs harbouring host trees with chlorpyrifos 20 EC 0.04% (20 ml/10 l water). ➤ To Collection and destruction of White grub adults from host trees around the field. ➤ For the control of sucking pest (aphid, thrips jassid), spray 3 ml Phosphamidon OR 15 ml Dimethoate OR 10 ml Methyl-O-Dameton OR 4 ml Imidacloprid OR 5 ml Fipronil OR 2 g Acetamiprid OR 4 g Thiomethoxam in 10 liters of water. If ladybird beetle is observed with aphid then postpone the spraying operation. ➤ Release egg parasite of Telenomus remus @ 50000/ ha, against S. litura. / Release of Trichogramma chilonis (50000/ha) against leaf miner and other defoliators. ➤ In Ground nut, spray 0.1% FeSO₄ @ 100 gram with 10 gram citric acid solution to correct yellowing.
Cotton	Stem borer, Aphid, Leafhopper and Mealy bug	Vegetative / Branching	➤ If irrigation facility is available, then apply irrigation to crop at critical growth stages. ➤ Interculturing and hand weeding should be carried out. ➤ Post Emergence application of Pyrethrin 6% + Quinalofop ethyl 4% MEC herbicide @ 2-2.5 ml/ litre OR Quinalofop ethyl 5 EC (0.01%) @ 50g a. i./ha (2 ml/lit of water) for control of weed.

			➤ If Stem borer infestation seen then give soil application of Carbofuran 3G @ 30 kg/ha and earthing up to support crop, drain excess water and management of stem borer. ➤ Apply top dressing of nitrogenous fertilizer in cotton to support healthy vegetative growth stage.
Soybean	Aphid, Jassid, Thrips, leaf miner, Spodoptera litura, Helicoverpa armigera	Vegetative	➤ If irrigation facility is available, then apply irrigation to crop at critical growth stages. ➤ Interculturing and hand weeding should be carried out. ➤ For the control of caterpillar, spray 25 ml chlorpyrifos 20% EC OR 10 ml methomyl 40% SP OR 16 ml trizophos 40% EC in 10 litre of water. OR use of integrated pest management like a pheromone trap, Verticillium lecanii fungi etc. ➤ To spray quinalphos 25 EC, oxydemeton methyl 25 EC, or dimethoate 30 EC @ 2ml /lit for control of Aphid, Jassid and Thrips.
Pigeon pea	Gujarat Pigeon pea – 101 & 103, BDN – 2 & 711	Germination / Vegetative	➤ If irrigation facility is available, then apply irrigation to crop at critical growth stages. ➤ Arrange for extensive drainage facilities to remove excess rain water from standing crop fields. ➤ Interculturing and hand weeding should be carried out in during vapsa condition.
Vegetable crops	-	Sowing/ Flowering/ Fruiting	➤ Install 50% to 75% green shade nets to protect nurseries, young plants, and vegetable crops from direct, intense sunlight. ➤ To prevent soil borne disease drenching of nursery bed should be done with Mencozeb 75 % WP @ 2.5 gm/ litre of water after 15 DAS. ➤ For white fly and leaf minor control in vegetables spray neem seed extract 5% or 40 ml neem oil with thiamethoxam 25 % WG 5 gm or imidacloprid 30.5 % SC 5 ml in 10 liters of water. ➤ Postpone harvesting of vegetables during forecast period.
Livestock	-	-	➤ Spray phenyl in the animal shed to avoid flies

			and mosquitoes. ➤ Feed 20 to 50 gram mineral mixture to each milch animal according their body weight to maintain health and to avoid the problems like milk fever and prolepses. ➤ Use chaff cutter for fodder. ➤ For the control of tick and other parasites, spray deltemethrin or aemitaz 2 ml in 1 litre of water. ➤ Udder of milking animals must be properly clean with zinc oxide or boric powder.
--	--	--	--

PORBANDAR

Crop	Variety/ Pest/Insect	Stage	Agromet Advisories
Kharif Groundnut	White grub, Early leaf Spot, Aphid, Jassid, Thrips, leaf miner, Armyworm, Red Hairy Caterpillar	Flowering / Pegging	➤ If irrigation facility is available, then apply irrigation to crop at critical growth stages. ➤ Interculturing and hand weeding should be carried out. ➤ To spraying white grubs harbouring host trees with chlorpyrifos 20 EC 0.04% (20 ml/10 l water). ➤ To Collection and destruction of White grub adults from host trees around the field. ➤ For the control of sucking pest (aphid, thrips jassid), spray 3 ml Phosphamidon OR 15 ml Dimethoate OR 10 ml Methyl-O-Dameton OR 4 ml Imidacloprid OR 5 ml Fipronil OR 2 g Acetamiprid OR 4 g Thiomethoxam in 10 liters of water. If ladybird beetle is observed with aphid then postpone the spraying operation. ➤ Release egg parasite of Telenomus remus @ 50000/ ha, against S. litura. / Release of Trichogramma chilonis (50000/ha) against leaf miner and other defoliators. ➤ In Ground nut, spray 0.1% FeSO₄ @ 100 gram with 10 gram citric acid solution to correct yellowing.
Cotton	Stem borer, Aphid, Leafhopper and Mealy bug	Vegetative / Branching	➤ If irrigation facility is available, then apply irrigation to crop at critical growth stages. ➤ Interculturing and hand weeding should be carried out. ➤ Post Emergence application of Pyrethriobac

			sodium 6% + Quizalofop ethyl 4% MEC herbicide @ 2-2.5 ml/ litre OR Quizalofop ethyl 5 EC (0.01%) @ 50g a. i./ha (2 ml/lit of water) for control of weed. ➤ If Stem borer infestation seen then give soil application of Carbofuran 3G @ 30 kg/ha and earthing up to support crop, drain excess water and management of stem borer. ➤ Apply top dressing of nitrogenous fertilizer in cotton to support healthy vegetative growth stage.
Soybean	Aphid, Jassid, Thrips, leaf miner, Spodoptera litura, Helicoverpa armigera	Vegetative	➤ If irrigation facility is available, then apply irrigation to crop at critical growth stages. ➤ Interculturing and hand weeding should be carried out. ➤ For the control of caterpillar, spray 25 ml chlorpyrifos 20% EC OR 10 ml methomyl 40% SP OR 16 ml trizophos 40% EC in 10 litre of water. OR use of integrated pest management like a pheromone trap, Verticillium lecanii fungi etc. ➤ To spray quinalphos 25 EC, oxydemeton methyl 25 EC, or dimethoate 30 EC @ 2ml /lit for control of Aphid, Jassid and Thrips.
Pigeon pea	Gujarat Pigeon pea – 101 & 103, BDN – 2 & 711	Germination / Vegetative	➤ If irrigation facility is available, then apply irrigation to crop at critical growth stages. ➤ Arrange for extensive drainage facilities to remove excess rain water from standing crop fields. ➤ Interculturing and hand weeding should be carried out in during vapsa condition.
Vegetable crops	-	Sowing/ Flowering/ Fruiting	➤ Install 50% to 75% green shade nets to protect nurseries, young plants, and vegetable crops from direct, intense sunlight. ➤ To prevent soil borne disease drenching of nursery bed should be done with Mencozeb 75 % WP @ 2.5 gm/ litre of water after 15 DAS. ➤ For white fly and leaf minor control in vegetables spray neem seed extract 5% or 40 ml neem oil with thiamethoxam 25 % WG 5 gm or imidacloprid 30.5 % SC 5 ml in 10 liters of water.

			➤ Postpone harvesting of vegetables during forecast period.
Livestock	-	-	➤ Spray phenyl in the animal shed to avoid flies and mosquitoes. ➤ Feed 20 to 50 gram mineral mixture to each milch animal according their body weight to maintain health and to avoid the problems like milk fever and prolepses. ➤ Use chaff cutter for fodder. ➤ For the control of tick and other parasites, spray deltemethrin or aemitaz 2 ml in 1 litre of water. ➤ Udder of milking animals must be properly clean with zinc oxide or boric powder.

AMFU Arnej:

AHMEDABAD

Advisory		
General advice	In prevailing dry spell condition, undertake the interculturing and weeding operations to conserve moisture in standing monsoon crops and apply need based light irrigation by considering weather and soil moisture condition in standing crops. Avoid application of fertilizer or top dressing or it should undertake only under optimum soil moisture condition in standing crops. Farmers are advised to download MEGHDOOT mobile application for acquiring information regarding Agromet Advisory in Gujaratilanguage. Click on following link to download this applicationuser:https://play.google.com/store/apps/details?id=com.aas.meghdoot	
Field Crops	Stage	Advisory
Rice	Booting/	• Maintain 5-7 cm stagnant water in rice field.

	Heading/ Flowering	• Spray mixture of streptocycline 1 g + copper oxychloride 20 g fungicides in 400-500 litres of water per hectare to control bacterial leaf blight disease. • Spray tricyclazole 75 WP 6 g or Iprobenfos 48 EC 10 ml in 10 litres of water at an interval of 15 to 20 days to control blast disease. • For proper nutrient management in transplanted rice, first top dressing of nitrogen fertilizer should be applied 30–40 days after transplanting. Apply 32 kg of nitrogen per hectare for early-maturing varieties, 40 kg of nitrogen per hectare for medium to late-maturing varieties and 48 kg of nitrogen per hectare for late-maturing varieties. • The rice yellow stem borer in rice can be effectively managed by installing five pheromone traps per hectare baited with lures that attract the male moths. Spray neem-based insecticides by mixing 10 ml (5 EC) to 60 ml (0.03 EC) in 10 litres of water. Alternatively, apply Cartap Hydrochloride 4G @ 8 kg per acre or Fipronil 0.3G with 10 kg per acre or Chlorantraniliprole 0.4G with 4 kg per acre after reducing the water level in the field. These granular insecticides should be applied twice, first application at the onset of pest infestation or 30–35 days after transplanting and second application at 15–20 days later. This schedule helps keep the pest infestation under control.
Cotton	Squaring/ Flowering/ Boll formation	• If irrigation facilities are available then apply need based irrigation through drip irrigation/alternate furrow irrigation method by considering soil moisture condition in standing crop. • Avoid application of fertilizers during long dry spell condition or it should apply under optimum soil moisture condition. Spray 2 % urea during dry spell in standing cotton crop. • Sucking pest infestation in cotton at initial stage can be controlled by spraying 500 g of neem seed kernel extract (5% NSKE) or 50 ml of neem oil or a neem-based insecticide by mixing 10 ml (5 EC) to 50 ml (0.03 EC) in 10 litres of water. If the infestation is severe, spray Flonicamid 50 WG with 3 g or Diafenthiuron 50 WP with 10 g or Dinotefuran 20 SG with 3 g by mixing the

		recommended quantity in 10 litres of water. To prevent an increase in the population of whiteflies and jassids (leafhoppers), do not use synthetic pyrethroid insecticides until the month of November.
Groundnut	Flowering/ Pod formation	- Leaf-eating caterpillar (Spodoptera) in groundnut can be controlled by installing one light trap per hectare in the affected area and destroy the adult moths and leaf-eating caterpillar moths attracted to the trap. - White grubs in groundnut can be controlled by spraying Chlorpyrifos 20EC with 4litres per hectare through drip irrigation or along with irrigation. If irrigation is not being given but intermittent rainfall is occurring, prepare the insecticide solution in a spray pump, remove the nozzle and apply the solution directly into the furrows in the required quantity. - Leaf-eating caterpillar (Spodoptera) in groundnut can be controlled by installing pheromone traps with 40 traps per hectare on a community basis to keep the pest population under control. At the initial stage of infestation, spray either neem seed kernel powder extract at 500 g (5% extract) or a neem-based insecticide at 20 ml (1 EC) to 40 ml (0.15 EC) mixed in 10 litres of water. If the infestation becomes severe, spray Chlorantraniliprole 18.5 SC at 3 ml or Flubendiamide 20 WG at 6 g or Thiamethoxam 12.6% + Lambda-cyhalothrin 9.5% ZC at 3 ml 10 litres of water. - Cutworms in groundnut can be controlled by installing one light trap per hectare to attract and destroy the moths. Prepare a 5% neem extract by mixing 500 g of neem seed kernel powder or neem leaves in 10 litres of water and spray it on the crop. If the infestation is severe, spray either Indoxacarb 14.5 SC with 10 ml per 10 litres of water or Emamectin benzoate 5 SG with 5 g per 10 litres of water.
Green gram	Flowering/ Pod formation	- Apply need based light irrigation by considering weather and soil moisture condition in green gram crop. - Mosaic of leaf curl disease is transmitted due to white fly so to control it, spray imidacloprid 17.8 SC 3 ml or acetamiprid 20 SP 2 g or azadirachtin 40 ml in 10 litres of water in green gram crop. - Hairy caterpillar in green gram can be controlled by installing one light trap per hectare to attract and

		destroy the adult moths of.Spray 500 g of neem seed kernel powder or neem leaves (5% extract) mixed in 10 litres of water to control hairy caterpillar in clear and calm weather condition.If the infestation is severe, spray Chlorantraniliprole 18.5SC with 4ml or Indoxacarb 15.8EC with 6ml or Emamectin Benzoate 5SG with 5g mixed in 10litres of water. • The infestation by aphids, whiteflies, leafhoppers and thrips in green gram at initial stage can be controlled by spraying 500 g of neem seed kernel powder (5% extract) or 40 g of the fungal biocontrol agent Verticillium lecanii powder mixed in 10 litres of water. • If there is a severe infestation of aphids in green gram and predatory insects are absent, spray Imidacloprid 17.8 SL with 2.5 ml or Thiamethoxam 25 WG with 4 g per 10 litres of water.
Black gram	Flowering/ Pod formation	• Apply need based light irrigation by considering weather and soil moisture condition in black gram crop. • Mosaic of leaf curl disease is transmitted due to white fly so to control it, spray imidacloprid 17.8 SC 3 ml or acetamiprid 20 SP 2 g or azadirachtin 40 ml in 10 litres of water in green gram crop. • Hairy caterpillar in black gram can be controlled by installing one light trap per hectare to attract and destroy the adult moths.Spray 500 g of neem seed kernel powder or neem leaves (5% extract) mixed in 10 litres of water to control hairy caterpillar in clear and calm weather condition.If the infestation is severe, spray Chlorantraniliprole 18.5SC with 4ml or Indoxacarb 15.8EC with 6ml or Emamectin Benzoate 5SG with 5g, mixed in 10litres of water. • The infestation by aphids, whiteflies, leafhoppers and thrips in black gram at initial stage can be controlled by spraying 500 g of neem seed kernel powder (5% extract) or 40 g of the fungal biocontrol agent Verticillium lecanii powder mixed in 10 litres of water. • If there is a severe infestation of aphids in black gram and predatory insects are absent, spray either Imidacloprid 17.8 SL with 2.5 ml per 10 litres of water or Thiamethoxam 25 WG with 4 g per 10 litres of water.
Pigeon pea	Vegetative	• Apply need based light irrigation by considering

		weather and soil moisture condition in pigeon pea crop. • Perform hand weeding three times at 40 and 60 days after sowing.
Pearl millet	Heading/ Flowering	• Shoot fly in pearl millet can be controlled by removing and destroying plants damaged by the pest during the early crop stage. At the initial stage of infestation, spray Beauveria bassiana 1.15% WP at 40 g mixed in 10 litres of water in clear and calm weather condition. In case of severe infestation, spray either Quinalphos 25 EC at 20 ml or Cypermethrin 10 EC at 5 ml mixed in 10 litres of water. • Downy mildew in pearl millet can be controlled by spraying Metalaxyl-MZ 72 WP with 15 g per 10 litres of water, as required, ensuring thorough coverage of both the upper and lower surfaces of the leaves.
Fruit Crop	Stage	Advisory
Chilli	Vegetative	• Apply need based light irrigation by considering weather and soil moisture condition in chilli crop. • Thrips in chilli can be controlled by spraying neem seed oil 30 to 50 ml in 10 litres. In case of heavy attack, spray of spinosad 45 SC 3 ml or spinetoram 11.7 SC 10 ml in 10 litres of water.
Tomato	Vegetative	• Apply need based light irrigation by considering weather and soil moisture condition in tomato crop. • Pinworm at initial incidence in tomato can be controlled by installing 40 yellow sticky traps per hectare. In heavy incidence, spray Chlorantraniliprole 18.5 SC 3 ml or Flubendiamide 480 SC 3 ml in 10 litres of water.
Brinjal	Vegetative	• Apply need based light irrigation by considering weather and soil moisture condition in brinjal crop • Sucking pest like white fly and jassid in brinjal can be controlled by spraying Difenthiuron 50 WP 4 ml or Thiamethoxam 25 WG 4 g in 10 litres of water.
Livestock		
• Keep animals at safer places in shed during abnormal weather condition. • Spray phenyl in the animal shed to avoid flies and mosquitoes. • Don't allow your animals to congregate under trees. • Vaccinate the animals against outbreaks of diseases like Foot & Mouth disease (FMD), Haemorrhagic Septicaemia (H.S.) and Black quarter (B.Q.) during onset of monsoon.		

BHAVNAGAR

Advisory		
General advice	In prevailing dry spell condition, undertake the interculturing and weeding operations to conserve moisture in standing monsoon crops and apply need based light irrigation by considering weather and soil moisture condition in standing crops. Avoid application of fertilizer or top dressing or it should undertake only under optimum soil moisture condition in standing crops. Farmers are advised to download MEGHDOOT mobile application for acquiring information regarding Agromet Advisory in Gujaratilanguage. Click on following link to download this applicationuser:https://play.google.com/store/apps/details?id=com.aas.me ghdoot	
Field Crops	Stage	Advisory
Cotton	Squaring/ Flowering/ Boll formation	• If irrigation facilities are available then apply need based irrigation through drip irrigation/alternate furrow irrigation method by considering soil moisture condition in standing crop. • Avoid application of fertilizers during long dry spell condition or it should apply under optimum soil moisture condition. Spray 2 % urea during dry spell in standing cotton crop. • Sucking pest infestation in cotton at initial stage can be controlled by spraying 500 g of neem seed kernel extract (5% NSKE) or 50 ml of neem oil or a neem-based insecticide by mixing 10 ml (5 EC) to 50 ml (0.03 EC) in 10 litres of water. If the infestation is severe, spray Flonicamid 50 WG with 3 g or Diafenthiuron 50 WP with 10 g or Dinotefuran 20 SG with 3 g by mixing the recommended quantity in 10 litres of water. • To prevent an increase in the population of whiteflies and jassids (leafhoppers), do not use synthetic pyrethroid insecticides until the month of November.
Groundnut	Flowering/ Pod formation	• Leaf-eating caterpillar (Spodoptera) in groundnut can be controlled by installing one light trap per hectare in the affected area and destroy the adult moths and leaf-eating caterpillar moths attracted to the trap. • White grubs in groundnut can be controlled by spraying Chlorpyrifos 20EC with 4litres per hectare through drip irrigation or along with irrigation. If irrigation is not being given but intermittent rainfall is occurring, prepare the insecticide solution in a spray pump, remove the nozzle and apply the solution directly into the furrows in the required quantity.

		• Leaf-eating caterpillar (Spodoptera) in groundnut can be controlled by installing pheromone traps with 40 traps per hectare on a community basis to keep the pest population under control. At the initial stage of infestation, spray either neem seed kernel powder extract at 500 g (5% extract) or a neem-based insecticide at 20 ml (1 EC) to 40 ml (0.15 EC) mixed in 10 litres of water. If the infestation becomes severe, spray Chlorantraniliprole 18.5 SC at 3 ml or Flubendiamide 20 WG at 6 g or Thiamethoxam 12.6% + Lambda-cyhalothrin 9.5% ZC at 3 ml 10 litres of water. • Cutworms in groundnut can be controlled by installing one light trap per hectare to attract and destroy the moths. Prepare a 5% neem extract by mixing 500 g of neem seed kernel powder or neem leaves in 10 litres of water and spray it on the crop. If the infestation is severe, spray either Indoxacarb 14.5 SC with 10 ml per 10 litres of water or Emamectin benzoate 5 SG with 5 g per 10 litres of water.
Green gram	Flowering/ Pod formation	• Apply need based light irrigation by considering weather and soil moisture condition in green gram crop. • Mosaic of leaf curl disease is transmitted due to white fly so to control it, spray imidacloprid 17.8 SC 3 ml or acetamiprid 20 SP 2 g or azadirachtin 40 ml in 10 litres of water in green gram crop. • Hairy caterpillar in green gram can be controlled by installing one light trap per hectare to attract and destroy the adult moths of. Spray 500 g of neem seed kernel powder or neem leaves (5% extract) mixed in 10 litres of water to control hairy caterpillar in clear and calm weather condition. If the infestation is severe, spray Chlorantraniliprole 18.5SC with 4ml or Indoxacarb 15.8EC with 6ml or Emamectin Benzoate 5SG with 5g mixed in 10litres of water. • The infestation by aphids, whiteflies, leafhoppers and thrips in green gram at initial stage can be controlled by spraying 500 g of neem seed kernel powder (5% extract) or 40 g of the fungal biocontrol agent Verticillium lecanii powder mixed in 10 litres of water. • If there is a severe infestation of aphids in green gram and predatory insects are absent, spray Imidacloprid 17.8 SL with 2.5 ml or Thiamethoxam 25 WG with 4 g

		per 10 litres of water.
Black gram	Flowering/ Pod formation	• Apply need based light irrigation by considering weather and soil moisture condition in black gram crop. • Mosaic of leaf curl disease is transmitted due to white fly so to control it, spray imidacloprid 17.8 SC 3 ml or acetamiprid 20 SP 2 g or azadirachtin 40 ml in 10 litres of water in green gram crop. • Hairy caterpillar in black gram can be controlled by installing one light trap per hectare to attract and destroy the adult moths. Spray 500 g of neem seed kernel powder or neem leaves (5% extract) mixed in 10 litres of water to control hairy caterpillar in clear and calm weather condition. If the infestation is severe, spray Chlorantraniliprole 18.5SC with 4ml or Indoxacarb 15.8EC with 6ml or Emamectin Benzoate 5SG with 5g, mixed in 10litres of water. • The infestation by aphids, whiteflies, leafhoppers and thrips in black gram at initial stage can be controlled by spraying 500 g of neem seed kernel powder (5% extract) or 40 g of the fungal biocontrol agent Verticillium lecanii powder mixed in 10 litres of water. • If there is a severe infestation of aphids in black gram and predatory insects are absent, spray either Imidacloprid 17.8 SL with 2.5 ml per 10 litres of water or Thiamethoxam 25 WG with 4 g per 10 litres of water.
Pigeon pea	Vegetative	• Apply need based light irrigation by considering weather and soil moisture condition in pigeon pea crop. • Perform hand weeding three times at 40 and 60 days after sowing.
Pearl millet	Heading/ Flowering	• Shoot fly in pearl millet can be controlled by removing and destroying plants damaged by the pest during the early crop stage. At the initial stage of infestation, spray Beauveria bassiana 1.15% WP at 40 g mixed in 10 litres of water in clear and calm weather condition. In case of severe infestation, spray either Quinalphos 25 EC at 20 ml or Cypermethrin 10 EC at 5 ml mixed in 10 litres of water. • Downy mildew in pearl millet can be controlled by spraying Metalaxyl-MZ 72 WP with 15 g per 10 litres of water, as required, ensuring thorough coverage of both the upper and lower surfaces of the leaves.
Fruit Crop	Stage	Advisory

Chilli	Vegetative	• Apply need based light irrigation by considering weather and soil moisture condition in chilli crop. • Thrips in chilli can be controlled by spraying neem seed oil 30 to 50 ml in 10 litres. In case of heavy attack, spray of spinosad 45 SC 3 ml or spinetoram 11.7 SC 10 ml in 10 litres of water.
Tomato	Vegetative	• Apply need based light irrigation by considering weather and soil moisture condition in tomato crop. • Pinworm at initial incidence in tomato can be controlled by installing 40 yellow sticky traps per hectare. In heavy incidence, spray Chlorantraniliprole 18.5 SC 3 ml or Flubendiamide 480 SC 3 ml in 10 litres of water.
Brinjal	Vegetative	• Apply need based light irrigation by considering weather and soil moisture condition in brinjal crop • Sucking pest like white fly and jassid in brinjal can be controlled by spraying Difenthiuron 50 WP 4 ml or Thiamethoxam 25 WG 4 g in 10 litres of water.
Livestock		
• Keep animals at safer places in shed during abnormal weather condition. • Spray phenyl in the animal shed to avoid flies and mosquitoes. • Don't allow your animals to congregate under trees. • Vaccinate the animals against outbreaks of diseases like Foot & Mouth disease (FMD), Haemorrhagic Septicaemia (H.S.) and Black quarter (B.Q.) during onset of monsoon.		

BOTAD

Advisory		
General advice	In prevailing dry spell condition, undertake the interculturing and weeding operations to conserve moisture in standing monsoon crops and apply need based light irrigation by considering weather and soil moisture condition in standing crops. Avoid application of fertilizer or top dressing or it should undertake only under optimum soil moisture condition in standing crops. Farmers are advised to download MEGHDOOT mobile application for acquiring information regarding Agromet Advisory in Gujaratilanguage. Click on following link to download this applicationuser:https://play.google.com/store/apps/details?id=com.aas.meghdoot	
Field Crops	Stage	Advisory
Cotton	Squaring/ Flowering/	• If irrigation facilities are available then apply need based irrigation through drip irrigation/alternate

	Boll formation	furrow irrigation method by considering soil moisture condition in standing crop. • Avoid application of fertilizers during long dry spell condition or it should apply under optimum soil moisture condition. Spray 2 % urea during dry spell in standing cotton crop. • Sucking pest infestation in cotton at initial stage can be controlled by spraying 500 g of neem seed kernel extract (5% NSKE) or 50 ml of neem oil or a neem-based insecticide by mixing 10 ml (5 EC) to 50 ml (0.03 EC) in 10 litres of water. If the infestation is severe, spray Flonicamid 50 WG with 3 g or Diafenthiuron 50 WP with 10 g or Dinotefuran 20 SG with 3 g by mixing the recommended quantity in 10 litres of water. • To prevent an increase in the population of whiteflies and jassids (leafhoppers), do not use synthetic pyrethroid insecticides until the month of November.
Groundnut	Flowering/ Pod formation	• Leaf-eating caterpillar (Spodoptera) in groundnut can be controlled by installing one light trap per hectare in the affected area and destroy the adult moths and leaf-eating caterpillar moths attracted to the trap. • White grubs in groundnut can be controlled by spraying Chlorpyrifos 20EC with 4litres per hectare through drip irrigation or along with irrigation. If irrigation is not being given but intermittent rainfall is occurring, prepare the insecticide solution in a spray pump, remove the nozzle and apply the solution directly into the furrows in the required quantity. • Leaf-eating caterpillar (Spodoptera) in groundnut can be controlled by installing pheromone traps with 40 traps per hectare on a community basis to keep the pest population under control. At the initial stage of infestation, spray either neem seed kernel powder extract at 500 g (5% extract) or a neem-based insecticide at 20 ml (1 EC) to 40 ml (0.15 EC) mixed in 10 litres of water. If the infestation becomes severe, spray Chlorantraniliprole 18.5 SC at 3 ml or Flubendiamide 20 WG at 6 g or Thiamethoxam 12.6% + Lambda-cyhalothrin 9.5% ZC at 3 ml 10 litres of water. • Cutworms in groundnut can be controlled by installing one light trap per hectare to attract and destroy the moths. Prepare a 5% neem extract by mixing 500 g of neem seed kernel powder or neem leaves in 10 litres of water and spray it on the crop. If the infestation is

		severe, spray either Indoxacarb 14.5 SC with 10 ml per 10 litres of water or Emamectin benzoate 5 SG with 5 g per 10 litres of water.
Green gram	Flowering/ Pod formation	• Apply need based light irrigation by considering weather and soil moisture condition in green gram crop. • Mosaic of leaf curl disease is transmitted due to white fly so to control it, spray imidacloprid 17.8 SC 3 ml or acetamiprid 20 SP 2 g or azadirachtin 40 ml in 10 litres of water in green gram crop. • Hairy caterpillar in green gram can be controlled by installing one light trap per hectare to attract and destroy the adult moths of. Spray 500 g of neem seed kernel powder or neem leaves (5% extract) mixed in 10 litres of water to control hairy caterpillar in clear and calm weather condition. If the infestation is severe, spray Chlorantraniliprole 18.5SC with 4ml or Indoxacarb 15.8EC with 6ml or Emamectin Benzoate 5SG with 5g mixed in 10 litres of water. • The infestation by aphids, whiteflies, leafhoppers and thrips in green gram at initial stage can be controlled by spraying 500 g of neem seed kernel powder (5% extract) or 40 g of the fungal biocontrol agent Verticillium lecanii powder mixed in 10 litres of water. • If there is a severe infestation of aphids in green gram and predatory insects are absent, spray Imidacloprid 17.8 SL with 2.5 ml or Thiamethoxam 25 WG with 4 g per 10 litres of water.
Black gram	Flowering/ Pod formation	• Apply need based light irrigation by considering weather and soil moisture condition in black gram crop. • Mosaic of leaf curl disease is transmitted due to white fly so to control it, spray imidacloprid 17.8 SC 3 ml or acetamiprid 20 SP 2 g or azadirachtin 40 ml in 10 litres of water in green gram crop. • Hairy caterpillar in black gram can be controlled by installing one light trap per hectare to attract and destroy the adult moths. Spray 500 g of neem seed kernel powder or neem leaves (5% extract) mixed in 10 litres of water to control hairy caterpillar in clear and calm weather condition. If the infestation is severe, spray Chlorantraniliprole 18.5SC with 4ml or

		Indoxacarb 15.8EC with 6ml or Emamectin Benzoate 5SG with 5g, mixed in 10litres of water. - The infestation by aphids, whiteflies, leafhoppers and thrips in black gram at initial stage can be controlled by spraying 500 g of neem seed kernel powder (5% extract) or 40 g of the fungal biocontrol agent Verticillium lecanii powder mixed in 10 litres of water. - If there is a severe infestation of aphids in black gram and predatory insects are absent, spray either Imidacloprid 17.8 SL with 2.5 ml per 10 litres of water or Thiamethoxam 25 WG with 4 g per 10 litres of water.
Pigeon pea	Vegetative	- Apply need based light irrigation by considering weather and soil moisture condition in pigeon pea crop. - Perform hand weeding three times at 40 and 60 days after sowing.
Pearl millet	Heading/ Flowering	- Shoot fly in pearl millet can be controlled by removing and destroying plants damaged by the pest during the early crop stage. At the initial stage of infestation, spray Beauveria bassiana 1.15% WP at 40 g mixed in 10 litres of water in clear and calm weather condition. In case of severe infestation, spray either Quinalphos 25 EC at 20 ml or Cypermethrin 10 EC at 5 ml mixed in 10 litres of water. - Downy mildew in pearl millet can be controlled by spraying Metalaxyl-MZ 72 WP with 15 g per 10 litres of water, as required, ensuring thorough coverage of both the upper and lower surfaces of the leaves.
Fruit Crop	Stage	Advisory
Chilli	Vegetative	- Apply need based light irrigation by considering weather and soil moisture condition in chilli crop. - Thrips in chilli can be controlled by spraying neem seed oil 30 to 50 ml in 10 litres. In case of heavy attack, spray of spinosad 45 SC 3 ml or spinetoram 11.7 SC 10 ml in 10 litres of water.
Tomato	Vegetative	- Apply need based light irrigation by considering weather and soil moisture condition in tomato crop. - Pinworm at initial incidence in tomato can be controlled by installing 40 yellow sticky traps per hectare. In heavy incidence, spray Chlorantraniliprole 18.5 SC 3 ml or Flubendiamide 480 SC 3 ml in 10 litres of water.
Brinjal	Vegetative	Apply need based light irrigation by considering weather and soil moisture condition in brinjal crop

- Sucking pest like white fly and jassid in brinjal can be controlled by spraying Difenthiuron 50 WP 4 ml or Thiamethoxam 25 WG 4 g in 10 litres of water.

Livestock

- Keep animals at safer places in shed during abnormal weather condition.
- Spray phenyl in the animal shed to avoid flies and mosquitoes.
- Don't allow your animals to congregate under trees.
- Vaccinate the animals against outbreaks of diseases like Foot & Mouth disease (FMD), Haemorrhagic Septicaemia (H.S.) and Black quarter (B.Q.) during onset of monsoon.



Annexure I

District level weather warning issued by M.C.Ahmedabad

THUNDERSTORM WARNING:

દિવસ - 4 ગુજરાત રાજ્ય અને દીવ, દમણ, દાદરા નગર હવેલી ની તારીખ 12-09-2026 માટે
મેઘગર્જનાની ચેતવણી



દિવસ - 5 ગુજરાત રાજ્ય અને દીવ, દમણ, દાદરા નગર હવેલી ની તારીખ 13-09-2026 માટે
મેઘગર્જનાની ચેતવણી



HEAVY RAINFALL WARNING:


દિવસ - 4 ગુજરાત રાજ્ય અને દીવ, દમણ, દાદરા નગર હવેલી ની 12/09/2026 માટે
'ભારે વરસાદ ની ચેતવણી'

દિવસ - 5 ગુજરાત રાજ્ય અને દીવ, દમણ, દાદરા નગર હવેલી ની 13/09/2026 માટે
'ભારે વરસાદ ની ચેતવણી'



Meteorological Subdivision Level Rainfall Weather Forecast for next 5 days

RAINFALL DISTRIBUTION



District Forecast & Warnings for Next 7 Days

Wednesday, 09 September 2026

Time of Issue: 12:00 PM IST

DISTRICT	Intensity / Probability	09 Sep (Wed) (Day-1)	10 Sep (Thu) (Day-2)	11 Sep (Fri) (Day-3)	12 Sep (Sat) (Day-4)	13 Sep (Sun) (Day-5)	14 Sep (Mon) (Day-6)	15 Sep (Tue) (Day-7)
BANASKANTHA	Intensity	Light to Moderate rain/thundershowers at isolated places	Light to Moderate rain/thundershowers at isolated places	Light to Moderate rain/thundershowers at isolated places	Light to Moderate rain/thundershowers at a few places	Light thunderstorm with lightning and surface wind 30-40 kmph accompanied with light to moderate rain at isolated places	Light to Moderate rain/thundershowers at many places	Light to Moderate rain/thundershowers at a few places
	Probability	Very Likely	Very Likely	Very Likely	Very Likely	Very Likely	Very Likely	Very Likely
PATAN	Intensity	Light to Moderate rain/thundershowers at isolated places	Light to Moderate rain/thundershowers at isolated places	Light to Moderate rain/thundershowers at isolated places	Light to Moderate rain/thundershowers at a few places	Light thunderstorm with lightning and surface	Light to Moderate rain/thundershowers at many places	Light to Moderate rain/thundershowers at a few places

						wind 30-40 kmph accompanied with light to moderate rain at isolated places		
	Probability	Very Likely	Very Likely	Very Likely	Very Likely	Very Likely	Very Likely	Very Likely
MEHSANA	Intensity	Light to Moderate rain/thundershowers at isolated places	Light to Moderate rain/thundershowers at isolated places	Light to Moderate rain/thundershowers at isolated places	Light thunders torm with lightning and surface wind 30-40 kmph accompanied with light to moderate rain at isolated places	Light thunderstorm with lightning and surface wind 30-40 kmph accompanied with light to moderate rain at isolated places	Light to Moderate rain/thundershowers at many places	Light to Moderate rain/thundershowers at a few places
	Probability	Very Likely	Very Likely	Very Likely	Very Likely	Very Likely	Very Likely	Very Likely
SABARKANTHA	Intensity	Light to Moderate rain/thun	Light to Moderate rain/thun	Light to Moderate rain/thun	Light thunders torm with lightning	Light thunderstorm	Light to Moderate rain/thun	Light to Moderate rain/thun

	ability	Likely	Likely	Likely	Likely	Likely	Likely	Likely
ARAVALI	Intensity	Light to Moderate rain/thundershowers at isolated places	Light to Moderate rain/thundershowers at isolated places	Light to Moderate rain/thundershowers at isolated places	Light thunders torm with lightning and surface wind 30-40 kmph accompanied with light to moderate rain at isolated places	Light thunders torm with lightning and surface wind 30-40 kmph accompanied with light to moderate rain at isolated places	Light to Moderate rain/thundershowers at a few places	Light to Moderate rain/thundershowers at a few places
	Probability	Very Likely	Very Likely	Very Likely	Very Likely	Very Likely	Very Likely	Very Likely
KHEDA	Intensity	Light to Moderate rain/thundershowers at isolated places	Light to Moderate rain/thundershowers at isolated places	Light to Moderate rain/thundershowers at isolated places	Light thunders torm with lightning and surface wind 30-40 kmph accompanied with light to moderate rain at isolated places	Light thunders torm with lightning and surface wind 30-40 kmph accompanied with light to moderate rain	Light to Moderate rain/thundershowers at a few places	Light to Moderate rain/thundershowers at a few places

						at isolated places		
	Probability	Very Likely	Very Likely	Very Likely	Very Likely	Very Likely	Very Likely	Very Likely
AHMEDABAD	Intensity	Light to Moderate rain/thundershowers at isolated places	Light to Moderate rain/thundershowers at isolated places	Light to Moderate rain/thundershowers at isolated places	Light thunders torm with lightning and surface wind 30-40 kmph accompanied with light to moderate rain at isolated places	Heavy rain at isolated places	Light to Moderate rain/thundershowers at a few places	Light to Moderate rain/thundershowers at a few places
	Probability	Very Likely	Very Likely	Very Likely	Very Likely	Likely	Very Likely	Very Likely
ANAND	Intensity	Light to Moderate rain/thundershowers at isolated places	Light to Moderate rain/thundershowers at isolated places	Light to Moderate rain/thundershowers at isolated places	Light thunders torm with lightning and surface wind 30-40 kmph accompanied with light to moderate rain at isolated places	Heavy rain at isolated places	Light to Moderate rain/thundershowers at a few places	Light to Moderate rain/thundershowers at a few places
	Probability	Very Likely	Very Likely	Very Likely	Very Likely	Likely	Very Likely	Very Likely
PANCHMAHAL	Intensity	Light to Moderate rain/thundershowers at isolated places	Light to Moderate rain/thundershowers at isolated places	Light to Moderate rain/thundershowers at isolated places	Light thunders torm with lightning and surface wind 30-40 kmph accompanied with	Heavy rain at isolated places	Light to Moderate rain/thundershowers at a few places	Light to Moderate rain/thundershowers at a few places

	ability	<i>Likely</i>	<i>Likely</i>	<i>Likely</i>	<i>Likely</i>	<i>Likely</i>	<i>Likely</i>	<i>Likely</i>
VADODARA	Intensity	<i>Light to Moderate rain/thundershowers at isolated places</i>	<i>Light to Moderate rain/thundershowers at isolated places</i>	<i>Light to Moderate rain/thundershowers at isolated places</i>	<i>Heavy rain at isolated places</i>	<i>Heavy rain at isolated places</i>	<i>Light to Moderate rain/thundershowers at a few places</i>	<i>Light to Moderate rain/thundershowers at a few places</i>
	Probability	<i>Very Likely</i>	<i>Very Likely</i>	<i>Very Likely</i>	<i>Likely</i>	<i>Likely</i>	<i>Very Likely</i>	<i>Very Likely</i>
CHHOTA UDEPUR	Intensity	<i>Light to Moderate rain/thundershowers at a few places</i>	<i>Light to Moderate rain/thundershowers at a few places</i>	<i>Light to Moderate rain/thundershowers at a few places</i>	<i>Heavy rain at isolated places</i>	<i>Heavy rain at isolated places</i>	<i>Light to Moderate rain/thundershowers at a few places</i>	<i>Light to Moderate rain/thundershowers at a few places</i>
	Probability	<i>Very Likely</i>	<i>Very Likely</i>	<i>Very Likely</i>	<i>Likely</i>	<i>Likely</i>	<i>Very Likely</i>	<i>Very Likely</i>
NARMADA	Intensity	<i>Light to Moderate rain/thundershowers at a few places</i>	<i>Light to Moderate rain/thundershowers at a few places</i>	<i>Light to Moderate rain/thundershowers at a few places</i>	<i>Heavy rain at isolated places</i>	<i>Heavy rain at isolated places</i>	<i>Light to Moderate rain/thundershowers at a few places</i>	<i>Light to Moderate rain/thundershowers at a few places</i>
	Probability	<i>Very Likely</i>	<i>Very Likely</i>	<i>Very Likely</i>	<i>Likely</i>	<i>Likely</i>	<i>Very Likely</i>	<i>Very Likely</i>
BHARUCH	Intensity	<i>Light to Moderate rain/thundershowers at isolated places</i>	<i>Light to Moderate rain/thundershowers at isolated places</i>	<i>Light to Moderate rain/thundershowers at isolated places</i>	<i>Heavy rain at isolated places</i>	<i>Heavy rain at isolated places</i>	<i>Heavy rain at isolated places</i>	<i>Light to Moderate rain/thundershowers at a few places</i>
	Probability	<i>Very Likely</i>	<i>Very Likely</i>	<i>Very Likely</i>	<i>Likely</i>	<i>Likely</i>	<i>Likely</i>	<i>Very Likely</i>
SURAT	Intensity	<i>Light to Moderate rain/thundershowers at isolated</i>	<i>Light to Moderate rain/thundershowers at isolated</i>	<i>Light to Moderate rain/thundershowers at isolated</i>	<i>Heavy rain at isolated places</i>	<i>Heavy rain at isolated places</i>	<i>Heavy rain at isolated places</i>	<i>Light to Moderate rain/thundershowers at a few</i>

		<i>places</i>	<i>places</i>	<i>places</i>				<i>places</i>
	Probability	<i>Very Likely</i>	<i>Very Likely</i>	<i>Very Likely</i>	<i>Likely</i>	<i>Likely</i>	<i>Likely</i>	<i>Very Likely</i>
DANGS	Intensity	<i>Light to Moderate rain/thundershowers at a few places</i>	<i>Light to Moderate rain/thundershowers at a few places</i>	<i>Light to Moderate rain/thundershowers at a few places</i>	<i>Heavy rain at isolated places</i>	<i>Heavy rain at isolated places</i>	<i>Heavy rain at isolated places</i>	<i>Light to Moderate rain/thundershowers at a few places</i>
	Probability	<i>Very Likely</i>	<i>Very Likely</i>	<i>Very Likely</i>	<i>Likely</i>	<i>Likely</i>	<i>Likely</i>	<i>Very Likely</i>
NAVSARI	Intensity	<i>Light to Moderate rain/thundershowers at a few places</i>	<i>Light to Moderate rain/thundershowers at a few places</i>	<i>Light to Moderate rain/thundershowers at a few places</i>	<i>Heavy rain at isolated places</i>	<i>Heavy rain at isolated places</i>	<i>Heavy rain at isolated places</i>	<i>Light to Moderate rain/thundershowers at a few places</i>
	Probability	<i>Very Likely</i>	<i>Very Likely</i>	<i>Very Likely</i>	<i>Likely</i>	<i>Likely</i>	<i>Likely</i>	<i>Very Likely</i>
VALSAD	Intensity	<i>Light to Moderate rain/thundershowers at a few places</i>	<i>Light to Moderate rain/thundershowers at a few places</i>	<i>Light to Moderate rain/thundershowers at a few places</i>	<i>Heavy rain at isolated places</i>	<i>Heavy rain at isolated places</i>	<i>Heavy rain at isolated places</i>	<i>Light to Moderate rain/thundershowers at a few places</i>
	Probability	<i>Very Likely</i>	<i>Very Likely</i>	<i>Very Likely</i>	<i>Likely</i>	<i>Likely</i>	<i>Likely</i>	<i>Very Likely</i>
TAPI	Intensity	<i>Light to Moderate rain/thundershowers at a few places</i>	<i>Light to Moderate rain/thundershowers at a few places</i>	<i>Light to Moderate rain/thundershowers at a few places</i>	<i>Heavy rain at isolated places</i>	<i>Heavy rain at isolated places</i>	<i>Heavy rain at isolated places</i>	<i>Light to Moderate rain/thundershowers at a few places</i>
	Probability	<i>Very Likely</i>	<i>Very Likely</i>	<i>Very Likely</i>	<i>Likely</i>	<i>Likely</i>	<i>Likely</i>	<i>Very Likely</i>
DAMAN	Intensity	<i>Light to Moderate rain/thundershow</i>	<i>Light to Moderate rain/thundershow</i>	<i>Light to Moderate rain/thundershow</i>	<i>Heavy rain at isolated places</i>	<i>Heavy rain at isolated places</i>	<i>Heavy rain at isolated places</i>	<i>Light to Moderate rain/thundershow</i>

		ers at a few places	ers at a few places	ers at a few places		place s		ers at a few places
	Prob ability	Very Likely	Very Likely	Very Likely	Likely	Likely	Likely	Very Likely
DADRA AND NAGAR HAVELI	Inten sity	Light to Moderate rain/thundershowers at a few places	Light to Moderate rain/thundershowers at a few places	Light to Moderate rain/thundershowers at a few places	Heavy rain at isolated places	Heavy rain at isolated places	Heavy rain at isolated places	Light to Moderate rain/thundershowers at a few places
	Prob ability	Very Likely	Very Likely	Very Likely	Likely	Likely	Likely	Very Likely
SURRENDRANAGAR	Inten sity	Light to Moderate rain/thundershowers at isolated places	Light to Moderate rain/thundershowers at isolated places	Light to Moderate rain/thundershowers at isolated places	Light thunders torm with lightning and surface wind 30-40 kmph accompanied with light to moderate rain at isolated places	Light thund erstor m with lightning and surfa ce wind 30-40 kmph accompan ied with light to mode rate rain at isolat ed place s	Light to Moderate rain/thundershowers at many places	Light to Moderate rain/thundershowers at a few places
	Prob ability	Very Likely	Very Likely	Very Likely	Very Likely	Very Likely	Very Likely	Very Likely
RAJKOT	Inten sity	Light to Moderate rain/thundershowers at isolated	Light to Moderate rain/thundershowers at isolated	Light to Moderate rain/thundershowers at isolated	Light to Moderate rain/thundershowers at a few	Heavy rain at isolated places	Heavy rain at isolated places	Light to Moderate rain/thundershowers at a few

		<i>places</i>	<i>places</i>	<i>places</i>	<i>places</i>			<i>places</i>
	Prob abilit y	Very Likely	Very Likely	Very Likely	Very Likely	Likely	Likely	Very Likely
JAMNAG AR	Inten sity	Light to Moderat e rain/thun dershow ers at isolated places	Light to Moderat e rain/thun dershow ers at isolated places	Light to Moderat e rain/thun dershow ers at isolated places	Light to Moderat e rain/thun dershow ers at isolated places	Light thund erstor m with lightni ng and surfa ce wind 30-40 kmph acco mpan ied with light to mode rate rain at isolat ed place s	Heavy rain at isolated places	Light to Moderat e rain/thun dershow ers at a few places
	Prob abilit y	Very Likely	Very Likely	Very Likely	Very Likely	Very Likely	Likely	Very Likely
PORBAN DAR	Inten sity	Light to Moderat e rain/thun dershow ers at isolated places	Light to Moderat e rain/thun dershow ers at isolated places	Light to Moderat e rain/thun dershow ers at isolated places	Light to Moderat e rain/thun dershow ers at isolated places	Heav y rain at isolat ed place s	Heavy rain at isolated places	Light to Moderat e rain/thun dershow ers at a few places
	Prob abilit y	Very Likely	Very Likely	Very Likely	Very Likely	Likely	Likely	Very Likely
JUNAGA DH	Inten sity	Light to Moderat e rain/thun dershow ers at isolated places	Light to Moderat e rain/thun dershow ers at isolated places	Light to Moderat e rain/thun dershow ers at isolated places	Light to Moderat e rain/thun dershow ers at a few places	Heav y rain at isolat ed place s	Heavy rain at isolated places	Light to Moderat e rain/thun dershow ers at a few places
	Prob	Very	Very	Very	Very	Likely	Likely	Very

	ability	Likely	Likely	Likely	Likely			Likely
AMRELI	Intensity	Light to Moderate rain/thundershowers at isolated places	Light to Moderate rain/thundershowers at isolated places	Light to Moderate rain/thundershowers at isolated places	Light thunders torm with lightning and surface wind 30-40 kmph accompanied with light to moderate rain at isolated places	Heavy rain at isolated places	Heavy rain at isolated places	Light to Moderate rain/thundershowers at a few places
	Probability	Very Likely	Very Likely	Very Likely	Very Likely	Likely	Likely	Very Likely
BHAVNAGAR	Intensity	Light to Moderate rain/thundershowers at isolated places	Light to Moderate rain/thundershowers at isolated places	Light to Moderate rain/thundershowers at isolated places	Light thunders torm with lightning and surface wind 30-40 kmph accompanied with light to moderate rain at isolated places	Heavy rain at isolated places	Heavy rain at isolated places	Light to Moderate rain/thundershowers at a few places
	Probability	Very Likely	Very Likely	Very Likely	Very Likely	Likely	Likely	Very Likely
MORBI	Intensity	Light to Moderate rain/thundershowers at isolated places	Light to Moderate rain/thundershowers at isolated places	Light to Moderate rain/thundershowers at isolated places	Light to Moderate rain/thundershowers at isolated places	Light thunderstorm with lightning and surface wind 30-40 kmph accompanied	Heavy rain at isolated places	Light to Moderate rain/thundershowers at a few places

						<i>with light to moderate rain at isolated places</i>		
	Probability	<i>Very Likely</i>	<i>Very Likely</i>	<i>Very Likely</i>	<i>Very Likely</i>	<i>Very Likely</i>	<i>Likely</i>	<i>Very Likely</i>
DEVBHOMI DWARKA	Intensity	<i>Light to Moderate rain/thundershowers at isolated places</i>	<i>Light to Moderate rain/thundershowers at isolated places</i>	<i>Light to Moderate rain/thundershowers at isolated places</i>	<i>Light to Moderate rain/thundershowers at isolated places</i>	<i>Light thunderstorm with lightning and surface wind 30-40 kmph accompanied with light to moderate rain at isolated places</i>	<i>Heavy rain at isolated places</i>	<i>Light to Moderate rain/thundershowers at a few places</i>
	Probability	<i>Very Likely</i>	<i>Very Likely</i>	<i>Very Likely</i>	<i>Very Likely</i>	<i>Very Likely</i>	<i>Likely</i>	<i>Very Likely</i>
GIR SOMNATH	Intensity	<i>Light to Moderate rain/thundershowers at isolated places</i>	<i>Light to Moderate rain/thundershowers at isolated places</i>	<i>Light to Moderate rain/thundershowers at isolated places</i>	<i>Light to Moderate rain/thundershowers at a few places</i>	<i>Heavy rain at isolated places</i>	<i>Heavy rain at isolated places</i>	<i>Light to Moderate rain/thundershowers at a few places</i>
	Probability	<i>Very Likely</i>	<i>Very Likely</i>	<i>Very Likely</i>	<i>Very Likely</i>	<i>Likely</i>	<i>Likely</i>	<i>Very Likely</i>

	y							
BOTAD	Inten sity	Light to Moderat e rain/thun dershow ers at isolated places	Light to Moderat e rain/thun dershow ers at isolated places	Light to Moderat e rain/thun dershow ers at isolated places	Light thunders torm with lightning and surface wind 30- 40 kmph accompa nied with light to moderat e rain at isolated places	Heav y rain at isolat ed place s	Heavy rain at isolated places	Light to Moderat e rain/thun dershow ers at a few places
	Prob abilit y	Very Likely	Very Likely	Very Likely	Very Likely	Likely	Likely	Very Likely
DIU	Inten sity	Light to Moderat e rain/thun dershow ers at isolated places	Light to Moderat e rain/thun dershow ers at isolated places	Light to Moderat e rain/thun dershow ers at isolated places	Light to Moderat e rain/thun dershow ers at a few places	Heav y rain at isolat ed place s	Heavy rain at isolated places	Light to Moderat e rain/thun dershow ers at a few places
	Prob abilit y	Very Likely	Very Likely	Very Likely	Very Likely	Likely	Likely	Very Likely
KUTCH	Inten sity	Light to Moderat e rain/thun dershow ers at isolated places	Light to Moderat e rain/thun dershow ers at isolated places	Light to Moderat e rain/thun dershow ers at isolated places	Light to Moderat e rain/thun dershow ers at isolated places	Light thund erstor m with lightni ng and surfa ce wind 30-40 kmph acco mpan ied with light to mode rate rain at isolat	Light to Moderat e rain/thun dershow ers at many places	Light to Moderat e rain/thun dershow ers at a few places

						ed place s		
	Prob abilit y	Very Likely	Very Likely	Very Likely	Very Likely	Very Likely	Very Likely	Very Likely

Warning Colour Codes	हिन्दी	मराठी
WARNING (TAKE ACTION)	चेतावनी	चेतावणी
ALERT (BE PREPARED)	सतर्क रहें	सतर्क असावे
WATCH (BE UPDATED)	निगरानी रखें	लक्षं असावे
NO WARNING (NO ACTION)	कोई चेतावनी नहीं	धोक्याची सूचना नाही



Heavy Rain



Heavy Snow



Thunderstorm



Dust Storm



Strong Winds



Visibility



Cyclone



Squall/ Hail



Frost



Cold Wave



Heat Wave



Sea State