

JEE1 OP 6th S2

JEE April 2024

| | |
|----------------|-------------------|
| Application No | |
| Candidate Name | |
| Roll No | |
| Test Date | 06/04/2024 |
| Test Time | 3:00 PM - 6:00 PM |
| Subject | B. Tech |

Section : Mathematics Section A

Q.1 $\lim_{n \rightarrow \infty} \frac{(1^2-1)(n-1) + (2^2-2)(n-2) + \dots + ((n-1)^2-(n-1)) \cdot 1}{(1^3+2^3+\dots+n^3)-(1^2+2^2+\dots+n^2)}$ is equal to :

Options

1. $\frac{2}{3}$
2. $\frac{3}{4}$
3. $\frac{1}{3}$
4. $\frac{1}{2}$

Question Type : MCQ

Question ID : 87827055886

Option 1 ID : 878270219625

Option 2 ID : 878270219626

Option 3 ID : 878270219624

Option 4 ID : 878270219623

Status : Answered

Chosen Option : 3

Q.2 Let ABC be an equilateral triangle. A new triangle is formed by joining the middle points of all sides of the triangle ABC and the same process is repeated infinitely many times. If P is the sum of perimeters and Q is be the sum of areas of all the triangles formed in this process, then :

Options

1. $P=36\sqrt{3}Q^2$
2. $P^2=6\sqrt{3}Q$
3. $P^2=72\sqrt{3}Q$
4. $P^2=36\sqrt{3}Q$

Question Type : MCQ

Question ID : 87827055893

Option 1 ID : 878270219651

Option 2 ID : 878270219652

Option 3 ID : 878270219654

Option 4 ID : 878270219653

Status : Answered

Chosen Option : 4

Q.3 Suppose the solution of the differential equation $\frac{dy}{dx} = \frac{(2 + \alpha)x - \beta y + 2}{\beta x - 2\alpha y - (\beta\gamma - 4\alpha)}$ represents a circle passing through origin. Then the radius of this circle is :

Options

1. $\frac{1}{2}$
2. 2
3. $\frac{\sqrt{17}}{2}$
4. $\sqrt{17}$

Question Type : **MCQ**

Question ID : **87827055890**

Option 1 ID : **878270219642**

Option 2 ID : **878270219640**

Option 3 ID : **878270219639**

Option 4 ID : **878270219641**

Status : **Not Answered**

Chosen Option : --

Q.4 If A is a square matrix of order 3 such that $\det(A) = 3$ and $\det(\text{adj}(-4 \text{ adj}(-3 \text{ adj}(3 \text{ adj}((2A)^{-1})))) = 2^m 3^n$, then $m + 2n$ is equal to :

Options

1. 4
2. 6
3. 2
4. 3

Question Type : **MCQ**

Question ID : **87827055881**

Option 1 ID : **878270219604**

Option 2 ID : **878270219606**

Option 3 ID : **878270219603**

Option 4 ID : **878270219605**

Status : **Answered**

Chosen Option : 1

Q.5

If z_1, z_2 are two distinct complex number such that $\left| \frac{z_1 - 2z_2}{\frac{1}{2} - z_1 \bar{z}_2} \right| = 2$, then

Options

1. both z_1 and z_2 lie on the same circle.
2. either z_1 lies on a circle of radius 1 or z_2 lies on a circle of radius $\frac{1}{2}$.
3. z_1 lies on a circle of radius $\frac{1}{2}$ and z_2 lies on a circle of radius 1.
4. either z_1 lies on a circle of radius $\frac{1}{2}$ or z_2 lies on a circle of radius 1.

Question Type : MCQ

Question ID : 87827055880

Option 1 ID : 878270219599

Option 2 ID : 878270219602

Option 3 ID : 878270219600

Option 4 ID : 878270219601

Status : Not Answered

Chosen Option : --

Q.6

If all the words with or without meaning made using all the letters of the word "NAGPUR" are arranged as in a dictionary, then the word at 315th position in this arrangement is :

Options

1. NRAGPU
2. NRAPUG
3. NRAPGU
4. NRAGUP

Question Type : MCQ

Question ID : 87827055882

Option 1 ID : 878270219607

Option 2 ID : 878270219610

Option 3 ID : 878270219608

Option 4 ID : 878270219609

Status : Answered

Chosen Option : 3

Q.7

Let $\vec{a} = 6\hat{i} + \hat{j} - \hat{k}$ and $\vec{b} = \hat{i} + \hat{j}$. If \vec{c} is a vector such that $|\vec{c}| \geq 6$, $\vec{a} \cdot \vec{c} = 6|\vec{c}|$, $|\vec{c} - \vec{a}| = 2\sqrt{2}$ and

the angle between $\vec{a} \times \vec{b}$ and \vec{c} is 60° , then $\left| \left(\vec{a} \times \vec{b} \right) \times \vec{c} \right|$ is equal to :

Options

1. $\frac{9}{2}(6 + \sqrt{6})$
2. $\frac{9}{2}(6 - \sqrt{6})$
3. $\frac{3}{2}\sqrt{3}$
4. $\frac{3}{2}\sqrt{6}$

Question Type : MCQ

Question ID : 87827055896

Option 1 ID : 878270219666

Option 2 ID : 878270219665

Option 3 ID : 878270219663

Option 4 ID : 878270219664

Status : Answered

Chosen Option : 1

Q.8

Suppose for a differentiable function h , $h(0) = 0$, $h(1) = 1$ and $h'(0) = h'(1) = 2$. If $g(x) = h(e^x)e^{h(x)}$, then $g'(0)$ is equal to :

Options

1. 3
2. 5
3. 8
4. 4

Question Type : MCQ

Question ID : 87827055887

Option 1 ID : 878270219627

Option 2 ID : 878270219629

Option 3 ID : 878270219630

Option 4 ID : 878270219628

Status : Answered

Chosen Option : 4

Q.9

If the function $f(x) = \left(\frac{1}{x}\right)^{2x}$; $x > 0$ attains the maximum value at $x = \frac{1}{e}$ then :

Options

1. $(2e)^\pi > \pi^{(2e)}$
2. $e^\pi < \pi^e$
3. $e^{2\pi} < (2\pi)^e$
4. $e^\pi > \pi^e$

Question Type : MCQ

Question ID : 87827055885

Option 1 ID : 878270219622

Option 2 ID : 878270219620

Option 3 ID : 878270219621

Option 4 ID : 878270219619

Status : Not Answered

Chosen Option : --

Q.10

If the area of the region $\left\{(x, y) : \frac{a}{x^2} \leq y \leq \frac{1}{x}, 1 \leq x \leq 2, 0 < a < 1\right\}$ is $(\log_e 2) - \frac{1}{7}$ then the value of $7a - 3$ is equal to :

Options

1. -1
2. 1
3. 2
4. 0

Question Type : MCQ

Question ID : 87827055888

Option 1 ID : 878270219631

Option 2 ID : 878270219633

Option 3 ID : 878270219634

Option 4 ID : 878270219632

Status : Answered

Chosen Option : 1

Q.11

Let $f(x) = \frac{1}{7 - \sin 5x}$ be a function defined on \mathbf{R} . Then the range of the function $f(x)$ is equal to :

Options

1. $\left[\frac{1}{7}, \frac{1}{6}\right]$

2. $\left[\frac{1}{8}, \frac{1}{6}\right]$

3. $\left[\frac{1}{7}, \frac{1}{5}\right]$

4. $\left[\frac{1}{8}, \frac{1}{5}\right]$

Question Type : **MCQ**

Question ID : **87827055878**

Option 1 ID : **878270219593**

Option 2 ID : **878270219594**

Option 3 ID : **878270219592**

Option 4 ID : **878270219591**

Status : **Answered**

Chosen Option : **2**

Q.12

If P(6, 1) be the orthocentre of the triangle whose vertices are A (5, -2), B(8, 3) and C (h, k), then the point C lies on the circle :

Options

1. $x^2 + y^2 - 61 = 0$

2. $x^2 + y^2 - 74 = 0$

3. $x^2 + y^2 - 52 = 0$

4. $x^2 + y^2 - 65 = 0$

Question Type : **MCQ**

Question ID : **87827055892**

Option 1 ID : **878270219648**

Option 2 ID : **878270219650**

Option 3 ID : **878270219647**

Option 4 ID : **878270219649**

Status : **Answered**

Chosen Option : **4**

Q.13

Let $\vec{a} = 2\hat{i} + \hat{j} - \hat{k}$, $\vec{b} = \left(\left(\vec{a} \times (\hat{i} + \hat{j}) \right) \times \hat{i} \right) \times \hat{i}$. Then the square of the projection of \vec{a} on \vec{b} is :

Options

1. 2
2. $\frac{1}{3}$
3. $\frac{1}{5}$
4. $\frac{2}{3}$

Question Type : MCQ

Question ID : 87827055895

Option 1 ID : 878270219660

Option 2 ID : 878270219659

Option 3 ID : 878270219662

Option 4 ID : 878270219661

Status : Answered

Chosen Option : 1

Q.14

Let P (α , β , γ) be the image of the point Q (3, -3, 1) in the line $\frac{x-0}{1} = \frac{y-3}{1} = \frac{z-1}{-1}$ and R be the point (2, 5, -1). If the area of the triangle PQR is λ and $\lambda^2 = 14K$, then K is equal to :

Options

1. 81
2. 36
3. 18
4. 72

Question Type : MCQ

Question ID : 87827055894

Option 1 ID : 878270219658

Option 2 ID : 878270219656

Option 3 ID : 878270219655

Option 4 ID : 878270219657

Status : Answered

Chosen Option : 1

Q.15 Let $A = \{1, 2, 3, 4, 5\}$. Let R be a relation on A defined by xRy if and only if $4x \leq 5y$. Let m be the number of elements in R and n be the minimum number of elements from $A \times A$ that are required to be added to R to make it a symmetric relation. Then $m + n$ is equal to :

Options

1. 23
2. 25
3. 26
4. 24

Question Type : MCQ

Question ID : 87827055879

Option 1 ID : 878270219595

Option 2 ID : 878270219597

Option 3 ID : 878270219598

Option 4 ID : 878270219596

Status : Answered

Chosen Option : 2

Q.16 If three letters can be posted to any one of the 5 different addresses, then the probability that the three letters are posted to exactly two addresses is :

Options

1. $\frac{12}{25}$
2. $\frac{4}{25}$
3. $\frac{6}{25}$
4. $\frac{18}{25}$

Question Type : MCQ

Question ID : 87827055897

Option 1 ID : 878270219669

Option 2 ID : 878270219667

Option 3 ID : 878270219668

Option 4 ID : 878270219670

Status : Not Answered

Chosen Option : --

Q.17 If the locus of the point, whose distances from the point (2, 1) and (1, 3) are in the ratio 5 : 4, is $ax^2 + by^2 + cxy + dx + ey + 170 = 0$, then the value of $a^2 + 2b + 3c + 4d + e$ is equal to :

Options

1. 5
2. -27
3. 437
4. 37

Question Type : MCQ

Question ID : 87827055891

Option 1 ID : 878270219643

Option 2 ID : 878270219645

Option 3 ID : 878270219644

Option 4 ID : 878270219646

Status : Answered

Chosen Option : 4

Q.18 A software company sets up m number of computer systems to finish an assignment in 17 days. If 4 computer systems crashed on the start of the second day, 4 more computer systems crashed on the start of the third day and so on, then it took 8 more days to finish the assignment. The value of m is equal to :

Options

1. 180
2. 125
3. 150
4. 160

Question Type : MCQ

Question ID : 87827055884

Option 1 ID : 878270219618

Option 2 ID : 878270219616

Option 3 ID : 878270219615

Option 4 ID : 878270219617

Status : Not Answered

Chosen Option : --

Q.19

If $\int \frac{1}{a^2 \sin^2 x + b^2 \cos^2 x} dx = \frac{1}{12} \tan^{-1}(3 \tan x) + \text{constant}$, then the maximum value of $a \sin x + b \cos x$, is :

Options

1. $\sqrt{42}$
2. $\sqrt{39}$
3. $\sqrt{41}$
4. $\sqrt{40}$

Question Type : **MCQ**

Question ID : **87827055889**

Option 1 ID : **878270219635**

Option 2 ID : **878270219636**

Option 3 ID : **878270219638**

Option 4 ID : **878270219637**

Status : **Answered**

Chosen Option : **4**

Q.20

Let $0 \leq r \leq n$. If ${}^{n+1}C_{r+1} : {}^nC_r : {}^{n-1}C_{r-1} = 55 : 35 : 21$, then $2n + 5r$ is equal to :

Options

1. **62**
2. 60
3. 55
4. 50

Question Type : **MCQ**

Question ID : **87827055883**

Option 1 ID : **878270219614**

Option 2 ID : **878270219613**

Option 3 ID : **878270219612**

Option 4 ID : **878270219611**

Status : **Answered**

Chosen Option : **4**

Section : **Mathematics Section B**

Q.21

If the shortest distance between the lines $\frac{x-\lambda}{3} = \frac{y-2}{-1} = \frac{z-1}{1}$ and $\frac{x+2}{-3} = \frac{y+5}{2} = \frac{z-4}{4}$ is $\frac{44}{\sqrt{30}}$, then the largest possible value of $|\lambda|$ is equal to _____.

Given **43.00**

Answer :

Question Type : **SA**

Question ID : **87827055905**

Status : **Answered**

Q.22 Let $[t]$ denote the largest integer less than or equal to t . If

$$\int_0^3 \left([x^2] + \left\lceil \frac{x^2}{2} \right\rceil \right) dx = a + b\sqrt{2} - \sqrt{3} - \sqrt{5} + c\sqrt{6} - \sqrt{7}, \text{ where } a, b, c \in \mathbf{Z}, \text{ then } a + b + c \text{ is equal to}$$

_____.

Given --

Answer :

Question Type : SA

Question ID : 87827055902

Status : Not Answered

Q.23

Let α, β be roots of $x^2 + \sqrt{2}x - 8 = 0$. If $U_n = \alpha^n + \beta^n$, then $\frac{U_{10} + \sqrt{2}U_9}{2U_8}$ is equal to _____.

Given 04.00

Answer :

Question Type : SA

Question ID : 87827055898

Status : Answered

Q.24

In a triangle ABC, $BC = 7$, $AC = 8$, $AB = \alpha \in \mathbf{N}$ and $\cos A = \frac{2}{3}$. If $49\cos(3C) + 42 = \frac{m}{n}$, where $\gcd(m, n) = 1$, then $m + n$ is equal to _____.

Given 43.00

Answer :

Question Type : SA

Question ID : 87827055907

Status : Answered

Q.25

The length of the latus rectum and directrices of a hyperbola with eccentricity e are 9 and $x = \pm \frac{4}{\sqrt{3}}$, respectively. Let the line $y - \sqrt{3}x + \sqrt{3} = 0$ touch this hyperbola at (x_0, y_0) . If m is the product of the focal distances of the point (x_0, y_0) , then $4e^2 + m$ is equal to _____.

Given --

Answer :

Question Type : SA

Question ID : 87827055904

Status : Not Answered

Q.26

If $S(x) = (1+x) + 2(1+x)^2 + 3(1+x)^3 + \dots + 60(1+x)^{60}$, $x \neq 0$, and $(60)^2 S(60) = a(b)^b + b$, where $a, b \in \mathbf{N}$, then $(a+b)$ equal to _____.

Given --

Answer :

Question Type : SA

Question ID : 87827055900

Status : Not Answered

Q.27 If the system of equations

$$2x + 7y + \lambda z = 3$$

$$3x + 2y + 5z = 4$$

$$x + \mu y + 32z = -1$$

has infinitely many solutions, then $(\lambda - \mu)$ is equal to _____ :

Given **38.00**

Answer :

Question Type : **SA**

Question ID : **87827055899**

Status : **Answered**

Q.28 Let $[t]$ denote the greatest integer less than or equal to t . Let $f: [0, \infty) \rightarrow \mathbf{R}$ be a function defined by

$$f(x) = \left[\frac{x}{2} + 3 \right] - [\sqrt{x}].$$
 Let S be the set of all points in the interval $[0, 8]$ at which f is not continuous.

Then $\sum_{a \in S} a$ is equal to _____.

Given --

Answer :

Question Type : **SA**

Question ID : **87827055901**

Status : **Not Answered**

Q.29 If the solution $y(x)$ of the given differential equation $(e^y + 1) \cos x \, dx + e^y \sin x \, dy = 0$ passes through

the point $\left(\frac{\pi}{2}, 0\right)$, then the value of $e^{y\left(\frac{\pi}{6}\right)}$ is equal to _____.

Given **03.00**

Answer :

Question Type : **SA**

Question ID : **87827055903**

Status : **Answered**

Q.30 From a lot of 12 items containing 3 defectives, a sample of 5 items is drawn at random. Let the random variable X denote the number of defective items in the sample. Let items in the sample be drawn one by one without replacement. If variance of X is $\frac{m}{n}$, where $\gcd(m, n) = 1$, then $n - m$ is equal to _____.

Given --

Answer :

Question Type : **SA**

Question ID : **87827055906**

Status : **Not Answered**

Q.31 A body projected vertically upwards with a certain speed from the top of a tower reaches the ground in t_1 . If it is projected vertically downwards from the same point with the same speed, it reaches the ground in t_2 . Time required to reach the ground, if it is dropped from the top of the tower, is :

Options

1. $\sqrt{t_1 - t_2}$
2. $\sqrt{t_1 t_2}$
3. $\sqrt{\frac{t_1}{t_2}}$
4. $\sqrt{t_1 + t_2}$

Question Type : MCQ

Question ID : 87827055910

Option 1 ID : 878270219691

Option 2 ID : 878270219689

Option 3 ID : 878270219692

Option 4 ID : 878270219690

Status : Answered

Chosen Option : 2

Q.32 A car of 800 kg is taking turn on a banked road of radius 300 m and angle of banking 30° . If coefficient of static friction is 0.2 then the maximum speed with which car can negotiate the turn safely : ($g = 10 \text{ m/s}^2$, $\sqrt{3} = 1.73$)

Options

1. 102.8 m/s
2. 51.4 m/s
3. 264 m/s
4. 70.4 m/s

Question Type : MCQ

Question ID : 87827055908

Option 1 ID : 878270219683

Option 2 ID : 878270219681

Option 3 ID : 878270219682

Option 4 ID : 878270219684

Status : Answered

Chosen Option : 2

Q.33 For the thin convex lens, the radii of curvature are at 15 cm and 30 cm respectively. The focal length the lens is 20 cm. The refractive index of the material is :

Options

1. 1.5
2. 1.4
3. 1.8
4. 1.2

Question Type : MCQ

Question ID : 87827055923

Option 1 ID : 878270219742

Option 2 ID : 878270219741

Option 3 ID : 878270219744

Option 4 ID : 878270219743

Status : Answered

Chosen Option : 1

Q.34 Assuming the earth to be a sphere of uniform mass density, a body weighed 300 N on the surface of earth. How much it would weigh at $R/4$ depth under surface of earth ?

Options

1. 300 N
2. 225 N
3. 375 N
4. 75 N

Question Type : MCQ

Question ID : 87827055914

Option 1 ID : 878270219707

Option 2 ID : 878270219706

Option 3 ID : 878270219708

Option 4 ID : 878270219705

Status : Answered

Chosen Option : 2

Q.35 A total of 48 J heat is given to one mole of helium kept in a cylinder. The temperature of helium increases by 2°C. The work done by the gas is :
Given, $R = 8.3 \text{ J K}^{-1} \text{ mol}^{-1}$.

Options

1. 23.1 J
2. 24.9 J
3. 48 J
4. 72.9 J

Question Type : **MCQ**

Question ID : **87827055916**

Option 1 ID : **878270219716**

Option 2 ID : **878270219713**

Option 3 ID : **878270219714**

Option 4 ID : **878270219715**

Status : **Answered**

Chosen Option : **1**

Q.36 Two identical conducting spheres P and S with charge Q on each, repel each other with a force 16 N. A third identical uncharged conducting sphere R is successively brought in contact with the two spheres. The new force of repulsion between P and S is :

Options

1. 1 N
2. 12 N
3. 4 N
4. 6 N

Question Type : **MCQ**

Question ID : **87827055919**

Option 1 ID : **878270219728**

Option 2 ID : **878270219726**

Option 3 ID : **878270219727**

Option 4 ID : **878270219725**

Status : **Answered**

Chosen Option : **3**

Q.37 When UV light of wavelength 300 nm is incident on the metal surface having work function 2.13 eV, electron emission takes place. The stopping potential is :
(Given $hc = 1240 \text{ eV nm}$)

Options

1. 4.1 V
2. 4 V
3. 2 V
4. 1.5 V

Question Type : MCQ

Question ID : 87827055924

Option 1 ID : 878270219745

Option 2 ID : 878270219748

Option 3 ID : 878270219747

Option 4 ID : 878270219746

Status : Answered

Chosen Option : 3

Q.38 Given below are two statements :

Statement (I) : Dimensions of specific heat is $[L^2 T^{-2} K^{-1}]$.

Statement (II) : Dimensions of gas constant is $[M L^2 T^{-1} K^{-1}]$.

In the light of the above statements, choose the **most appropriate** answer from the options given below.

Options

1. Both **statement (I)** and **statement (II)** are incorrect
2. Both **statement (I)** and **statement (II)** are correct
3. **Statement (I)** is incorrect but **statement (II)** is correct
4. **Statement (I)** is correct but **statement (II)** is incorrect

Question Type : MCQ

Question ID : 87827055909

Option 1 ID : 878270219686

Option 2 ID : 878270219685

Option 3 ID : 878270219688

Option 4 ID : 878270219687

Status : Answered

Chosen Option : 1

Q.39 In a coil, the current changes from -2 A to $+2\text{ A}$ in 0.2 s and induces an emf of 0.1 V . The self inductance of the coil is :

Options

1. 1 mH
2. 5 mH
3. 4 mH
4. 2.5 mH

Question Type : **MCQ**

Question ID : **87827055921**

Option 1 ID : **878270219733**

Option 2 ID : **878270219736**

Option 3 ID : **878270219735**

Option 4 ID : **878270219734**

Status : **Answered**

Chosen Option : **2**

Q.40 Match List-I with List-II :

List-I

Y vs X


List-II

Shape of Graph

(A) Y = magnetic susceptibility

X = magnetising field


(I)



(B) Y = magnetic field

X = distance from centre of a current carrying wire for $x < a$
(where a = radius of wire)


(II)



(C) Y = magnetic field

X = distance from centre of a current carrying wire for $x > a$
(where a = radius of wire)


(III)



(D) Y = magnetic field inside solenoid

X = distance from centre

(IV)



Choose the **correct** answer from the options given below :

Options

1. (A)-(I), (B)-(III), (C)-(II), (D)-(IV)
2. (A)-(III), (B)-(I), (C)-(IV), (D)-(II)
3. (A)-(IV), (B)-(I), (C)-(III), (D)-(II)
4. (A)-(III), (B)-(IV), (C)-(I), (D)-(II)

Question Type : MCQ

Question ID : 87827055912

Option 1 ID : 878270219699

Option 2 ID : 878270219698

Option 3 ID : 878270219697

Option 4 ID : 878270219700

Status : Answered

- Q.41** In finding out refractive index of glass slab the following observations were made through travelling microscope 50 vernier scale division = 49 MSD; 20 divisions on main scale in each cm
- For mark on paper
MSR = 8.45 cm, VC=26
- For mark on paper seen through slab
MSR = 7.12 cm, VC=41
- For powder particle on the top surface of the glass slab
MSR = 4.05 cm, VC=1
- (MSR = Main Scale Reading, VC = Vernier Coincidence)
- Refractive index of the glass slab is :

Options

1. 1.52
2. 1.42
3. 1.24
4. 1.35

Question Type : MCQ

Question ID : 87827055927

Option 1 ID : 878270219760

Option 2 ID : 878270219758

Option 3 ID : 878270219757

Option 4 ID : 878270219759

Status : Not Answered

Chosen Option : --

- Q.42** The acceptor level of a p-type semiconductor is 6 eV. The maximum wavelength of light which can create a hole would be : Given $hc = 1242 \text{ eV nm}$.

Options

1. 103.5 nm
2. 407 nm
3. 414 nm
4. 207 nm

Question Type : MCQ

Question ID : 87827055926

Option 1 ID : 878270219756

Option 2 ID : 878270219754

Option 3 ID : 878270219755

Option 4 ID : 878270219753

Status : Answered

Chosen Option : 4

Q.43 In a vernier calliper, when both jaws touch each other, zero of the vernier scale shifts towards left and its 4th division coincides exactly with a certain division on main scale. If 50 vernier scale divisions equal to 49 main scale divisions and zero error in the instrument is 0.04 mm then how many main scale divisions are there in 1 cm ?

Options

1. 5
2. 20
3. 40
4. 10

Question Type : MCQ

Question ID : 87827055917

Option 1 ID : 878270219720

Option 2 ID : 878270219718

Option 3 ID : 878270219719

Option 4 ID : 878270219717

Status : Answered

Chosen Option : 2

Q.44 A body of weight 200 N is suspended from a tree branch through a chain of mass 10 kg. The branch pulls the chain by a force equal to (if $g = 10 \text{ m/s}^2$) :

Options

1. 100 N
2. 200 N
3. 150 N
4. 300 N

Question Type : MCQ

Question ID : 87827055911

Option 1 ID : 878270219694

Option 2 ID : 878270219693

Option 3 ID : 878270219696

Option 4 ID : 878270219695

Status : Answered

Chosen Option : 4

Q.45 Energy of 10 non rigid diatomic molecules at temperature T is :

Options

1. $35 RT$
2. $70 K_B T$
3. $35 K_B T$
4. $\frac{7}{2} RT$

Question Type : MCQ

Question ID : 87827055918

Option 1 ID : 878270219721

Option 2 ID : 878270219722

Option 3 ID : 878270219724

Option 4 ID : 878270219723

Status : Answered

Chosen Option : 3

Q.46 The number of electrons flowing per second in the filament of a 110 W bulb operating at 220 V is :
(Given $e = 1.6 \times 10^{-19} \text{ C}$)

Options

1. 6.25×10^{17}
2. 6.25×10^{18}
3. 31.25×10^{17}
4. 1.25×10^{19}

Question Type : MCQ

Question ID : 87827055920

Option 1 ID : 878270219729

Option 2 ID : 878270219732

Option 3 ID : 878270219731

Option 4 ID : 878270219730

Status : Answered

Chosen Option : 3

Q.47 The longest wavelength associated with Paschen series is : (Given $R_H = 1.097 \times 10^7$ SI unit)

Options

1. $1.094 \times 10^{-6} \text{ m}$
2. $1.876 \times 10^{-6} \text{ m}$
3. $3.646 \times 10^{-6} \text{ m}$
4. $2.973 \times 10^{-6} \text{ m}$

Question Type : MCQ

Question ID : 87827055925

Option 1 ID : 878270219751

Option 2 ID : 878270219750

Option 3 ID : 878270219752

Option 4 ID : 878270219749

Status : Answered

Chosen Option : 2

Q.48 When kinetic energy of a body becomes 36 times of its original value, the percentage increase in the momentum of the body will be :

Options

1. 500%
2. 6%
3. 60%
4. 600%

Question Type : MCQ

Question ID : 87827055913

Option 1 ID : 878270219702

Option 2 ID : 878270219701

Option 3 ID : 878270219703

Option 4 ID : 878270219704

Status : Answered

Chosen Option : 1

Q.49 Pressure inside a soap bubble is greater than the pressure outside by an amount :
(given : R = Radius of bubble, S = Surface tension of bubble)

Options

1. $\frac{2S}{R}$
2. $\frac{S}{R}$
3. $\frac{4S}{R}$
4. $\frac{4R}{S}$

Question Type : MCQ

Question ID : 87827055915

Option 1 ID : 878270219709

Option 2 ID : 878270219711

Option 3 ID : 878270219710

Option 4 ID : 878270219712

Status : Answered

Chosen Option : 1

Q.50 In the given electromagnetic wave $E_y = 600 \sin(\omega t - kx) \text{ Vm}^{-1}$, intensity of the associated light beam is (in W/m^2) : (Given $\epsilon_0 = 9 \times 10^{-12} \text{ C}^2\text{N}^{-1}\text{m}^{-2}$)

Options

1. 972
2. 486
3. 243
4. 729

Question Type : MCQ

Question ID : 87827055922

Option 1 ID : 878270219738

Option 2 ID : 878270219740

Option 3 ID : 878270219739

Option 4 ID : 878270219737

Status : Answered

Chosen Option : 2

Section : Physics Section B

Q.51 Two coherent monochromatic light beams of intensities I and 4 I are superimposed. The difference between maximum and minimum possible intensities in the resulting beam is x I. The value of x is _____.

Given 08.00

Answer :

Question Type : SA

Question ID : 87827055931

Status : Answered

Q.52 A coil having 100 turns, area of $5 \times 10^{-3} \text{ m}^2$, carrying current of 1 mA is placed in uniform magnetic field of 0.20 T such a way that plane of coil is perpendicular to the magnetic field. The work done in turning the coil through 90° is _____ μJ .

Given **100.00**

Answer :

Question Type : SA

Question ID : 87827055935

Status : Answered

Q.53 For a given series LCR circuit it is found that maximum current is drawn when value of variable capacitance is 2.5 nF. If resistance of 200Ω and 100 mH inductor is being used in the given circuit. The frequency of ac source is _____ $\times 10^3 \text{ Hz}$. (given $\pi^2 = 10$)

Given **10.00**

Answer :

Question Type : SA

Question ID : 87827055936

Status : Answered

Q.54 A particle moves in a straight line so that its displacement x at any time t is given by $x^2 = 1 + t^2$. Its acceleration at any time t is x^{-n} where $n =$ _____.

Given --


Answer :

Question Type : SA

Question ID : 87827055934

Status : Not Answered

Q.55 In the given figure an ammeter A consists of a 240Ω coil connected in parallel to a 10Ω shunt. The reading of the ammeter is _____ mA.



Given **160.00**


Answer :

Question Type : SA

Question ID : 87827055930

Status : Answered

Q.56 A capacitor of $10\mu\text{F}$ capacitance whose plates are separated by 10 mm through air and each plate has area 4 cm^2 is now filled equally with two dielectric media of $K_1 = 2$, $K_2 = 3$ respectively as shown in figure. If new force between the plates is 8 N. The supply voltage is _____ V.



Given --


Answer :

Question Type : SA

Question ID : 87827055929

Status : Not Answered

- Q.57** A wire of cross sectional area A , modulus of elasticity $2 \times 10^{11} \text{ Nm}^{-2}$ and length 2 m is stretched between two vertical rigid supports. When a mass of 2 kg is suspended at the middle it sags lower from its original position making angle $\theta = \frac{1}{100}$ radian on the points of support. The value of A is $\text{---} \times 10^{-4} \text{ m}^2$ (consider $x \ll L$).
(given : $g = 10 \text{ m/s}^2$)



Given --
Answer :

Question Type : SA
Question ID : 87827055937
Status : Not Answered

- Q.58** Three balls of masses 2 kg , 4 kg and 6 kg respectively are arranged at centre of the edges of an equilateral triangle of side 2 m . The moment of inertia of the system about an axis through the centroid and perpendicular to the plane of triangle, will be $\text{---} \text{ kg m}^2$.

Given --
Answer :

Question Type : SA
Question ID : 87827055933
Status : Not Answered

- Q.59** Two open organ pipes of lengths 60 cm and 90 cm resonate at 6^{th} and 5^{th} harmonics respectively. The difference of frequencies for the given modes is $\text{---} \text{ Hz}$.
(Velocity of sound in air = 333 m/s)

Given **740.00**
Answer :


Question Type : SA
Question ID : 87827055928
Status : Answered

- Q.60** In Franck-Hertz experiment, the first dip in the current-voltage graph for hydrogen is observed at 10.2 V . The wavelength of light emitted by hydrogen atom when excited to the first excitation level is $\text{---} \text{ nm}$. (Given $hc = 1245 \text{ eV nm}$, $e = 1.6 \times 10^{-19} \text{ C}$).

Given --
Answer :





Question Type : SA
Question ID : 87827055932
Status : Not Answered

Q.61



Consider the above chemical reaction. Product "A" is :

Options



Question Type : MCQ

Question ID : 87827055953

Option 1 ID : 878270219831

Option 2 ID : 878270219832

Option 3 ID : 878270219833

Option 4 ID : 878270219834

Status : Answered

Chosen Option : 1

Q.62 Match List - I with List - II.

List - I

Reaction

- (A) $\text{N}_2(\text{g}) + \text{O}_2(\text{g}) \rightarrow 2\text{NO}(\text{g})$
(B) $2\text{Pb}(\text{NO}_3)_2(\text{s}) \rightarrow 2\text{PbO}(\text{s}) + 4\text{NO}_2(\text{g}) + \text{O}_2(\text{g})$
(C) $2\text{Na}(\text{s}) + 2\text{H}_2\text{O}(\text{l}) \rightarrow 2\text{NaOH}(\text{aq}) + \text{H}_2(\text{g})$
(D) $2\text{NO}_2(\text{g}) + 2^-\text{OH}(\text{aq.}) \rightarrow \text{NO}_2^-(\text{aq.}) + \text{NO}_3^-(\text{aq.}) + \text{H}_2\text{O}(\text{l})$

List - II

Type of redox reaction

- (I) Decomposition
(II) Displacement
(III) Disproportionation
(IV) Combination

Choose the **correct** answer from the options given below :

Options

1. (A)-(II), (B)-(III), (C)-(IV), (D)-(I)
2. (A)-(IV), (B)-(I), (C)-(II), (D)-(III)
3. (A)-(III), (B)-(II), (C)-(I), (D)-(IV)
4. (A)-(I), (B)-(II), (C)-(III), (D)-(IV)

Question Type : **MCQ**

Question ID : **87827055941**

Option 1 ID : **878270219784**

Option 2 ID : **878270219783**

Option 3 ID : **878270219785**

Option 4 ID : **878270219786**

Status : **Answered**

Chosen Option : **2**

Q.63 Given below are two statements :

Statement I : PF_5 and BrF_5 both exhibit sp^3d hybridisation.

Statement II : Both SF_6 and $[\text{Co}(\text{NH}_3)_6]^{3+}$ exhibit sp^3d^2 hybridisation.

In the light of the above statements, choose the **correct** answer from the options given below :

Options

1. **Statement I** is false but **Statement II** is true
2. Both **Statement I** and **Statement II** are false
3. Both **Statement I** and **Statement II** are true
4. **Statement I** is true but **Statement II** is false

Question Type : **MCQ**

Question ID : **87827055939**

Option 1 ID : **878270219778**

Option 2 ID : **878270219776**


Option 3 ID : **878270219775**

Option 4 ID : **878270219777**

Status : **Answered**

Chosen Option : **2**

Q.64



The **correct** arrangement for decreasing order of electrophilic substitution for above compounds is :

Options

1. (III) > (I) > (II) > (IV)
2. (IV) > (I) > (II) > (III)
3. (III) > (IV) > (II) > (I)
4. (II) > (IV) > (III) > (I)

Question Type : MCQ

Question ID : 87827055951

Option 1 ID : 878270219823

Option 2 ID : 878270219824

Option 3 ID : 878270219825

Option 4 ID : 878270219826

Status : Answered

Chosen Option : 1

Q.65

During the detection of acidic radical present in a salt, a student gets a pale yellow precipitate soluble with difficulty in NH_4OH solution when sodium carbonate extract was first acidified with dil. HNO_3 and then AgNO_3 solution was added. This indicates presence of :

Options

1. CO_3^{2-}
2. I^-
3. Br^-
4. Cl^-

Question Type : MCQ

Question ID : 87827055949

Option 1 ID : 878270219818

Option 2 ID : 878270219817

Option 3 ID : 878270219816

Option 4 ID : 878270219815

Status : Answered

Chosen Option : 3

Q.66 The incorrect statement regarding the geometrical isomers of 2-butene is :

Options

1. cis-2-butene and trans-2-butene are not interconvertible at room temperature.
2. cis-2-butene and trans-2-butene are stereoisomers.
3. trans-2-butene is more stable than cis-2-butene.
4. cis-2-butene has less dipole moment than trans-2-butene.

Question Type : **MCQ**

Question ID : **87827055952**

Option 1 ID : **878270219828**

Option 2 ID : **878270219827**

Option 3 ID : **878270219829**


Option 4 ID : **878270219830**

Status : **Answered**

Chosen Option : **4**

Q.67

The major products formed :



A and B respectively are :

Options

1. and
2. and
3. and
4. and

Question Type : MCQ

Question ID : 87827055956


Option 1 ID : 878270219844

Option 2 ID : 878270219843

Option 3 ID : 878270219845
Option 4 ID : 878270219846
Status : Answered
Chosen Option : 3

Q.68

The ratio $\frac{K_P}{K_C}$ for the reaction :



Options

1. 1
2. $(RT)^{1/2}$
3. $\frac{1}{\sqrt{RT}}$
4. RT

Question Type : MCQ
Question ID : 87827055940
Option 1 ID : 878270219782
Option 2 ID : 878270219780
Option 3 ID : 878270219779
Option 4 ID : 878270219781
Status : Answered
Chosen Option : 3

Q.69

The number of ions from the following that are expected to behave as oxidising agent is :
 Sn^{4+} , Sn^{2+} , Pb^{2+} , Tl^{3+} , Pb^{4+} , Tl^{+}

Options

1. 3
2. 1
3. 2
4. 4

Question Type : MCQ
Question ID : 87827055944
Option 1 ID : 878270219795
Option 2 ID : 878270219798
Option 3 ID : 878270219796
Option 4 ID : 878270219797
Status : Answered
Chosen Option : 3

- Q.70** Evaluate the following statements related to group 14 elements for their correctness.
- (A) Covalent radius decreases down the group from C to Pb in a regular manner.
 - (B) Electronegativity decreases from C to Pb down the group gradually.
 - (C) Maximum covalance of C is 4 whereas other elements can expand their covalance due to presence of d orbitals.
 - (D) Heavier elements do not form $p\pi-p\pi$ bonds.
 - (E) Carbon can exhibit negative oxidation states.

Choose the **correct** answer from the options given below :

Options

1. (A) and (B) Only
2. (A), (B) and (C) Only
3. (C), (D) and (E) Only
4. (C) and (D) Only

Question Type : **MCQ**

Question ID : **87827055945**

Option 1 ID : **878270219799**

Option 2 ID : **878270219801**

Option 3 ID : **878270219802**

Option 4 ID : **878270219800**

Status : **Answered**

Chosen Option : **3**

- Q.71** The incorrect statements regarding enzymes are :

- (A) Enzymes are biocatalysts.
- (B) Enzymes are non-specific and can catalyse different kinds of reactions.
- (C) Most Enzymes are globular proteins.
- (D) Enzyme - oxidase catalyses the hydrolysis of maltose into glucose.

Choose the **correct** answer from the option given below :

Options

1. (A), (B) and (C)
2. (B), (C) and (D)
3. (B) and (C)
4. (B) and (D)

Question Type : **MCQ**

Question ID : **87827055957**

Option 1 ID : **878270219849**

Option 2 ID : **878270219848**

Option 3 ID : **878270219850**

Option 4 ID : **878270219847**

Status : **Answered**

Chosen Option : **1**

Q.72

Match List - I with List - II.

| List - I | | List - II | |
|--------------|----|---------------------------|-------|
| Alkali Metal | | Emission Wavelength in nm | |
| (A) | Li | (I) | 589.2 |
| (B) | Na | (II) | 455.5 |
| (C) | Rb | (III) | 670.8 |
| (D) | Cs | (IV) | 780.0 |

Choose the **correct** answer from the options given below :

Options

1. (A)-(II), (B)-(IV), (C)-(III), (D)-(I)
2. (A)-(I), (B)-(IV), (C)-(III), (D)-(II)
3. (A)-(IV), (B)-(II), (C)-(I), (D)-(III)
4. (A)-(III), (B)-(I), (C)-(IV), (D)-(II)

Question Type : MCQ

Question ID : 87827055943

Option 1 ID : 878270219793

Option 2 ID : 878270219791

Option 3 ID : 878270219794

Option 4 ID : 878270219792

Status : Not Answered

Chosen Option : --

Q.73

The correct statement among the following, for a "chromatography" purification method is :

Options 1.

1. R_f of a polar compound is smaller than that of a non-polar compound.
2. R_f is an integral value.
3. Organic compounds run faster than solvent in the thin layer chromatographic plate.
4. Non-polar compounds are retained at top and polar compounds come down in column chromatography.

Question Type : MCQ

Question ID : 87827055950

Option 1 ID : 878270219822

Option 2 ID : 878270219820

Option 3 ID : 878270219821


Option 4 ID : 878270219819

Status : Answered

Chosen Option : 1

Q.74

Consider the given reaction, identify the major product P.



Options

1. $\text{CH}_3-\text{CH}_2-\overset{\text{O}}{\parallel}{\text{C}}-\text{NH}_2$
2. $\text{CH}_3-\text{CH}_2-\text{CH}_2-\text{OH}$
3. $\text{CH}_3-\overset{\text{O}}{\parallel}{\text{C}}-\text{CH}_2\text{CH}_3$
4. $\text{CH}_3-\underset{\text{OH}}{\text{CH}}-\text{COOH}$

Question Type : MCQ

Question ID : 87827055955

Option 1 ID : 878270219841

Option 2 ID : 878270219842

Option 3 ID : 878270219840

Option 4 ID : 878270219839

Status : Not Answered

Chosen Option : --

Q.75

Molality (m) of 3 M aqueous solution of NaCl is :

(Given : Density of solution = 1.25 g mL^{-1} , Molar mass in g mol^{-1} : Na-23, Cl-35.5)

Options

1. 2.79 m
2. 2.90 m
3. 1.90 m
4. 3.85 m

Question Type : MCQ

Question ID : 87827055938

Option 1 ID : 878270219771

Option 2 ID : 878270219773

Option 3 ID : 878270219774

Option 4 ID : 878270219772

Status : Answered

Chosen Option : 1

Q.76

The correct IUPAC name of $[\text{PtBr}_2(\text{PMe}_3)_2]$ is :

Options

1. dibromobis(trimethylphosphine)platinum(II)
2. bis[bromo(trimethylphosphine)]platinum(II)
3. dibromodi(trimethylphosphine)platinum(II)
4. bis(trimethylphosphine)dibromoplatinum(II)

Question Type : MCQ

Question ID : 87827055947

Option 1 ID : 878270219807

Option 2 ID : 878270219810

Option 3 ID : 878270219809

Option 4 ID : 878270219808

Status : Answered

Chosen Option : 1

Q.77

How can an electrochemical cell be converted into an electrolytic cell ?

Options 1.

1. Applying an external opposite potential greater than E^0_{cell} .
2. Applying an external opposite potential lower than E^0_{cell} .
3. Reversing the flow of ions in salt bridge.
4. Exchanging the electrodes at anode and cathode.

Question Type : MCQ

Question ID : 87827055942

Option 1 ID : 878270219790

Option 2 ID : 878270219789

Option 3 ID : 878270219788

Option 4 ID : 878270219787

Status : Answered

Chosen Option : 1

Q.78

Match List - I with List - II.

List - I
Tetrahedral Complex

List - II
Electronic configuration

- | | |
|----------------------------|--------------------|
| (A) TiCl_4 | (I) e^2, t_2^0 |
| (B) $[\text{FeO}_4]^{2-}$ | (II) e^4, t_2^3 |
| (C) $[\text{FeCl}_4]^-$ | (III) e^0, t_2^0 |
| (D) $[\text{CoCl}_4]^{2-}$ | (IV) e^2, t_2^3 |

Choose the **correct** answer from the options given below :

Options

- (A)-(III), (B)-(IV), (C)-(II), (D)-(I)
- (A)-(I), (B)-(III), (C)-(IV), (D)-(II)
- (A)-(III), (B)-(I), (C)-(IV), (D)-(II)
- (A)-(IV), (B)-(III), (C)-(I), (D)-(II)

Question Type : MCQ

Question ID : 87827055948

Option 1 ID : 878270219812

Option 2 ID : 878270219814

Option 3 ID : 878270219811

Option 4 ID : 878270219813

Status : Answered

Chosen Option : 3

Q.79

Arrange the following elements in the increasing order of number of unpaired electrons in it.

- (A) Sc
(B) Cr
(C) V
(D) Ti
(E) Mn

Choose the **correct** answer from the options given below :

Options

- (C) < (E) < (B) < (A) < (D)
- (B) < (C) < (D) < (E) < (A)
- (A) < (D) < (C) < (E) < (B)
- (A) < (D) < (C) < (B) < (E)

Question Type : MCQ

Question ID : 87827055946

Option 1 ID : 878270219806

Option 2 ID : 878270219803

Option 3 ID : 878270219805


Option 4 ID : 878270219804

Status : Answered





Chosen Option : 4

Q.80

Identify the product (A) in the following reaction.



Options



Question Type : MCQ

Question ID : 87827055954

Option 1 ID : 878270219838

Option 2 ID : 878270219836

Option 3 ID : 878270219837

Option 4 ID : 878270219835

Status : Answered

Chosen Option : 3

- Q.81** Consider the two different first order reactions given below
 $A + B \rightarrow C$ (Reaction 1)
 $P \rightarrow Q$ (Reaction 2)
 The ratio of the half life of Reaction 1 : Reaction 2 is 5 : 2. If t_1 and t_2 represent the time taken to complete $\frac{2}{3}^{\text{rd}}$ and $\frac{4}{5}^{\text{th}}$ of Reaction 1 and Reaction 2, respectively, then the value of the ratio $t_1 : t_2$ is _____ $\times 10^{-1}$ (nearest integer).
 [Given : $\log_{10}(3) = 0.477$ and $\log_{10}(5) = 0.699$]

Given --
 Answer :

Question Type : SA
 Question ID : 87827055962
 Status : Not Answered

- Q.82** Consider the following reactions
 $\text{NiS} + \text{HNO}_3 + \text{HCl} \rightarrow \text{A} + \text{NO} + \text{S} + \text{H}_2\text{O}$
 $\text{A} + \text{NH}_4\text{OH} + \text{H}_3\text{C}-\underset{\text{H}_3\text{C}-\text{C}=\text{N}-\text{OH}}{\text{C}=\text{N}-\text{OH}} \rightarrow \text{B} + \text{NH}_4\text{Cl} + \text{H}_2\text{O}$
 The number of protons that do not involve in hydrogen bonding in the product B is _____.


Given --
 Answer :

Question Type : SA
 Question ID : 87827055964
 Status : Not Answered

- Q.83** Total number of species from the following with central atom utilising sp^2 hybrid orbitals for bonding is _____.
 NH_3 , SO_2 , SiO_2 , BeCl_2 , C_2H_2 , C_2H_4 , BCl_3 , HCHO , C_6H_6 , BF_3 , $\text{C}_2\text{H}_4\text{Cl}_2$

Given --
 Answer :

Question Type : SA
 Question ID : 87827055959
 Status : Not Answered



The ratio of number of oxygen atoms to bromine atoms in the product Q is _____ $\times 10^{-1}$.

Given 15.00
 Answer :

Question Type : SA
 Question ID : 87827055966
 Status : Answered

- Q.85** An amine (X) is prepared by ammonolysis of benzyl chloride. On adding p-toluenesulphonyl chloride to it the solution remains clear. Molar mass of the amine (X) formed is _____ g mol^{-1} .
 (Given molar mass in gmol^{-1} C : 12, H : 1, O : 16, N : 14)

Given 107.00
 Answer :

Question Type : SA
 Question ID : 87827055967
 Status : Answered

Q.86 For the reaction at 298 K, $2A + B \rightarrow C$. $\Delta H = 400 \text{ kJ mol}^{-1}$ and $\Delta S = 0.2 \text{ kJ mol}^{-1} \text{ K}^{-1}$. The reaction will become spontaneous above _____ K.

Given **2000.00**


Answer :

Question Type : SA

Question ID : 87827055960

Status : Answered

Q.87 When ' x ' $\times 10^{-2}$ mL methanol (molar mass = 32 g; density = 0.792 g/cm^3) is added to 100 mL water (density = 1 g/cm^3), the following diagram is obtained.



$x =$ _____ (nearest integer).

[Given : Molal freezing point depression constant of water at 273.15 K is $1.86 \text{ K kg mol}^{-1}$]

Given --


Answer :

Question Type : SA

Question ID : 87827055961

Status : Not Answered

Q.88 Number of carbocations from the following that are **not** stabilized by hyperconjugation is _____.



Given **05.00**

Answer :

Question Type : SA

Question ID : 87827055965

Status : Answered

Q.89 For hydrogen atom, energy of an electron in first excited state is -3.4 eV , K.E. of the same electron of hydrogen atom is $x \text{ eV}$. Value of x is _____ $\times 10^{-1} \text{ eV}$. (Nearest integer)

Given --

Answer :

Question Type : SA

Question ID : 87827055958

Status : Not Answered

Q.90

Among VO_2^+ , MnO_4^- and $\text{Cr}_2\text{O}_7^{2-}$, the spin-only magnetic moment value of the species with least oxidising ability is _____ BM (Nearest integer).
(Given atomic number V = 23, Mn = 25, Cr = 24)

Given **00.00**

Answer :

Question Type : **SA**

Question ID : **87827055963**

Status : **Answered**