

# 12.8.3.11

EE24BTECH11024 - G. Abhimanyu Koushik

**Question:** Using the method of integration find the area bounded by the curve  $|x|+|y| = 1$   
**Solution:**

Theoretical Solution:

The curve  $|x| + |y| = 1$  consists of 4 lines

$$x + y = 1 \quad (1)$$

$$-x + y = 1 \quad (2)$$

$$x - y = 1 \quad (3)$$

$$-x - y = 1 \quad (4)$$

These lines intersect at points  $(1, 0)$ ,  $(-1, 0)$ ,  $(0, 1)$  and  $(0, -1)$  forming a square. To find the area we can integrate  $x + y = 1$  from  $x = 0$  to  $x = 1$  and then multiply the area by 4 to get the total area

$$A_0 = \int_0^1 (1 - x) dx \quad (5)$$

$$A_0 = \left( x - \frac{x^2}{2} \right) \Big|_0^1 \quad (6)$$

$$A_0 = \frac{1}{2} \quad (7)$$

$$A = 4A_0 \quad (8)$$

$$A = 2 \quad (9)$$

Computational Solution:

Taking trapezoid shaped strips of small area and adding them all up. Say we have to find the area of  $y(x)$  from  $x = x_0$  to  $x = x_n$ , discretize points on the  $x$  axis  $x_0, x_1, x_2, \dots, x_n$  such that they are equally spaced with step-size  $h$ .

Sum of all trapezoidal areas is given by,

$$A = \frac{1}{2}h(y(x_1) + y(x_0)) + \frac{1}{2}h(y(x_2) + y(x_1)) + \dots + \frac{1}{2}h(y(x_n) + y(x_{n-1})) \quad (10)$$

$$= h \left[ \frac{1}{2}(y(x_0) + y(x_n)) + y(x_1) + \dots + y(x_{n-1}) \right] \quad (11)$$

Let  $A(x_n)$  be the area enclosed by the curve  $y(x)$  from  $x = x_0$  to  $x = x_n$ ,  $(x_0, x_1, \dots, x_n)$  be equidistant points with step-size  $h$ .

$$A(x_n + h) = A(x_n) + \frac{1}{2}h(y(x_n + h) + y(x_n)) \quad (12)$$

We can repeat this till we get required area.

Discretizing the steps, making  $A(x_n) = A_n$ ,  $y(x_n) = y_n$  we get,

$$A_{n+1} = A_n + \frac{1}{2}h(y_{n+1} + y_n) \quad (13)$$

We can write  $y_{n+1}$  in terms of  $y_n$  using first principle of derivative.  $y_{n+1} = y_n + hy'_n$

$$A_{n+1} = A_n + \frac{1}{2}h((y_n + hy'_n) + y_n) \quad (14)$$

$$A_{n+1} = A_n + \frac{1}{2}h(2y_n + hy'_n) \quad (15)$$

$$A_{n+1} = A_n + hy_n + \frac{1}{2}h^2y'_n \quad (16)$$

$$x_{n+1} = x_n + h \quad (17)$$

In the given question,  $y_n = 1 - x_n$  and  $y'_n = -1$

General Difference Equation will be given by,

$$A_{n+1} = A_n + hy_n + \frac{1}{2}h^2y'_n \quad (18)$$

$$A_{n+1} = A_n + h(1 - x_n) + \frac{1}{2}h^2(-1) \quad (19)$$

$$A_{n+1} = A_n - hx_n + \left(h - \frac{h^2}{2}\right) \quad (20)$$

$$x_{n+1} = x_n + h \quad (21)$$

Iterating till we reach  $x_n = 1$  will return required area. Note, Area obtained is to be multiplied by 4 as the calculated area only accounts for one quarter of the graph.




Fig. 1: Graph of the parabola  $|x| + |y| = 1$  and  $x + y = 1$  and the area of which the integral is calculated