

11.16.3.8.4

EE24BTECH11024 - G. Abhimanyu Koushik

Question:

A three coins are tossed each once, what is the probability of getting atmost 2 heads?

Solution:

Define a discrete random variable X = number of heads

We will take our random variable as a sum of outcomes of three bernoulli random variables

$$X = X_1 + X_2 + X_3 \quad (0.1)$$

Where

$$X_i = \begin{cases} 1, & \text{Outcome in Heads} \\ 0, & \text{Outcome in Tails} \end{cases} \quad (0.2)$$

$$p_{X_i}(n) = \begin{cases} 1 - p, & n = 0 \\ p, & n = 1 \end{cases} \quad (0.3)$$

Where $p = \frac{1}{2}$

Using properties of Z-Transform of PMF

$$M_X(z) = M_{X_1}(z)M_{X_2}(z)M_{X_3}(z) \quad (0.4)$$

$$M_{X_1}(z) = \sum_{n=-\infty}^{\infty} p_{X_1}(n)z^{-n} = (1 - p) + pz^{-1} \quad (0.5)$$

$$M_{X_2}(z) = \sum_{n=-\infty}^{\infty} p_{X_2}(n)z^{-n} = (1 - p) + pz^{-1} \quad (0.6)$$

$$M_{X_3}(z) = \sum_{n=-\infty}^{\infty} p_{X_3}(n)z^{-n} = (1 - p) + pz^{-1} \quad (0.7)$$

$$M_X(z) = ((1 - p) + pz^{-1})^3 \quad (0.8)$$

$$= \sum_{n=-\infty}^{\infty} {}^3C_n(1 - p)^{3-n}p^n z^{-n} \quad (0.9)$$

$$p_X(n) = {}^3C_n p^n (1 - p)^{3-n} \quad (0.10)$$

$$p_X(n) = \frac{{}^3C_n}{8} \quad (0.11)$$

The Probability Mass Function (PMF) for the given random variable is

$$p_X(n) = \begin{cases} \frac{1}{8}, & n = 0 \\ \frac{3}{8}, & n = 1 \\ \frac{3}{8}, & n = 2 \\ \frac{1}{8}, & n = 3 \end{cases} \quad (0.12)$$

The Cumulative Distribution Function (CDF) for the given random variable is

$$F_X(n) = \sum_{k=-\infty}^n {}^3C_k \left(\frac{1}{2}\right)^3 = \begin{cases} 0 & x < 0 \\ {}^3C_0 \left(\frac{1}{2}\right)^3 = \frac{1}{8} & 0 \leq x < 1 \\ {}^3C_1 \left(\frac{1}{2}\right)^3 + {}^3C_0 \left(\frac{1}{2}\right)^3 = \frac{4}{8} & 1 \leq x < 2 \\ {}^3C_2 \left(\frac{1}{2}\right)^3 + {}^3C_1 \left(\frac{1}{2}\right)^3 + {}^3C_0 \left(\frac{1}{2}\right)^3 = \frac{7}{8} & 2 \leq x < 3 \\ {}^3C_3 \left(\frac{1}{2}\right)^3 + {}^3C_2 \left(\frac{1}{2}\right)^3 + {}^3C_1 \left(\frac{1}{2}\right)^3 + {}^3C_0 \left(\frac{1}{2}\right)^3 = 1 & 3 \leq x \end{cases} \quad (0.13)$$

The probability of getting atmost 2 heads is

$$F_X(2) = \frac{7}{8} \quad (0.14)$$

Simulation:

To run a simulation we need to generate random numbers with uniform probability, which is done as shown below:

- 1) Generate a random number by calling rand(). It generates a random number between 0 and RANDMAX
- 2) Divide the generated number by RANDMAX so that it becomes a real number in the range [0, 1)
- 3) If the number is less than p , take it as event happened, else the event did not happen




Fig. 3.1: Probability Mass Function of given Random variable




Fig. 3.2: Cumulative Distribution Function of given Random variable