

10.3.2.5

EE24BTECH11006 - Arnav Mahishi

Question:

Half the perimeter of a rectangular garden, whose length is $4m$ more than its width, is $36m$. Find the dimensions of the garden

Variable	Description
A	Matrix consisting of coefficients in the linear equation
L	Lower triangular matrix
U	Upper triangular matrix
\mathbf{x}	Solution to the linear equation

TABLE 0: Variables Used

Theoretical Solution:

Taking x to be length and y to be width We get:

$$x - y = 4 \quad (0.1)$$

$$x + y = 72 \quad (0.2)$$

$$\Rightarrow 2x = 76 \quad (0.3)$$

$$\Rightarrow x = 38 \quad (0.4)$$

$$\Rightarrow y = 34 \quad (0.5)$$

Computational Solution:

The set of linear equations $x - y - 4 = 0$ and $x + y - 72 = 0$ can be represented by the following equation

$$\begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 4 \\ 72 \end{pmatrix} \quad (0.6)$$

Any non-singular matrix can be represented as a product of a lower triangular matrix L and an upper triangular matrix U :

$$A = L \cdot U \quad (0.7)$$

1. Initialization: - Start by initializing L as the identity matrix $L = I$ and U as a copy of A .

2. Iterative Update: - For each pivot $k = 1, 2, \dots, n$: - Compute the entries of U using the first update equation. - Compute the entries of L using the second update equation.

3. Result: - After completing the iterations, the matrix A is decomposed into $L \cdot U$, where L is a lower triangular matrix with ones on the diagonal, and U is an upper triangular matrix.

1. Update for $U_{k,j}$ (Entries of U)

For each column $j \geq k$, the entries of U in the k -th row are updated as:

$$U_{k,j} = A_{k,j} - \sum_{m=1}^{k-1} L_{k,m} \cdot U_{m,j}, \quad \text{for } j \geq k.$$

This equation computes the elements of the upper triangular matrix U by eliminating the lower triangular portion of the matrix.

2. Update for $L_{i,k}$ (Entries of L)

For each row $i > k$, the entries of L in the k -th column are updated as:

$$L_{i,k} = \frac{1}{U_{k,k}} \left(A_{i,k} - \sum_{m=1}^{k-1} L_{i,m} \cdot U_{m,k} \right), \quad \text{for } i > k.$$

This equation computes the elements of the lower triangular matrix L , where each entry in the column is determined by the values in the rows above it.

Using a code we get L, U as

Step-by-Step Process:

1. Initial Matrix:

$$A = \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix} \quad (0.8)$$

2. Compute U (Upper Triangular Matrix):

Using the update equation for U :

$$U_{11} = A_{11} = 1, \quad U_{12} = A_{12} = -1 \quad (0.9)$$

For U_{22} :

$$U_{22} = A_{22} - L_{21} \cdot U_{12} = 1 - 1 \cdot (-1) = 2 \quad (0.10)$$

3. Compute L (Lower Triangular Matrix):

Using the update equation for L :

$$L_{21} = \frac{A_{21}}{U_{11}} = 1 \quad (0.11)$$

The final L matrix is:

$$L = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} \quad (0.12)$$

4. Solving the System:

Using the equations $Ly = \mathbf{b}$ and $Ux = y$:

• Forward Substitution:

$$\begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} = \begin{pmatrix} 4 \\ 72 \end{pmatrix} \quad (0.13)$$

Solving gives:

$$y_1 = 4, \quad y_2 = 68 \quad (0.14)$$

• **Backward Substitution:**

$$\begin{pmatrix} 1 & -1 \\ 0 & 2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 4 \\ 68 \end{pmatrix} \quad (0.15)$$

Solving gives:

$$x_2 = 34, \quad x_1 = 38 \quad (0.16)$$

Thus, the solution is:

$$\mathbf{x} = \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 38 \\ 34 \end{pmatrix} \quad (0.17)$$




Fig. 0.1: Solution to set of linear equations