

10.3.4.1.3

EE24BTECH11013 - MANIKANTA D

Problem:

Solve the system of linear equations:

$$3x - 5y - 4 = 0 \text{ and } 9x = 2y + 7. \quad (0.1)$$

Step 1: Represent the system in matrix form

The system of equations can be written as:

$$A\mathbf{x} = \mathbf{b}, \quad (0.2)$$

where

$$A = \begin{pmatrix} 3 & -5 \\ 9 & -2 \end{pmatrix}, \mathbf{x} = \begin{pmatrix} x \\ y \end{pmatrix}, \mathbf{b} = \begin{pmatrix} 4 \\ -7 \end{pmatrix}. \quad (0.3)$$

Step 2: Perform LU Decomposition

We decompose the matrix A into the product of a lower triangular matrix L and an upper triangular matrix U , i.e.,

$$A = LU, \quad (0.4)$$

where

$$L = \begin{pmatrix} 1 & 0 \\ l_{21} & 1 \end{pmatrix}, U = \begin{pmatrix} u_{11} & u_{12} \\ 0 & u_{22} \end{pmatrix}. \quad (0.5)$$

Now, let's compute the LU decomposition step by step.

First, we find the elements of U and L :

$$u_{11} = a_{11} = 3, u_{12} = a_{12} = -5. \quad (0.6)$$

Next, we compute l_{21} and u_{22} :

$$l_{21} = \frac{a_{21}}{u_{11}} = \frac{9}{3} = 3, \quad (0.7)$$

$$u_{22} = a_{22} - l_{21}u_{12} = -2 - 3 \times (-5) = 13. \quad (0.8)$$

So the LU decomposition is:

$$L = \begin{pmatrix} 1 & 0 \\ 3 & 1 \end{pmatrix}, U = \begin{pmatrix} 3 & -5 \\ 0 & 13 \end{pmatrix}. \quad (0.9)$$

Step 3: Solve for \mathbf{x} using LU decomposition

Now we solve the system in two steps using forward substitution and backward substitution.

First, solve $L\mathbf{y} = \mathbf{b}$ for \mathbf{y} :

$$\begin{pmatrix} 1 & 0 \\ 3 & 1 \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} = \begin{pmatrix} 4 \\ -7 \end{pmatrix}. \quad (0.10)$$

This gives:

$$y_1 = 4, 3y_1 + y_2 = -7 \Rightarrow 3(4) + y_2 = -7 \Rightarrow y_2 = -19. \quad (0.11)$$

Thus, $\mathbf{y} = \begin{pmatrix} 4 \\ -19 \end{pmatrix}$.

Next, solve $U\mathbf{x} = \mathbf{y}$ for \mathbf{x} :

$$\begin{pmatrix} 3 & -5 \\ 0 & 13 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 4 \\ -19 \end{pmatrix}. \quad (0.12)$$

This gives:

$$13y = -19 \Rightarrow y = -\frac{19}{13}, \quad (0.13)$$

$$3x - 5y = 4 \Rightarrow 3x - 5\left(-\frac{19}{13}\right) = 4 \Rightarrow 3x = 4 - \frac{95}{13} \Rightarrow x = \frac{147}{39}. \quad (0.14)$$

Thus, the solution is $x = \frac{147}{39}$ and $y = -\frac{19}{13}$.

LU Decomposition using Doolittle's algorithm:

The LU decomposition can be efficiently computed using Doolittle's algorithm. This method generates the matrices L (lower triangular) and U (upper triangular) such that $A = LU$. The elements of these matrices are calculated as follows:

Elements of the U Matrix:

For each column j :

$$U_{ij} = A_{ij} \text{ if } i = 0, \quad (0.15)$$

$$U_{ij} = A_{ij} - \sum_{k=0}^{i-1} L_{ik} U_{kj} \text{ if } i > 0. \quad (0.16)$$

Elements of the L Matrix:

For each row i :

$$L_{ij} = \frac{A_{ij}}{U_{jj}} \text{ if } j = 0, \quad (0.17)$$

$$L_{ij} = \frac{A_{ij} - \sum_{k=0}^{j-1} L_{ik} U_{kj}}{U_{jj}} \text{ if } j > 0. \quad (0.18)$$

This systematic approach ensures that the matrix A is decomposed into L and U without requiring row swaps, provided A is nonsingular.




Fig. 0.1: Solving the system of equations