

Probability and Statistics (MA6.101)

Monsoon 2021, IIIT Hyderabad
11 November, Friday (Lecture 21)

Taught by Prof. Pawan Kumar

Probability

Joint Probability

Joint Probability Mass Function

The joint PMF of two discrete random variables X and Y is denoted $P_{XY}(x, y)$ and defined $P(X = x, Y = y)$.

The joint range for X and Y is $R_{XY} = \{(x, y) \mid P_{XY} > 0\}$, i.e., $R_{XY} \subset R_X \times R_Y$.

The sum of joint probabilities must sum to 1.

To compute the probability of some event set $A \subset \mathbb{R}^2$, we take the sum:

$$P((X, Y) \in A) = \sum_{(x_i, y_j) \in A \cap R_{XY}} P_{XY}(x_i, y_j).$$

We can also obtain the PMF of X from its joint PMF with Y as

$$P_X(x) = \sum_{y_j \in R_Y} P_{XY}(x, y_j).$$

We call the PMF obtained in this way the marginal PMF of X .

Joint Cumulative Distribution Function

Analogously, we have

$$F_{XY}(x, y) = P(X \leq x, Y \leq y).$$

This applies to both discrete and continuous variables.

Note that we have the property $F_{XY}(\infty, \infty) = 1$, and $F_{XY}(-\infty, y) = 0$.

For the marginal CDF, we can find

$$F_X(x) = F_{XY}(x, \infty).$$

For the probability of ranges, we can use the following result:

$$P(x_1 < X \leq x_2, y_1 < Y \leq y_2) = F_{XY}(x_2, y_2) - F_{XY}(x_1, y_2) - F_{XY}(x_2, y_1) + F_{XY}(x_1, y_1).$$

Independent Random Variables

If X, Y are two RVs, they are independent if

$$\forall x, y : P_{XY}(x, y) = P_X(x)P_Y(y).$$

Note that if X and Y are independent, then

$$P(x_i|y_j) = P_X(x_i).$$