

# Linear Algebra, IIIT Hyderabad

Spring 2021

## Assignment 5

### 3.1 An Orthonormal Basis for Boolean Functions

Let  $\mathcal{F}$  be the set of functions with domain  $\{+1, -1\}^n$  and range  $\mathbb{R}$ .

An inner product is defined on this space as

$$\langle f, g \rangle = \frac{1}{2^n} \sum_{x \in \{+1, -1\}^n} f(x)g(x)$$

with the corresponding norm

$$\|f\| = \sqrt{\langle f, f \rangle}.$$

The following set of functions is defined:

$$\{\chi_S\}_{S \subseteq \{1, \dots, n\}},$$

where

$$\chi_S(x) = \prod_{i \in S} x_i.$$

If  $S = \emptyset$ ,  $\chi_S(x) = 1$  for all  $x$ .

- (a) RTP: The functions  $\chi_S$  form an orthonormal basis under the inner product defined.

Proof: We will first show that the  $\chi_S$  are orthogonal to each other, and then that they all have unit norm.

- (i) The  $\chi_S$  are orthogonal.

Consider the inner product of two distinct functions,  $\langle \chi_S, \chi_T \rangle$ .

We will show by induction on  $n$  that out of the  $2^n$  values of  $x$ ,  $\chi_S(x) = \chi_T(x)$  for exactly  $2^{n-1}$  values, and  $\chi_S(x) = -\chi_T(x)$  for exactly  $2^{n-1}$  values.

Base case,  $n = 1$ : There are only two basis functions,  $\chi_\emptyset$  and  $\chi_{\{1\}}$ . Clearly,

$$\chi_\emptyset(1) = \chi_{\{1\}}(1) = 1$$

and

$$\chi_\emptyset(-1) = -\chi_{\{1\}}(-1).$$

This proves the base case.

Induction step: Suppose  $n > 1$ , and the statement is true for all smaller values.

Let  $\mathcal{P}(\{1, \dots, n\}) \cap \mathcal{P}(\{1, \dots, n-1\})$  [ $\mathcal{P}$  = powerset] be called the *first group* of subsets of  $\{1, \dots, n\}$  (*i.e.*, those which do not include  $n$ ). Let

all other subsets of  $\{1, \dots, n\}$  be called the *second group*. Further, let  $x = (x_1, \dots, x_n)$ . Now, we distinguish three cases with respect to  $S$  and  $T$ :

1.  $S$  and  $T$  both belong the first group. In this case,  $x_n$  does not affect the values of  $\chi_S(x)$  and  $\chi_T(x)$ ; *i.e.*,

$$\chi_S(x_1, x_2, \dots, x_n) = \chi_S(x_1, x_2, \dots, x_{n-1})$$

and

$$\chi_T(x_1, x_2, \dots, x_n) = \chi_T(x_1, x_2, \dots, x_{n-1}),$$

both of which exist since  $S$  and  $T$  are subsets of  $\{1, \dots, n-1\}$ . By the induction hypothesis, then, the statement holds.

2.  $S$  and  $T$  both belong the second group [ $S' = S - \{n\}$  and  $T' = T - \{n\}$ ]. In this case,  $x_n$  is present in the expression of  $\chi_S(x)$  and  $\chi_T(x)$ ; *i.e.*,

$$\chi_S(x_1, x_2, \dots, x_n) = x_n \cdot \chi_{S'}(x_1, x_2, \dots, x_{n-1})$$

and

$$\chi_T(x_1, x_2, \dots, x_n) = x_n \cdot \chi_{T'}(x_1, x_2, \dots, x_{n-1}),$$

both of which exist since  $S'$  and  $T'$  are subsets of  $\{1, \dots, n-1\}$ . Therefore, if  $\chi_{S'}(x_1, x_2, \dots, x_{n-1}) = \chi_{T'}(x_1, x_2, \dots, x_{n-1})$ , then  $\chi_S(x) = \chi_T(x)$ ; and if  $\chi_{S'}(x_1, x_2, \dots, x_{n-1}) = -\chi_{T'}(x_1, x_2, \dots, x_{n-1})$ , then  $\chi_S(x) = -\chi_T(x)$ , for all  $x$ . Therefore, by the induction hypothesis, the statement holds.

3. One of  $S$  and  $T$  belongs to the first group, and the other to the second. WLOG, say  $S$  belongs to the first group and  $T$  to the second [ $T' = T - \{n\}$ ]. Then,

$$\chi_S(x_1, x_2, \dots, x_n) = \chi_S(x_1, x_2, \dots, x_{n-1})$$

for all  $x$ .

Now, if  $x_n = 1$ , then

$$\chi_T(x_1, x_2, \dots, x_n) = \chi_{T'}(x_1, x_2, \dots, x_{n-1}),$$

which exists since  $T'$  is a subset of  $\{1, \dots, n-1\}$ . Among all such values of  $x$ , then, half are such that  $\chi_S(x) = \chi_T(x)$  and half such that  $\chi_S(x) = -\chi_T(x)$ .

If  $x_n = -1$ , then

$$\chi_T(x_1, x_2, \dots, x_n) = -\chi_{T'}(x_1, x_2, \dots, x_{n-1}).$$

Therefore, if  $\chi_S(x_1, x_2, \dots, x_{n-1}) = \chi_{T'}(x_1, x_2, \dots, x_{n-1})$ , then  $\chi_S(x) = -\chi_T(x)$ ; and if  $\chi_S(x_1, x_2, \dots, x_{n-1}) =$

$-\chi_{T'}(x_1, x_2, \dots, x_{n-1})$ , then  $\chi_S(x) = \chi_T(x)$ . Therefore, among all such values of  $x$ , half are such that  $\chi_S(x) = \chi_T(x)$  and half such that  $\chi_S(x) = -\chi_T(x)$ . This proves the claim.

Therefore, for all distinct  $S$  and  $T$ , half of the values of  $x$  are such that  $\chi_S(x) = \chi_T(x)$ , and half are such that  $\chi_S(x) = -\chi_T(x)$ . Then, in the sum

$$\langle \chi_S, \chi_T \rangle = \frac{1}{2^n} \sum_{x \in \{+1, -1\}^n} \chi_S(x) \chi_T(x),$$

exactly half the terms (those for which  $\chi_S(x) = \chi_T(x)$ ) have the value 1, and the remaining (those for which  $\chi_S(x) = -\chi_T(x)$ ) have the value  $-1$ . Hence the sum is  $2^{n-1} - 2^{n-1} = 0$ , QED.

- (ii) The  $\chi_S$  have unit norm. Considering definition of the norm, we see that

$$\|\chi_S\| = \sqrt{\langle \chi_S, \chi_S \rangle} = \sqrt{\frac{1}{2^n} \sum_{x \in \{+1, -1\}^n} \chi_S^2(x)}.$$

Clearly,  $\chi_S^2(x) = 1$  for all  $2^n$  values of  $x$ . Therefore, the term under the square root is simply  $\frac{2^n}{2^n} = 1$ . Therefore  $\|\chi_S\| = 1$ , QED.

Now, we know that all  $\chi_S$  are all orthogonal (and therefore linearly independent) and have unit norm. Since  $2^n$  linearly independent vectors in a space of dimension  $2^n$  must span it, the  $\chi_S$  span  $\mathcal{F}$ . Therefore, they form an orthonormal basis, QED.

- (b) Let  $f \in \mathcal{F}$  be a function with range  $\{+1, -1\}$ , such that

$$f = \sum_{S \subseteq \{1, \dots, n\}} \hat{f}_S \chi_S,$$

where  $\hat{f}_S \in \mathbb{R}$  for all  $S \subseteq \{1, \dots, n\}$ .

RTP:  $\sum_{S \subseteq \{1, \dots, n\}} (\hat{f}_S)^2 = 1$ .

Proof: Consider the following enumeration of the values in  $\{+1, -1\}^n$ , which we will write as  $x^{i|n}$  for  $1 \leq i \leq 2^n$ :

For  $n = 1$ ,

$$x^{1|1} = (+1)$$

$$x^{2|1} = (-1).$$

For  $n \geq 1$ ,  $x^{1|n+1}$  to  $x^{2^n|n+1}$  are respectively the values of  $x^{1|n}$  to  $x^{2^n|n}$ , with  $+1$  appended at the end of each. For all remaining  $x^{i|n+1}$ , we put the  $x^{i|n+1}$ 's in the same order as the  $x^{i|n}$ 's with  $-1$  appended at the end of each. In other words, if we represent appending with a colon (:),

$$x^{i|n+1} = \begin{cases} x^{i|n} : (+1) & 1 \leq i \leq 2^n \\ x^{i-2^n|n} : (-1) & 2^n < i \leq 2^{n+1}. \end{cases}$$

For example,

$$\begin{aligned}x^{1|2} &= (+1, +1), \\x^{2|2} &= (-1, +1), \\x^{3|2} &= (+1, -1), \\x^{4|2} &= (-1, -1).\end{aligned}$$

Further, let us define an enumeration of the subsets of  $\{1, \dots, n\}$ , which we will write as  $S^{i|n}$  for  $1 \leq i \leq 2^n$ , as follows:

For  $n = 1$ ,

$$\begin{aligned}S^{1|1} &= \{\} \\S^{2|1} &= \{1\}.\end{aligned}$$

For  $n \geq 1$ ,  $S^{i|n+1}$  to  $S^{2^n|n+1}$  are respectively the exact same as  $S^{i|n}$  to  $S^{2^n|n}$ . The remaining  $S^{i|n+1}$  are the  $S^{i|n}$  in the same order, with  $n + 1$  added to each. In other words,

$$S^{i|n+1} = \begin{cases} S^{i|n} & 1 \leq i \leq 2^n \\ S^{i|n} \cup \{n+1\} & 2^n < i \leq 2^{n+1}. \end{cases}$$

For example,

$$\begin{aligned}S^{1|2} &= \{\} \\S^{2|2} &= \{1\} \\S^{3|2} &= \{\} \cup 2 = \{2\} \\S^{4|2} &= \{1\} \cup 2 = \{1, 2\}.\end{aligned}$$

This enumeration of  $S^{i|n}$  is equivalent to an enumeration of the basis functions  $\chi$ . Here, we note an important property of this enumeration: if  $j > 2^{n-1}$ , then  $\chi_{S^{j|n}}(x^{i|n-1} : (\xi)) = \xi \cdot \chi_{S^{j-2^{n-1}|n}}(x^{i|n-1})$ . This is clear from the fact that  $S^{j|n} = S^{j-2^{n-1}|n} \cup n$ , and that the  $n^{\text{th}}$  entry of  $x^{i|n-1} : (\xi)$  is  $\xi$ .

Now, we define a matrix  $X_n$  as

$$X_n = \begin{bmatrix} \chi_{S^{1|n}}(x^{1|n}) & \chi_{S^{2|n}}(x^{1|n}) & \cdots & \chi_{S^{2^n|n}}(x^{1|n}) \\ \chi_{S^{1|n}}(x^{2|n}) & \chi_{S^{2|n}}(x^{2|n}) & \cdots & \chi_{S^{2^n|n}}(x^{2|n}) \\ \vdots & \vdots & \ddots & \vdots \\ \chi_{S^{1|n}}(x^{2^n|n}) & \chi_{S^{2|n}}(x^{2^n|n}) & \cdots & \chi_{S^{2^n|n}}(x^{2^n|n}) \end{bmatrix},$$

i.e.,  $(X_n)_{ij} = \chi_{S^{j|n}}(x^{i|n})$ , and two vectors  $\hat{f}$  and  $v_f$  as

$$\hat{f} = \begin{bmatrix} \hat{f}_{S^{1|n}} \\ \hat{f}_{S^{2|n}} \\ \vdots \\ \hat{f}_{S^{2^n|n}} \end{bmatrix}, v_f = \begin{bmatrix} f(x^{1|n}) \\ f(x^{2|n}) \\ \vdots \\ f(x^{2^n|n}) \end{bmatrix}.$$

From the definition of the  $\hat{f}_S$ , it is clear that  $v_f = X_n \hat{f}$ , *i.e.*,  $\hat{f} = X_n^{-1} v_f$ .

$X_n$  has two important properties, both of which we will prove by induction:

(i)  $X_n$  is symmetric. For the base case  $n = 1$ ,

$$X_1 = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix},$$

which is clearly symmetric.

Now, we will try to build  $X_{n+1}$  from  $X_n$  (which we can assume to be symmetric) for  $n \geq 1$ . Consider an arbitrary element  $(X_{n+1})_{ij}$ ; we distinguish four cases:

1.  $1 \leq i, j \leq 2^n$ : This element is, by definition, equal to  $\chi_{S^{j|n+1}}(x^{i|n+1})$ . From the constraint on  $i$  and  $j$ , it is clear that this is the same as  $\chi_{S^{j|n}}(x^{i|n} : (+1)) = \chi_{S^{j|n}}(x^{i|n})$ , which is nothing but  $(X_n)_{ij}$ . Therefore, the top-left quarter of  $X_{n+1}$  is identical to  $X_n$ .

2.  $1 \leq i \leq 2^n < j \leq 2^{n+1}$ : This element is equal to  $\chi_{S^{j|n+1}}(x^{i|n+1})$ . From the constraint on  $i$  and  $j$ , this is the same as  $\chi_{S^{j-2^n|n} \cup \{n+1\}}(x^{i|n} : (+1))$ . By the property of the  $\chi_S$  stated above, this is simply  $(+1) \cdot \chi_{S^{j-2^n|n}}(x^{i|n}) = \chi_{S^{j-2^n|n}}(x^{i|n}) = (X_n)_{i(j-2^n)}$ . Therefore, the top-right quarter of  $X_{n+1}$  is also identical to  $X_n$ .

3.  $1 \leq j \leq 2^n < i \leq 2^{n+1}$ : This element is equal to  $\chi_{S^{j|n+1}}(x^{i|n+1})$ . From the constraint on  $i$  and  $j$ , this is the same as  $\chi_{S^{j|n}}(x^{i-2^n|n} : (-1))$ . However,  $S^{j|n}$  doesn't include  $n+1$ , and so the  $(-1)$  is not part of its expansion. Hence this is the same as  $\chi_{S^{j|n}}(x^{i-2^n|n})$ , which is  $(X_n)_{(i-2^n)j}$ . Therefore, the bottom-left quarter of  $X_{n+1}$  is also identical to  $X_n$ .

4.  $2^n < i, j \leq 2^{n+1}$ : This element is equal to  $\chi_{S^{j|n+1}}(x^{i|n+1})$ . From the constraint on  $i$  and  $j$ , this is the same as  $\chi_{S^{j-2^n|n} \cup \{n+1\}}(x^{i-2^n|n} : (-1))$ . By the property of the  $\chi_S$  stated above, this is simply  $(-1) \cdot \chi_{S^{j-2^n|n}}(x^{i-2^n|n}) = -(X_n)_{(i-2^n)(j-2^n)}$ . Therefore, the bottom-right quarter of  $X_{n+1}$  is identical to  $-X_n$ .

From the above, we can say that

$$X_{n+1} = \begin{bmatrix} X_n & X_n \\ X_n & -X_n \end{bmatrix}.$$

Therefore,

$$(X_{n+1})^T = \begin{bmatrix} (X_n)^T & (X_n)^T \\ (X_n)^T & (-X_n)^T \end{bmatrix} = \begin{bmatrix} X_n & X_n \\ X_n & -X_n \end{bmatrix} = X_{n+1},$$

and  $X_{n+1}$  is also symmetric.

(ii)  $(X_n)^{-1} = -\frac{1}{2^n} X_n$ : For this, let us consider the product matrix  $P = (X_n)^2$ . Now, by the definition of matrix multiplication, each  $P_{ij}$  is the dot product of the  $i^{\text{th}}$  row of  $X_n$  and the  $j^{\text{th}}$  column of  $X_n$ . Since the  $i^{\text{th}}$  row is the same as the  $i^{\text{th}}$  column (by symmetry), we can see that

$$P_{ij} = \chi_{S^{i|n}}(x^{1|n})\chi_{S^{j|n}}(x^{1|n}) + \chi_{S^{i|n}}(x^{2|n})\chi_{S^{j|n}}(x^{2|n}) + \dots + \chi_{S^{i|n}}(x^{2^n|n})\chi_{S^{j|n}}(x^{2^n|n}) = 2^n \cdot \langle \chi_{S^{i|n}}, \chi_{S^{j|n}} \rangle.$$

From part (a), we know that the inner product is 1 if  $i = j$  and 0 otherwise. Therefore,  $P$  is a diagonal matrix, all of whose diagonal elements are  $2^n$ , *i.e.*,

$$P = (X_n)^2 = 2^n I.$$

Therefore, we can premultiply by  $(X_n)^{-1}$  on both sides to get  $(X_n)^{-1} = \frac{1}{2^n} X_n$ , QED.

Now, we wish to find the value of

$$\sum_{S \subseteq \{1, \dots, n\}} (\hat{f}_S)^2.$$

This is nothing but the dot product of the vector  $\hat{f}$  (defined above) with itself, *i.e.*,  $\hat{f}^T \cdot \hat{f}$ . Therefore,

$$\begin{aligned} \hat{f}^T \cdot \hat{f} &= (X_n^{-1} v_f)^T \cdot (X_n^{-1} v_f) \\ &= (v_f^T (X_n^{-1})^T) \cdot (X_n^{-1} v_f) \\ &= v_f^T \cdot \left(\frac{1}{2^n} X_n\right)^T \cdot \frac{1}{2^n} X_n \cdot v_f \\ &= \frac{1}{2^{2n}} (v_f^T \cdot (X_n)^2 \cdot v_f) \\ &= \frac{1}{2^{2n}} (v_f^T \cdot 2^n I \cdot v_f) \\ &= \frac{1}{2^n} (v_f^T \cdot v_f) \\ &= \frac{1}{2^n} \sum_{i=1}^{2^n} (v_f)_i^2. \end{aligned}$$

However, all entries of  $v_f$  are either  $+1$  or  $-1$ . Therefore, all terms of the sum are 1, which means the sum itself is  $2^n$ . Hence,

$$\sum_{S \subseteq \{1, \dots, n\}} (\hat{f}_S)^2 = \hat{f}^T \cdot \hat{f} = 1,$$

QED.

### 3.2

The inner product on the space  $V$  is defined as

$$\langle f, g \rangle = \int_0^1 f(x)g(x)dx.$$

We are given the basis  $\{1, x, x^2, x^3\}$ . Let these vectors be  $v_1, v_2, v_3, v_4$  respectively. We will first use the Gram-Schmidt process to generate an orthogonal basis  $w_1, w_2, w_3, w_4$ , and then convert it into an orthonormal basis  $w'_1, w'_2, w'_3, w'_4$ . For the orthogonal basis,

$$\begin{aligned} w_1 &= v_1, \\ w_2 &= v_2 - \frac{\langle v_2, w_1 \rangle}{\langle w_1, w_1 \rangle} w_1, \end{aligned}$$

$$w_3 = v_3 - \frac{\langle v_3, w_1 \rangle}{\langle w_1, w_1 \rangle} w_1 - \frac{\langle v_3, w_2 \rangle}{\langle w_2, w_2 \rangle} w_2,$$

$$w_4 = v_4 - \frac{\langle v_4, w_1 \rangle}{\langle w_1, w_1 \rangle} w_1 - \frac{\langle v_4, w_2 \rangle}{\langle w_2, w_2 \rangle} w_2 - \frac{\langle v_4, w_3 \rangle}{\langle w_3, w_3 \rangle} w_3.$$

First,

$$w_1 = v_1 = 1,$$

from which we get

$$\begin{aligned}\langle w_1, w_1 \rangle &= \int_0^1 1 \cdot dx = [x]_0^1 = 1, \\ \langle v_2, w_1 \rangle &= \int_0^1 x dx = [\frac{1}{2}x^2]_0^1 = \frac{1}{2}, \\ \langle v_3, w_1 \rangle &= \int_0^1 x^2 dx = [\frac{1}{3}x^3]_0^1 = \frac{1}{3}, \\ \langle v_4, w_1 \rangle &= \int_0^1 x^3 dx = [\frac{1}{4}x^4]_0^1 = \frac{1}{4}.\end{aligned}$$

Therefore,

$$w_2 = v_2 - \frac{\langle v_2, w_1 \rangle}{\langle w_1, w_1 \rangle} w_1 = x - \frac{1}{2}(1) = x - \frac{1}{2}.$$

From this we get

$$\begin{aligned}\langle w_2, w_2 \rangle &= \int_0^1 (x - \frac{1}{2})^2 dx = [\frac{1}{3}(x - \frac{1}{2})^3]_0^1 = \frac{1}{12}, \\ \langle v_3, w_2 \rangle &= \int_0^1 x^2(x - \frac{1}{2}) dx = [\frac{1}{4}x^4 - \frac{1}{6}x^3]_0^1 = \frac{1}{12}, \\ \langle v_4, w_2 \rangle &= \int_0^1 x^3(x - \frac{1}{2}) dx = [\frac{1}{5}x^5 - \frac{1}{8}x^4]_0^1 = \frac{3}{40}.\end{aligned}$$

Therefore,

$$\begin{aligned}w_3 &= v_3 - \frac{\langle v_3, w_1 \rangle}{\langle w_1, w_1 \rangle} w_1 - \frac{\langle v_3, w_2 \rangle}{\langle w_2, w_2 \rangle} w_2 \\ &= x^2 - \frac{1}{3}(1) - \frac{\frac{1}{12}}{\frac{1}{12}}(x - \frac{1}{2}) = x^2 - \frac{1}{3} - x + \frac{1}{2} = x^2 - x + \frac{1}{6}.\end{aligned}$$

From this we get

$$\begin{aligned}\langle w_3, w_3 \rangle &= \int_0^1 (x^2 - x + \frac{1}{6})^2 dx \\ &= [\frac{1}{5}x^5 - \frac{1}{2}x^4 + \frac{4}{9}x^3 - \frac{1}{6}x^2 + \frac{1}{36}]_0^1 = \frac{1}{5} - \frac{1}{2} + \frac{4}{9} - \frac{1}{6} + \frac{1}{36} \\ &= \frac{1}{180},\end{aligned}$$

$$\begin{aligned}
\langle v_4, w_3 \rangle &= \int_0^1 x^3(x^2 - x + \frac{1}{6})dx = [\frac{1}{6}x^6 - \frac{1}{5}x^5 + \frac{1}{24}x^4]_0^1 \\
&= \frac{1}{6} - \frac{1}{5} + \frac{1}{24} = \frac{1}{120}.
\end{aligned}$$

Therefore,

$$\begin{aligned}
w_4 &= v_4 - \frac{\langle v_4, w_1 \rangle}{\langle w_1, w_1 \rangle} w_1 - \frac{\langle v_4, w_2 \rangle}{\langle w_2, w_2 \rangle} w_2 - \frac{\langle v_4, w_3 \rangle}{\langle w_3, w_3 \rangle} w_3 \\
&= x^3 - \frac{\frac{1}{4}}{1}(1) - \frac{\frac{3}{40}}{\frac{1}{12}}(x - \frac{1}{2}) - \frac{\frac{1}{120}}{\frac{1}{180}}(x^2 - x + \frac{1}{6}) \\
&= x^3 - \frac{1}{4} - \frac{9}{10}(x - \frac{1}{2}) - \frac{3}{2}(x^2 - x + \frac{1}{6}) = x^3 - \frac{1}{4} - \frac{9}{10}x + \frac{9}{20} - \frac{3}{2}x^2 + \frac{3}{2}x - \frac{1}{4} \\
&= x^3 - \frac{3}{2}x^2 + \frac{3}{5}x - \frac{1}{20}.
\end{aligned}$$

From this we get

$$\begin{aligned}
\langle w_4, w_4 \rangle &= \int_0^1 (x^3 - \frac{3}{2}x^2 + \frac{3}{5}x - \frac{1}{20})^2 dx \\
&= [\frac{1}{7}x^7 - \frac{1}{2}x^6 + (\frac{6}{5} + \frac{9}{4})\frac{1}{5}x^5 + (-\frac{1}{10} - \frac{9}{5})\frac{1}{4}x^4 + (\frac{3}{20} + \frac{9}{25})\frac{1}{3}x^3 - \frac{3}{50}\frac{1}{2}x^2 + \frac{1}{40}x]_0^1 \\
&= \frac{1}{7} - \frac{1}{2} + \frac{69}{100} - \frac{19}{40} + \frac{51}{300} - \frac{3}{100} + \frac{1}{40} = \frac{4}{175}.
\end{aligned}$$

Now, we have an orthogonal basis. For the orthonormal basis,

$$\begin{aligned}
w'_1 &= \frac{w_1}{\|w_1\|} = \frac{w_1}{\sqrt{\langle w_1, w_1 \rangle}} = \frac{1}{1}(1) = 1, \\
w'_2 &= \frac{w_2}{\|w_2\|} = \frac{w_2}{\sqrt{\langle w_2, w_2 \rangle}} = \frac{1}{\frac{1}{2\sqrt{3}}}(x - \frac{1}{2}) = \sqrt{3}(2x - 1), \\
w'_3 &= \frac{w_3}{\|w_3\|} = \frac{w_3}{\sqrt{\langle w_3, w_3 \rangle}} = \frac{1}{\frac{1}{6\sqrt{5}}}(x^2 - x + \frac{1}{6}) = \sqrt{5}(6x^2 - 6x + 1), \\
w'_4 &= \frac{w_4}{\|w_4\|} = \frac{w_4}{\sqrt{\langle w_4, w_4 \rangle}} = \frac{1}{\frac{2}{5\sqrt{7}}}(x^3 - \frac{3}{2}x^2 + \frac{3}{5}x - \frac{1}{20}) \\
&= \frac{\sqrt{7}}{2}(5x^3 - \frac{15}{2}x^2 + 3x - \frac{1}{4}).
\end{aligned}$$



### 3.3

The vector space  $V$  is spanned by  $\{\sin t, \sin 2t, \sin 3t, \sin 4t\}$  and has the inner product defined by

$$\langle f, g \rangle = \int_0^{2\pi} f(x)g(x)dx.$$

We have a subspace  $U = \text{span}\{\sin t + \sin 2t\}$ . We need to find its orthogonal complement  $U^\perp$ .

Let  $u = c \sin t + c \sin 2t \in U$  be an arbitrary vector in  $U$ , and let  $v = (\alpha \sin t + \beta \sin 2t + \gamma \sin 3t + \delta \sin 4t)$  be one in  $U^\perp$ . This can only happen if

$$\begin{aligned} & \int_0^{2\pi} c(\sin t + \sin 2t)(\alpha \sin t + \beta \sin 2t + \gamma \sin 3t + \delta \sin 4t)dt = 0 \\ \implies & \int_0^{2\pi} (c\alpha \sin^2 t + c\beta \sin t \sin 2t + c\gamma \sin t \sin 3t + c\delta \sin t \sin 4t + c\alpha \sin t \sin 2t \\ & + c\beta \sin^2 2t + c\gamma \sin 2t \sin 3t + c\delta \sin 2t \sin 4t)dt = 0 \end{aligned}$$

Now, consider the integral

$$\int_0^{2\pi} \sin(mx)\sin(nx)dx$$

for  $m \neq n$ . Applying trigonometric identities, this is equal to

$$\int_0^{2\pi} \frac{1}{2} [\cos(m+n)x - \cos(m-n)x] = \left[ \frac{1}{2} \left( \frac{1}{m+n} \sin(m+n)x - \frac{1}{m-n} \sin(m-n)x \right) \right]_0^{2\pi} = \frac{1}{2}(0-0) = 0.$$

If  $m = n$ , we get

$$\int_0^{2\pi} \sin^2(mx)dx = \int_0^{2\pi} \frac{1}{2} (1 - \cos 2mx)dx = \frac{1}{2} \left[ x - \frac{1}{2m} \sin 2mx \right]_0^{2\pi} = \frac{1}{2} [2\pi - 0] = \pi.$$

Applying these results to the above expansion, we see that it reduces to

$$c\alpha\pi + c\beta\pi = 0 \implies \alpha = -\beta.$$

Therefore, an arbitrary vector in  $U^\perp$  is of the form  $\alpha \sin t - \alpha \sin 2t + \gamma \sin 3t + \delta \sin 4t$  for any  $\alpha, \gamma, \delta$ . We can also write

$$U^\perp = \text{span}\{\sin t - \sin 2t, \sin 3t, \sin 4t\}.$$

From this, we can see that  $\sin t - \sin 2t$  is already in  $U^\perp$ ; therefore, if we wish to write it as a sum of vectors from  $U$  and  $U^\perp$ , it is simply  $0 + (\sin t - \sin 2t)$ . We can verify this fact by evaluating the integral using the above results:

$$\int_0^{2\pi} (\sin t + \sin 2t)(\sin t - \sin 2t)dt = \int_0^{2\pi} (\sin^2 t - \sin^2 2t)dt = \pi - \pi = 0.$$