

9/10/24

LAB - 1

classmate

Date _____

Page _____

- Q) Develop a java program that prints all real solutions to the quadratic equation $ax^2 + bx + c = 0$. Read a, b, c and use the quadratic formula. If the discriminate $b^2 - 4ac$ is negative, display a message stating that there are no real solutions.

```
import java.util.*;
class Quad {
    Scanner sc = new Scanner(System.in);
    int a, b, c, d;
    double r1, r2, d-sq;

    void input() {
        System.out.println("Enter coefficients a, b, c=");
        a = sc.nextInt();
        b = sc.nextInt();
        c = sc.nextInt();
    }
}
```

```
void calc() {
    int d = b*b - 4*a*c;
    System.out.println(d);
    if (d == 0) {
        r1 = -b/(2*a);
        System.out.println("Roots are real and equal");
        System.out.println("Root 1 = " + r1 + " \n Root 2 = " + r1);
    }
}
```

```

else if (d > 0) {
    d-sq = Math.sqrt(d);
    r1 = (-b + d-sq) / (2.0 * a);
    r2 = (-b - d-sq) / (2.0 * a);    real and distinct
    System.out.println("Roots are imaginary");
    System.out.println("Root 1 = " + r1 + " \n Root 2 = " + r2);
}

```

```

else {
    d-sq = Math.sqrt(-d);
    r1 = -b / (2.0 * a);
    r2 = d-sq / (2.0 * a);
    System.out.println("Roots are imaginary");
    System.out.println("Root 1 = " + r1 + " + " + r2 + "i" +
        "\n Root 2 = " + r1 + " - " + r2 + "i");
} } }

```

```

class Quadratic {
    public static void main (String [] args) {
        Quad quad = new Quad();
        quad.input();
        quad.calc();
    }
}

```

Output

enter coefficients a, b, c

1

5

6

Roots are real and distinct

Root 1 = 2.0

Root 2 = 2.0

enter coefficients a, b, c

1

1

1

Roots are imaginary

$$\text{Root 1} = -0.5 + 0.8660254037844386i$$

$$\text{Root 2} = -0.5 + 0.8660254037844386i$$

Enter coefficient a, b, c

4

12

9

Roots are real and equal

$$\text{Root 1} = -1.5$$

$$\text{Root 2} = -1.5$$

o/p seen

gk
9/10/24