

BSDS 100: Intro to Data Science with R

Assignment 4

by James D. Wilson (University of San Francisco)

Directions: For all questions in this assignment, write complete sentences and fully answer any question that is asked. Provide all R code and solutions by *knitting* your final RStudio file into a single file named [your_name]_CA4.pdf. This assignment is due next Tuesday at the beginning of class. Late assignments will automatically have 10 points deducted.

1. Create the vector

```
> myAtomicVector <- c(1, 4, 3, 2, NA, 3.22, -44, 2, NA, 0, 22, 34)
```

- (a) How many positive numbers (> 0) are there in this vector?
 - (b) How many negative numbers (< 0) are there in this vector?
 - (c) How many 0's are there in this vector?
 - (d) How many NAs are there in this vector?
 - (e) How many numbers in the vector are non-zero **and** not NAs?
 - (f) What is the sum of the positive numbers in this vector?
 - (g) What is the sum of the negative numbers in this vector?
2. Consider a vector of length 1000, where F_n is the n th number in the sequence. Then the [Fibonacci sequence](#) is the vector where the following recursion holds:

$$F_n = F_{n-1} + F_{n-2}$$

That is, the n th number in the sequence will be the sum of the previous two numbers.

- (a) Using the `rep()` and `seq()` commands, create a vector (but don't print it out!) that contains the first 1000 Fibonacci numbers starting from the first two numbers (1, 1).
- (b) Using the Fibonacci numbers generated above, generate a vector (of length 999) with values (again, don't print these out)

$$Z_n = \frac{F_{n+1}}{F_n}$$

- (c) Plot the vector Z_n using the command `plot(Z_n)`. Then add a line to the plot using the following command `abline(h = (1 + sqrt(5))/2)`. This value is known as the [golden ratio](#) in mathematics.
- (d) Comment on the plot that you obtain. What do you observe?