



AIR QUALITY ANALYSIS

Comprehensive Data Analysis & Future Projections

A detailed report on air quality metrics,
pollutant distribution, and environmental trends

September 2026

Table of Contents

1. Executive Summary
2. Understanding Air Quality Metrics
3. AQI Categories Explained
4. Data Analysis & Current Situation
5. Key Findings
6. Pollutant Analysis
7. Geographic Breakdown
8. Trend Analysis
9. Future Projections
10. Recommendations

1. Executive Summary

This comprehensive report analyzes air quality data across multiple regions and time periods, examining pollutant concentrations, AQI (Air Quality Index) trends, and making evidence-based projections for future air quality conditions.

Key Highlights:

✦ Critical Findings

- Average AQI Status:** The data shows varying air quality levels across different geographic regions and time periods, with some areas experiencing moderate to poor air quality conditions.
- Primary Pollutants:** Multiple harmful pollutants are identified as primary contributors to air quality degradation, requiring targeted mitigation strategies.
- Temporal Patterns:** AQI values show significant fluctuations throughout the month, indicating changing atmospheric and meteorological conditions.
- Geographic Variation:** Different states and areas demonstrate different pollution levels, suggesting localized sources or regional weather patterns.
- Health Implications:** Current air quality levels in some regions may impact public health, particularly for vulnerable populations.

2. Understanding Air Quality Metrics

Air quality is measured using several important metrics and indices. This section explains the key terms and measurements used throughout this report.

AQI - Air Quality Index

Definition: The Air Quality Index (AQI) is a scale used to communicate how polluted the air currently is or how polluted it is forecast to become. It is calculated based on concentrations of major air pollutants. A higher AQI value indicates worse air quality.

Scale Range: 0-500+ (values over 500 indicate hazardous conditions)

Unit: Numeric index value (unitless)

Primary Pollutants Measured

PM2.5 - Fine Particulate Matter

Particles with diameter ≤ 2.5 micrometers. These microscopic particles can penetrate deep into lungs and bloodstream, causing serious respiratory and cardiovascular health problems. Common sources: vehicle emissions, industrial pollution, burning of fossil fuels, dust.

Measurement Unit: $\mu\text{g}/\text{m}^3$ (micrograms per cubic meter)

PM10 - Coarse Particulate Matter

Particles with diameter ≤ 10 micrometers. Larger than PM2.5, can lodge in airways and cause respiratory issues. Sources: dust storms, construction, grinding, vehicular emissions.

Measurement Unit: $\mu\text{g}/\text{m}^3$ (micrograms per cubic meter)

NO2 - Nitrogen Dioxide

A reddish-brown toxic gas formed from nitrogen oxide emissions. Major source: traffic emissions from vehicles. Can cause bronchitis and reduced lung function, especially in children and elderly.

Measurement Unit: ppb (parts per billion) or $\mu\text{g}/\text{m}^3$

SO2 - Sulfur Dioxide

A colorless gas with pungent odor. Released when burning fossil fuels containing sulfur. Aggravates asthma and respiratory conditions. Sources: power plants, refineries, industrial facilities.

Measurement Unit: ppb (parts per billion) or $\mu\text{g}/\text{m}^3$

O3 - Ozone (at ground level)

Ground-level ozone is formed when pollutants react with sunlight. Reduces lung function and causes respiratory disease. Not to be confused with protective ozone in upper atmosphere.

Measurement Unit: ppb (parts per billion)

🔥 CO - Carbon Monoxide

Colorless, odorless, toxic gas. Reduces oxygen-carrying capacity of blood. Primary source: vehicle emissions and incomplete combustion. Particularly dangerous in enclosed spaces.

Measurement Unit: ppm (parts per million) or mg/m³

3. AQI Categories Explained

Air Quality Index is categorized into six levels. Each level indicates different health impacts and recommended actions for different population groups.

AQI Range	Category	Health Effects	Recommended Action
0-50	GOOD	No health impacts expected. Air quality is satisfactory.	Enjoy outdoor activities freely. No precautions needed.
51-100	MODERATE	Air quality is acceptable for most people. Unusually sensitive people may experience issues.	Unusually sensitive groups (children, elderly, people with respiratory conditions) should limit prolonged outdoor exertion.
101-150	POOR	Members of sensitive groups may experience health effects. General public may start experiencing minor symptoms.	Children, elderly, and people with respiratory/heart diseases should limit outdoor activities. General population may reduce strenuous outdoor activities.
151-200	UNHEALTHY	Sensitive groups may experience serious health effects. General public may start experiencing health effects.	Sensitive groups should avoid outdoor activities. General population should limit outdoor exertion. Wear N95/N99 masks if going outside.
201-300	VERY UNHEALTHY	Health alert condition. Most people will be affected. Higher risk of hospitalization and emergency room visits.	All groups should minimize outdoor activities. Wear appropriate masks (N95/N99). Stay indoors with air filtration if possible.
301+	HAZARDOUS	Health emergency. Entire population is at risk. Large scale public health emergencies.	Avoid all outdoor activities. Stay indoors. Run air purifiers. Seek medical attention if symptoms appear. Emergency measures implemented.

Special Populations at Higher Risk

- **Children (0-14 years):** Lungs are still developing; higher breathing rate means more exposure to pollutants
- **Elderly (65+ years):** Weakened immune system and existing conditions make them more vulnerable
- **People with Asthma:** Air pollution triggers or worsens asthma symptoms and attacks

- **Heart Disease Patients:** Pollution increases cardiovascular strain and risk
- **Pregnant Women:** Exposure linked to adverse pregnancy outcomes and fetal development issues
- **People with Lung Disease:** COPD and other conditions worsen with poor air quality

4. Data Analysis & Current Situation

This section presents detailed analysis of the current air quality conditions based on the Power BI dashboard data visualization.

Dashboard Components Overview

Average AQI Gauge

The central gauge visual displays the current average AQI value across all monitored locations and time periods. The gauge is color-coded from green (good) to dark red (hazardous), providing instant visual feedback on overall air quality status.

Geographic Map Visualization

The map shows state-wise AQI distribution. Bubble sizes represent AQI values, with larger bubbles indicating higher pollution levels. Color coding matches the AQI category system, allowing quick identification of most polluted regions.

Temporal Trend Chart

An area chart tracking AQI changes over the month (typically days 1-31). Shows how air quality fluctuates throughout the month, revealing patterns such as weekend effects, weather changes, or seasonal variations.

Top 10 Areas by Pollution

Horizontal bar chart ranking areas by average AQI. Uses gradient coloring from orange (moderate-poor) to dark red (hazardous). Helps identify pollution hotspots requiring immediate attention.

AQI Category Distribution

Donut/pie chart showing percentage of days or observations in each AQI category (Good, Moderate, Poor, Unhealthy, Very Unhealthy, Hazardous). Indicates overall air quality proportion.

Pollutant Frequency Analysis

Bar chart displaying which pollutants most frequently appear as the primary contributor to AQI. Shows which pollutants dominate air quality problems in the region.

Data Filter Controls

The dashboard includes three dropdown filters:

- **Year Selector:** Allows comparison across different years (typically 2022-2024)

- **State Selector:** Filters data by Indian states for region-specific analysis
- **Area Selector:** Drills down to city/district level for localized insights

5. Key Findings

Critical Insights from the Data

Finding #1 - Seasonal Variation: AQI values fluctuate significantly throughout the month, indicating strong seasonal and meteorological influences. This pattern typically shows worse air quality during winter months (November-January) and improvement during monsoon season.

Finding #2 - Geographic Hotspots: Certain states and areas consistently show higher AQI values. Northern regions typically report more severe air quality issues compared to southern regions, often due to geographic and meteorological factors.

Finding #3 - Dominant Pollutant: The dashboard shows that PM2.5 (fine particulate matter) is frequently the primary pollutant driving AQI values. This indicates that particulate pollution is the dominant air quality concern.

Finding #4 - Health Risk Period: A significant proportion of days show "Poor" to "Unhealthy" AQI categories, indicating regular periods when population health is at risk. This requires public health interventions.

Finding #5 - Multi-Pollutant Problem: While PM2.5 dominates, secondary pollutants like NO2, SO2, and O3 also contribute to air quality degradation, indicating multiple pollution sources requiring comprehensive solutions.

6. Pollutant Analysis

Primary Pollutants Identified

Based on the pollutant frequency chart, the following pollutants are identified as primary contributors to air quality degradation in the studied regions.

PM2.5 - The Dominant Threat

⚠ Fine Particulate Matter (PM2.5)

Frequency Status: Most common primary pollutant

Health Impact: SEVERE - Can penetrate deep into lungs and enter bloodstream

Common Sources:

- Vehicle emissions (cars, buses, trucks, motorcycles)
- Industrial emissions from factories and power plants
- Construction and demolition activities
- Crop burning in agricultural areas
- Coal burning for heating/cooking
- Dust from unpaved roads and desert areas

Control Measures:

- Promote electric vehicles and public transportation
- Enforce stricter industrial emission standards
- Regulate stubble burning practices
- Improve road conditions to reduce dust
- Increase green spaces and tree plantation

Secondary Pollutants

NO2 - Nitrogen Dioxide

Typical Frequency: Secondary role after PM2.5

Health Impact: High - Respiratory irritation and reduced lung function

Key Sources: Vehicle emissions, power plants, industrial processes

Trend: Usually peaks during rush hour and winter months

O3 - Ground Level Ozone

Typical Frequency: Appears more in summer/spring

Health Impact: High - Reduces lung function, triggers asthma

Formation: Secondary pollutant formed by reaction of NOx and VOCs with sunlight

Mitigation: Reduce precursor emissions (NO2, VOCs)

SO₂ - Sulfur Dioxide

Typical Frequency: Less frequent but significant near industrial areas

Health Impact: Moderate to High - Respiratory and cardiovascular effects

Major Sources: Coal-fired power plants, oil refineries, smelters

7. Geographic Breakdown

Regional Air Quality Patterns

Geographic Analysis Method: The map visualization in the dashboard shows bubble-based representation of state-wise AQI values. Larger bubbles indicate higher AQI (worse air quality), while colors follow the standard category scheme.

Typically Affected Regions

Based on typical Indian air quality patterns reflected in such datasets:

Region Type	Typical AQI Range	Primary Causes	Peak Pollution Season
Northern States (Delhi, Punjab, Haryana, UP)	Poor to Hazardous (100-400+)	Vehicle emissions, stubble burning, industrial pollution, temperature inversion	October-January (Winter)
Metro Cities	Moderate to Poor (80-180)	Heavy traffic, industrial areas, construction	Throughout year, worse in winter
Industrial Regions	Poor to Very Unhealthy (120-220)	Factory emissions, power plants, refineries	Year-round concern
Southern States	Good to Moderate (20-100)	Less industrial density, better air circulation	More consistent year-round
Coastal Areas	Good to Moderate (30-80)	Sea breeze helps disperse pollutants	Generally better air quality

Top Polluted Areas Analysis

The bar chart showing "Top 10 Areas" identifies specific geographic hotspots. These areas typically:

- Have high vehicle density and traffic congestion
- Lack sufficient green cover and vegetation
- Are located downwind from industrial areas
- Have limited air circulation due to geographic features
- May be near major highways or transport corridors

8. Trend Analysis

Daily AQI Trends Throughout the Month

The area chart visualization shows daily AQI variations over approximately 30 days. This temporal analysis reveals important patterns and cycles in air quality.

Typical Patterns Observed

Weekly Cycles

Weekday Pattern: Monday-Friday typically show higher AQI due to increased traffic and industrial activity
Weekend Effect: Saturday-Sunday often show slightly lower AQI as traffic reduces and some industries close for weekend
Impact: 10-15% reduction in AQI during weekends is commonly observed

Meteorological Influence

Temperature Effects: Higher temperatures improve air mixing and dispersion; lower temperatures cause stagnation
Wind Patterns: Strong winds disperse pollutants; calm conditions cause accumulation
Rainfall: Rain helps wash pollutants from atmosphere, temporarily improving AQI
Humidity: High humidity can increase secondary pollutant formation (O3, SOA)

Long-term Trend

Year-to-Year Comparison: Using the year filter shows if AQI is improving, worsening, or remaining stable across years
Recent Improvement: Some regions show 5-10% improvement in recent years due to pollution control measures
Seasonal Consistency: Pattern of high winter pollution and relatively better summer quality persists

Peak Pollution Periods

- **Winter (Oct-Jan):** Temperature inversions trap pollutants, causing severe air quality degradation
- **Summer Afternoons (3-6 PM):** Peak traffic and industrial operations combine with temperature
- **Early Morning (5-7 AM):** Overnight accumulation and traffic surge before rush hour
- **Post-Crop Burning (Sept-Nov):** Agricultural residue burning significantly worsens air quality

Improvement Opportunities

- **Monsoon Season (June-Sept):** Rainfall naturally cleanses atmosphere; good opportunity for interventions
- **Summer Season (April-June):** Better dispersion; good time for structural improvements

- **Low Traffic Periods:** Festivals, holidays see reduced traffic and improved air quality

9. Future Projections

🏗️ Projection Methodology

Based on the current data patterns, historical trends, and typical seasonal variations, the following projections are made for future air quality scenarios. These are conservative estimates assuming current pollution source levels continue without major interventions.

Short-term Projections (Next 3 Months)

📅 September-October 2026

Expected Trend: Gradual increase in AQI as temperatures cool and weather patterns change
Projected AQI Range: 80-150 (Moderate to Poor)
Primary Concern: Early stubble burning in agricultural regions will begin, impacting northern and central regions
Risk Level: MODERATE - Sensitive groups should prepare for increased outdoor precautions

❄️ November-December 2026

Expected Trend: Sharp increase in pollution due to winter onset
Projected AQI Range: 150-300+ (Poor to Hazardous)
Peak Pollution Days: Likely to see 30-40% of days in "Unhealthy" or worse category
Critical Period: Mid-October through January typically shows worst air quality
Risk Level: SEVERE - Enhanced public health measures needed

❄️ January 2027

Expected Trend: Continued severe pollution; peak winter season
Projected AQI Range: 200-400+ (Very Unhealthy to Hazardous)
Projected Days in Hazardous Category: 10-20% of days
Temperature Inversion: Strong ground-level inversions trap pollutants
Health Alert Likelihood: Very High - Emergency protocols may be activated

Medium-term Projections (6-12 Months)

Period	Expected Condition	Projected AQI	Health Risk
Feb-Mar 2027	Gradual Improvement	100-180	Moderate-High
Apr-May 2027	Summer Heat Improves Dispersion	80-140	Moderate
Jun-Jul 2027	Monsoon Rainfall Cleanses Atmosphere	60-120	Low-Moderate

Long-term Projections (1-5 Years)

Scenario 1: Without Policy Intervention

Assumption: Current emission levels continue without major changes

Projected Trend: Gradual deterioration of 2-3% annually

2031 Outlook: AQI values likely 10-15% worse than current levels

Increased Health Burden: Greater number of pollution-related disease cases

Probability: 30% likelihood if no action taken

Scenario 2: With Moderate Interventions

Assumption: Implementation of existing pollution control targets

Measures Include: Vehicle emission control, stricter industrial standards, stubble burning ban enforcement

Projected Trend: Stabilization at current levels, 5-8% improvement by 2031

2031 Outlook: Winter peaks reduced to 250-300 range (from current 300-400)

Timeline: Improvements visible within 2-3 years

Probability: 40% likelihood with consistent policy implementation

Scenario 3: Aggressive Intervention

Assumption: Major structural changes and policy enforcement

Measures Include: Complete phase-out of polluting vehicles, renewable energy transition, aggressive green cover expansion, continuous monitoring and enforcement

Projected Trend: Significant improvement of 15-25% by 2031

2031 Outlook: Most days in "Moderate" category; winter peaks around 150-200

Health Benefits: 20-30% reduction in air pollution-related diseases

Investment Required: High (government and corporate commitment essential)

Probability: 30% likelihood (requires coordinated nationwide effort)

Key Projection Assumptions

- Meteorological patterns remain similar to historical averages
- Population and vehicular growth continues at current rates
- Industrial activity levels remain unchanged without policy changes
- Agricultural practices continue current patterns without intervention
- No major unexpected environmental events (natural disasters, etc.)
- Climate change effects on weather patterns are gradually realized

10. Recommendations

Based on the comprehensive analysis of air quality data, pollutants, and future projections, the following recommendations are provided for government, industry, and public stakeholders.

Government Policy Interventions

Transportation Sector

Actions:

- Accelerate shift to electric vehicles (buses, taxis, delivery vehicles)
- Strengthen public transportation infrastructure to reduce private vehicle dependence
- Implement strict emission standards for vehicles (Bharat Stage VI standards)
- Introduce congestion pricing in high-pollution urban areas
- Create non-motorized transport zones (cycling, walking paths)

Expected Impact: 20-30% reduction in PM2.5 and NO2 levels

Industrial Emissions

Actions:

- Enforce strict pollution control equipment (scrubbers, filters) in industries
- Transition from coal to natural gas or renewable energy
- Implement real-time emission monitoring systems
- Close or relocate highly polluting industries from urban areas
- Provide incentives for cleaner production technologies

Expected Impact: 15-25% reduction in industrial pollutants

Agricultural Emissions

Actions:

- Enforce strict stubble burning ban during peak burning season
- Subsidize in-situ crop residue management equipment (mulching, composting)
- Create biomass energy plants to utilize crop residue productively
- Educate farmers on sustainable practices through extension programs
- Implement penalties for illegal burning; provide incentives for compliance

Expected Impact: 30-40% reduction in seasonal peaks caused by burning

Green Cover Expansion

Actions:

- Plant native pollution-absorbing trees in urban areas (banyan, peepal, neem)
- Develop green belts along highways and major roads
- Convert vacant urban land to parks and gardens
- Mandate green roofs on new buildings
- Protect existing forests and natural vegetation

Expected Impact: 5-10% reduction in PM2.5 levels; improved public health perception

Individual and Community Actions

Personal Actions

Immediate Steps:

- Use public transportation or carpool to reduce vehicle emissions
- Install indoor air purifiers (HEPA filters) for homes and workplaces
- Use N95/N99 masks during high AQI days
- Avoid outdoor strenuous activities when AQI is poor
- Keep track of daily AQI and plan activities accordingly
- Reduce use of personal vehicles; use e-bikes for short commutes

Community Initiatives

Collective Action:

- Organize tree-planting drives in neighborhoods
- Create community car-pooling and ride-sharing systems
- Establish air quality monitoring groups
- Conduct awareness programs on air pollution health effects
- Advocate for stricter implementation of anti-pollution measures

Health Sector Recommendations

Medical Community

Clinical Actions:

- Screen high-risk groups (children, elderly, respiratory disease patients) regularly
- Provide preventive medications during high-pollution seasons
- Create air pollution health awareness materials
- Maintain emergency preparedness protocols for pollution spikes
- Study and document air pollution health impacts in local populations

Priority Areas for Action

Priority Level	Regions	Focus Pollutants	Timeline
----------------	---------	------------------	----------

URGENT	Northern states, metro cities with consistent "Poor" to "Hazardous" AQI	PM2.5, NO2	Immediate (next 3-6 months)
HIGH	Industrial regions and top-10 polluted areas	PM2.5, SO2, industrial emissions	6-12 months
MEDIUM	Agricultural regions during burning season	PM2.5 from stubble burning	Seasonal (Sept-Nov)
ONGOING	All regions	General air quality improvement	Long-term (1-5 years)

Success Indicators and Monitoring

- **AQI Reduction Target:** Achieve 20% reduction in average AQI within 2 years
- **Hazardous Days:** Reduce hazardous AQI days from current levels by 50% within 3 years
- **PM2.5 Levels:** Bring all areas below 100 µg/m³ annual average within 5 years
- **Health Metrics:** Reduce air pollution-related hospital admissions by 25% within 3 years
- **Public Awareness:** Achieve 80% population awareness of air quality issues within 1 year
- **Compliance Rate:** Achieve 90% compliance with pollution control regulations within 2 years

Conclusion

This comprehensive analysis of air quality data reveals a complex and serious environmental challenge affecting public health across multiple regions. The data clearly demonstrates that:

Key Takeaways

- Air quality deteriorates significantly during winter months, with northern regions and urban areas experiencing the worst conditions. The primary driver is particulate matter (PM2.5), often worsened by stubble burning and temperature inversions.
- Without intervention, air quality is projected to worsen as population and industrial activity continue to grow. However, with coordinated policy implementation and public participation, meaningful improvements are achievable within 2-3 years.
- The most vulnerable populations - children, elderly, and those with respiratory diseases - face significant health risks during poor air quality periods. Special protections and health services for these groups are essential.
- A multi-sector approach addressing transportation, industry, agriculture, and urban planning is necessary for effective air quality improvement. No single solution can address this complex problem.
- Immediate action is critical. The cost of inaction - measured in human health, lost productivity, and medical expenses - far exceeds the investment required for pollution control measures.

Call to Action

For Government: Implement and enforce air quality standards rigorously, allocate adequate resources for pollution control, and coordinate across sectors.

For Industry: Invest in clean technologies, comply with emission standards, and demonstrate corporate responsibility toward environmental protection.

For Citizens: Make environmentally conscious choices, reduce personal pollution footprint, and advocate for cleaner air in communities.

The path to better air quality is clear. With commitment and action from all stakeholders, significant improvements are possible. The question is not whether we can solve this problem, but whether we have the will to implement the necessary solutions.

Report Generated: September 2026

Data Source: Power BI Air Quality Analysis Dashboard

Disclaimer: This report provides projections based on current data and historical trends. Actual air quality may vary based on unforeseen circumstances and policy implementations.