



# Discrete Mathematics

## Counting



# The Basics of Counting


## Section 6.1



# Basic Counting Principles: The Product Rule

Discrete  
Mathematics

**The Product Rule:** A procedure can be broken down into a sequence of two tasks. There are  $n_1$  ways to do the first task and  $n_2$  ways to do the second task. Then there are  $n^1 \cdot n^2$  ways to do the procedure.



**Example:** How many bit strings of length seven are there?(arrangement)

**Solution:** Since each of the seven bits is either a 0 or a 1, the answer is  $2_7 = 128$ .



# Counting Functions

Discrete  
Mathematics

**Counting Functions:** How many functions are there from a set with  $m$  elements to a set with  $n$  elements?

**Solution:** Since a function represents a choice of one of the  $n$  elements of the codomain for each of the  $m$  elements in the domain, the product rule tells us that there are

$n \cdot n \cdot \dots \cdot n = n^m$  functions.



# Counting Subsets of a Finite Set

Discrete  
Mathematics

**Counting Subsets of a Finite Set:** Use the product rule to show that the number of different subsets of a finite set  $S$  is  $2^{|S|}$ .

**Solution:** When the elements of  $S$  are listed in an arbitrary order, there is a one-to-one correspondence between subsets of  $S$  and bit strings of length  $|S|$ . When the  $i$ th element is in the subset, the bit string has a 1 in the  $i$ th position and a 0 otherwise.

By the product rule, there are  $2^{|S|}$  such bit strings, and therefore  $2^{|S|}$  subsets.



# Product Rule in Terms of Sets

Discrete  
Mathematics


- If  $A_1, A_2, \dots, A_m$  are finite sets, then the number of elements in the Cartesian product of these sets is the product of the number of elements of each set.
- The task of choosing an element in the Cartesian product  $A_1 \times A_2 \times \dots \times A_m$  is done by choosing an element in  $A_1$ , an element in  $A_2$ , ..., and an element in  $A_m$ .
- By the product rule, it follows that:  
$$|A_1 \times A_2 \times \dots \times A_m| = |A_1| \cdot |A_2| \cdot \dots \cdot |A_m|.$$



# Basic Counting Principles: The Sum Rule

Discrete  
Mathematics

**The Sum Rule:** If a task can be done either in one of  $n_1$  ways or in one of  $n_2$  ways to do the second task, where none of the set of  $n_1$  ways is the same as any of the  $n_2$  ways, then there are  $n_1 + n_2$  ways to do the task.





# Basic Counting Principles: The Sum Rule

---

Discrete  
Mathematics

**Example:** The mathematics department must choose either a student or a faculty member as a representative for a university committee. How many choices are there for this representative if there are 37 members of the mathematics faculty and 83 mathematics majors and no one is both a faculty member and a student.

**Solution:** By the sum rule it follows that there are  $37 + 83 = 120$  possible ways to pick a representative.





# The Sum Rule in terms of sets.

Discrete  
Mathematics

- The sum rule can be phrased in terms of sets.

$|A \cup B| = |A| + |B|$  as long as  $A$  and  $B$  are disjoint sets.

- Or more generally,

$$|A_1 \cup A_2 \cup \cdots \cup A_m| = |A_1| + |A_2| + \cdots + |A_m|$$

when  $A_i \cap A_j = \emptyset$  for all  $i, j$ .



## Combining the Sum and Product Rule

---

Discrete  
Mathematics

**Example:** Suppose statement labels in a programming language can be either a single letter or a letter followed by a digit. Find the number of possible labels.

**Solution:** Use the product rule.

$$26 + 26 \cdot 10 = 286$$



# Basic Counting Principles: Subtraction Rule

---

Discrete  
Mathematics

**Subtraction Rule:** If a task can be done either in one of  $n_1$  ways or in one of  $n_2$  ways, then the total number of ways to do the task is  $n_1 + n_2$  minus the number of ways to do the task that are common to the two different ways.

- Also known as, *the principle of inclusion-exclusion*:

$$|A \cup B| = |A| + |B| - |A \cap B|$$




# Counting Bit Strings

Discrete  
Mathematics

**Example:** How many bit strings of length eight either start with a 1 bit or end with the two bits 00?

**Solution:** Use the subtraction rule.

- Number of bit strings of length eight that start with a 1 bit:  $2^7 = 128$
- Number of bit strings of length eight that start with bits 00:  $2^6 = 64$
- Number of bit strings of length eight that start with a 1 bit and end with bits 00 :  $2^5 = 32$



Hence, the number is  $128 + 64 - 32 = 160$ .



# Basic Counting Principles:

## Division Rule

---

Discrete  
Mathematics

**Division Rule:** There are  $n/d$  ways to do a task if it can be done using a procedure that can be carried out in  $n$  ways, and **for every way  $w$ , exactly  $d$  of the  $n$  ways correspond to way  $w$ .**

- Restated in **terms of sets**: If the finite set  $A$  is the union of  $n$  pairwise disjoint subsets each with  $d$  elements, then  $n = |A|/d$ .
- In **terms of functions**: If  $f$  is a function from  $A$  to  $B$ , where both are finite sets, and for every value  $y \in B$  there are exactly  $d$  values  $x \in A$  such that  $f(x) = y$ , then  $|B| = |A|/d$ .



# Division Rule

**Example:** How many ways are there to seat four people around a circular table, where two seatings are considered the same when each person has the same left and right neighbor?

**Solution:** Number the seats around the table from 1 to 4 proceeding clockwise. There are four ways to select the person for seat 1, 3 for seat 2, 2, for seat 3, and one way for seat 4. Thus there are  $4! = 24$  ways to order the four people. But since two seatings are the same when each person has the same left and right neighbor, for every choice for seat 1, we get the same seating.

Therefore, by the division rule, there are  $24/4 = 6$  different seating arrangements.




# Tree Diagrams

Discrete  
Mathematics

- **Tree Diagrams:** We can solve many counting problems through the use of *tree diagrams*, where a branch represents a possible choice and the leaves represent possible outcomes.
- **Example:** Suppose that “I Love Discrete Math” T-shirts come in five different sizes: S,M,L,XL, and XXL. Each size comes in four colors (white, red, green, and black), except XL, which comes only in red, green, and black, and XXL, which comes only in green and black. What is the minimum number of stores that the campus book store needs to stock to have one of each size and color available?
- **Solution:** Draw the tree diagram.

W = white, R = red, G = green, B = black



- The store must stock 17 T-shirts.



西北工业大学  
NORTHWESTERN POLYTECHNICAL UNIVERSITY



# The Pigeonhole Principle (鸽笼原理)

Section 6.2







# The Pigeonhole Principle

Discrete  
Mathematics


- If a flock of 20 pigeons roosts in a set of 19 pigeonholes, one of the pigeonholes must have more than 1 pigeon.



(a)



(b)



(c)

**Pigeonhole Principle:** If  $k$  is a positive integer and  $k + 1$  objects are placed into  $k$  boxes, then at least one box contains two or more objects.

**Proof:** We use a proof by contraposition. Suppose none of the  $k$  boxes has more than one object. Then the total number of objects would be at most  $k$ . This contradicts the statement that we have  $k + 1$  objects.





# The Pigeonhole Principle

Discrete  
Mathematics

**Corollary 1:** A function  $f$  from a set with  $k + 1$  elements to a set with  $k$  elements is not one-to-one.

**Proof:** Use the pigeonhole principle.

- Create a box for each element  $y$  in the codomain of  $f$ .
- Put in the box for  $y$  all of the elements  $x$  from the domain such that  $f(x) = y$ .
- Because there are  $k + 1$  elements and only  $k$  boxes, at least one box has two or more elements.

Hence,  $f$  can't be one-to-one.





# Pigeonhole Principle

---

Discrete  
Mathematics

**Example:** Among any group of 367 people, there must be at least two with the same birthday, because there are only 366 possible birthdays.



# The Generalized Pigeonhole Principle

Discrete  
Mathematics

**The Generalized Pigeonhole Principle:** If  $N$  objects are placed into  $k$  boxes, then there is at least one box containing at least  $\lceil N/k \rceil$  objects.

**Proof:** We use a proof by contraposition. Suppose that none of the boxes contains more than  $\lceil N/k \rceil - 1$  objects. Then the total number of objects is at most

$$k \left( \left\lceil \frac{N}{k} \right\rceil - 1 \right) < k \left( \left( \frac{N}{k} + 1 \right) - 1 \right) = N,$$

where the inequality  $\lceil N/k \rceil < \lceil N/k \rceil + 1$  has been used. This is a contradiction because there are a total of  $n$  objects.

**Example:** Among 100 people there are at least  $\lceil 100/12 \rceil = 9$  who were born in the same month.





# The Generalized Pigeonhole Principle

---

Discrete  
Mathematics

## Example

What is the minimum number of students required in a discrete mathematics class to be sure that at least six will receive the same grade, if there are five possible grades A, B, C, D and F?

$$\lceil N/5 \rceil = 6 \quad N = 5 * 5 + 1 = 26$$



# Permutations and Combinations (排列组合)

Section 6.3



# Permutations

Discrete  
Mathematics

**Definition:** A *permutation* of a set of distinct objects is an *ordered* arrangement of these objects. An ordered arrangement of  $r$  elements of a set is called an  *$r$ -permutation*.

**Example:** Let  $S = \{1, 2, 3\}$ .

- The ordered arrangement 3,1,2 is a permutation of  $S$ .
- The ordered arrangement 3,2 is a 2-permutation of  $S$ .
- The number of  $r$ -permutations of a set with  $n$  elements is denoted by  $P(n, r)$ .
  - The 2-permutations of  $S = \{1, 2, 3\}$  are 1,2; 1,3; 2,1; 2,3; 3,1; and 3,2. Hence,  $P(3, 2) = 6$ .



# A Formula for the Number of Permutations

Discrete  
Mathematics

**Theorem 1:** If  $n$  is a positive integer and  $r$  is an integer with  $1 \leq r \leq n$ , then there are

$$P(n, r) = n(n - 1)(n - 2) \cdots (n - r + 1)$$

$r$ -permutations of a set with  $n$  distinct elements.

**Proof:** Use the product rule. The first element can be chosen in  $n$  ways. The second in  $n - 1$  ways, and so on until there are  $(n - (r - 1))$  ways to choose the last element.

- Note that  $P(n, 0) = 1$ , since there is only one way to order zero elements. Empty list

**Corollary 1:** If  $n$  and  $r$  are integers with  $1 \leq r \leq n$ , then

$$P(n, r) = \frac{n!}{(n-r)!}$$





# Solving Counting Problems by Counting Permutations

---

Discrete  
Mathematics

**Example:** How many ways are there to select a first-prize winner, a second prize winner, and a third-prize winner from 100 different people who have entered a contest?

**Solution:**

$$P(100,3) = 100 \cdot 99 \cdot 98 = 970,200$$



# Solving Counting Problems by Counting Permutations (*continued*)

---

Discrete  
Mathematics

**Example:** How many permutations of the letters *ABCDEFGH* contain the string *ABC* ?

**Solution:** We solve this problem by counting the permutations of six objects, *ABC*, *D*, *E*, *F*, *G*, and *H*.

$$6! = 6 \cdot 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1 = 720$$



# Combinations

Discrete  
Mathematics

**Definition:** An  *$r$ -combination* of elements of a set is an **unordered selection** of  $r$  elements from the set. Thus, an  $r$ -combination is simply a subset of the set with  $r$  elements.

- The number of  $r$ -combinations of a set with  $n$  distinct elements is denoted by  $C(n, r)$ . The notation  $\binom{n}{r}$  is also used and is called a ***binomial coefficient***. (We will see the notation again in the binomial theorem in Section 6.4.)

**Example:** Let  $S$  be the set  $\{a, b, c, d\}$ . Then  $\{a, c, d\}$  is a 3-combination from  $S$ . It is the same as  $\{d, c, a\}$  since the order listed does not matter.

- $C(4, 2) = 6$  because the 2-combinations of  $\{a, b, c, d\}$  are the six subsets  $\{a, b\}$ ,  $\{a, c\}$ ,  $\{a, d\}$ ,  $\{b, c\}$ ,  $\{b, d\}$ , and  $\{c, d\}$ .



# Combinations

Discrete  
Mathematics

**Theorem 2:** The number of  $r$ -combinations of a set with  $n$  elements, where  $n \geq r \geq 0$ , equals

$$C(n, r) = \frac{n!}{(n-r)!r!}.$$

**Proof:** By the product rule  $P(n, r) = C(n, r) \cdot P(r, r)$ . Therefore,

$$C(n, r) = \frac{P(n, r)}{P(r, r)} = \frac{n!/(n-r)!}{r!/(r-r)!} = \frac{n!}{(n-r)!r!}.$$



# Combinations

Discrete  
Mathematics

**Example:** How many poker hands of five cards can be dealt from a standard deck of 52 cards? Also, how many ways are there to select 47 cards from a deck of 52 cards?

**Solution:** Since the order in which the cards are dealt does not matter, the number of five card hands is:

$$\begin{aligned} C(52, 5) &= \frac{52!}{5!47!} \\ &= \frac{52 \cdot 51 \cdot 50 \cdot 49 \cdot 48}{5 \cdot 4 \cdot 3 \cdot 2 \cdot 1} = 26 \cdot 17 \cdot 10 \cdot 49 \cdot 12 = 2,598,960 \end{aligned}$$

- The different ways to select 47 cards from 52 is

$$C(52, 47) = \frac{52!}{47!5!} = C(52, 5) = 2,598,960.$$

*This is a special case of a general result. →*



# Combinations

Discrete  
Mathematics

**Corollary 2:** Let  $n$  and  $r$  be nonnegative integers with  $r \leq n$ . Then  $C(n, r) = C(n, n - r)$ .

**Proof:** From Theorem 2, it follows that

$$C(n, r) = \frac{n!}{(n-r)!r!}$$

and

$$C(n, n - r) = \frac{n!}{(n-r)![n-(n-r)]!} = \frac{n!}{(n-r)!r!} .$$



Hence,  $C(n, r) = C(n, n - r)$ .



# Combinations

Discrete  
Mathematics

**Example:** How many ways are there to select five players from a 10-member tennis team to make a trip to a match at another school.

**Solution:** By Theorem 2, the number of combinations is

$$C(10, 5) = \frac{10!}{5!5!} = 252.$$

**Example:** A group of 30 people have been trained as astronauts to go on the first mission to Mars. How many ways are there to select a crew of six people to go on this mission?

**Solution:** By Theorem 2, the number of possible crews is

$$C(30, 6) = \frac{30!}{6!24!} = \frac{30 \cdot 29 \cdot 28 \cdot 27 \cdot 26 \cdot 25}{6 \cdot 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1} = 593,775 .$$



# Binomial Coefficients and Identities (二项式系数)

Section 6.4





# Powers of Binomial Expressions

Discrete  
Mathematics

**Definition:** A *binomial expression* is the sum of two terms, such as  $x + y$ . (More generally, these terms can be products of constants and variables.)

- We can use counting principles to find the coefficients in the expansion of  $(x + y)^n$  where  $n$  is a positive integer.
- To illustrate this idea, we first look at the process of expanding  $(x + y)^3$ .
- $(x + y)(x + y)(x + y)$  expands into a sum of terms that are the product of a term from each of the three sums.
- Terms of the form  $x^3, x^2y, xy^2, y^3$  arise. The question is what are the coefficients?
  - To obtain  $x^3$ , an  $x$  must be chosen from each of the sums. There is only one way to do this. So, the coefficient of  $x^3$  is 1.
  - To obtain  $x^2y$ , an  $x$  must be chosen from two of the sums and a  $y$  from the other. There are  $\binom{3}{2}$  ways to do this and so the coefficient of  $x^2y$  is 3.
  - To obtain  $xy^2$ , an  $x$  must be chosen from one of the sums and a  $y$  from the other two. There are  $\binom{3}{1}$  ways to do this and so the coefficient of  $xy^2$  is 3.
  - To obtain  $y^3$ , a  $y$  must be chosen from each of the sums. There is only one way to do this. So, the coefficient of  $y^3$  is 1.
- We have used a counting argument to show that  $(x + y)^3 = x^3 + 3x^2y + 3xy^2 + y^3$ .
- Next we present the binomial theorem gives the coefficients of the terms in the expansion of  $(x + y)^n$ .



# Binomial Theorem

Discrete  
Mathematics

**Binomial Theorem:** Let  $x$  and  $y$  be variables, and  $n$  a nonnegative integer. Then:

$$(x+y)^n = \sum_{j=0}^n \binom{n}{j} x^{n-j} y^j = \binom{n}{0} x^n + \binom{n}{1} x^{n-1} y + \cdots + \binom{n}{n-1} x y^{n-1} + \binom{n}{n} y^n.$$

**Proof:** We use combinatorial reasoning . The terms in the expansion of  $(x + y)^n$  are of the form  $x^{n-k}y^k$  for  $k = 0, 1, 2, \dots, n$ . To form the term  $x^{n-k}y^k$ , it is necessary to choose  $n-k$   $x$ s from the  $n$  sums.





# Using the Binomial Theorem

Discrete  
Mathematics

**Example:** What is the coefficient of  $x^{12}y^{13}$  in the expansion of  $(2x - 3y)^{25}$ ?

**Solution:** We view the expression as  $(2x + (-3y))^{25}$ . By the binomial theorem

$$(2x + (-3y))^{25} = \sum_{j=0}^{25} \binom{25}{j} (2x)^{25-j} (-3y)^j.$$

Consequently, the coefficient of  $x^{12}y^{13}$  in the expansion is obtained when  $j = 13$ .

$$\binom{25}{13} 2^{12} (-3)^{13} = -\frac{25!}{13!12!} 2^{12} 3^{13}.$$



# A Useful Identity

Discrete  
Mathematics

**Corollary 1:** With  $n \geq 0$ ,

$$\sum_{k=0}^n \binom{n}{k} = 2^n.$$

**Proof (using binomial theorem):** With  $x = 1$  and  $y = 1$ , from the binomial theorem we see that:

$$2^n = (1 + 1)^n = \sum_{k=0}^n \binom{n}{k} 1^k 1^{(n-k)} = \sum_{k=0}^n \binom{n}{k}.$$

**Proof (combinatorial):** Consider the subsets of a set with  $n$  elements. There are  $\binom{n}{0}$  subsets with zero elements,  $\binom{n}{1}$  with one element,  $\binom{n}{2}$  with two elements, ..., and  $\binom{n}{n}$  with  $n$  elements. Therefore the total is


$$\sum_{k=0}^n \binom{n}{k}.$$

Since, we know that a set with  $n$  elements has  $2^n$  subsets, we conclude:

$$\sum_{k=0}^n \binom{n}{k} = 2^n.$$



# Pascal's Identity



discrete  
mathematics

**Pascal's Identity:** If  $n$  and  $k$  are integers with  $n \geq k \geq 0$ , then

$$\binom{n+1}{k} = \binom{n}{k-1} + \binom{n}{k}.$$

Blaise Pascal  
(1623-1662)

**Proof (combinatorial):** Let  $T$  be a set where  $|T| = n + 1$ ,  $a \in T$ , and  $S = T - \{a\}$ . There are  $\binom{n+1}{k}$  subsets of  $T$  containing  $k$  elements. Each of these subsets either:

- contains  $a$  with  $k - 1$  other elements, or
- contains  $k$  elements of  $S$  and not  $a$ .

There are

- $\binom{n}{k-1}$  subsets of  $k$  elements that contain  $a$ , since there are  $\binom{n}{k-1}$  subsets of  $k - 1$  elements of  $S$ ,
- $\binom{n}{k}$  subsets of  $k$  elements of  $T$  that do not contain  $a$ , because there are  $\binom{n}{k}$  subsets of  $k$  elements of  $S$ .

Hence,

$$\binom{n+1}{k} = \binom{n}{k-1} + \binom{n}{k}.$$



See Exercise 19  
for an algebraic  
proof.




# Pascal's Triangle

Discrete  
Mathematics

杨辉三角

The  $n$ th row in the triangle consists of the binomial coefficients  $k = 0, 1, \dots, n$ .  $\binom{n}{k}$



By Pascal's identity, adding two adjacent binomial coefficients results in the binomial coefficient in the next row between these two coefficients.



# Homework

---

Discrete  
Mathematics

- § 6.1 16,51,55  
§ 6.2 8,16,21  
§ 6.3 11, 24,28,30  
§ 6.4 7,8