



Sequences and Summations


Section 2.4



Section Summary

Discrete
Mathematics

- Sequences.
 - Examples: Geometric Progression, Arithmetic Progression
- Recurrence Relations
 - Example: Fibonacci Sequence
- Summations



Sequences

Discrete
Mathematics

Definition: A *sequence* is a function from a subset of the integers (usually either the set $\{0, 1, 2, 3, 4, \dots\}$ or $\{1, 2, 3, 4, \dots\}$) to a set S .

- The notation a_n is used to denote the image of the integer n . We can think of a_n as the equivalent of $f(n)$ where f is a function from $\{0, 1, 2, \dots\}$ to S . We call a_n a *term* of the sequence.



Sequences

Discrete
Mathematics

Example: Consider the sequence $\{a_n\}$ where

$$a_n = \frac{1}{n}$$

$$1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4} \cdots$$



Geometric Progression

Discrete
Mathematics

Definition: A *geometric progression* (等比数列) is a sequence of the form:
where the *initial term* a and the *common ratio* r are real numbers. $a, ar, ar^2, \dots, ar^n, \dots$

Examples:

1. Let $a = 1$ and $r = -1$. Then:

$$\{b_n\} = \{b_0, b_1, b_2, b_3, b_4, \dots\} = \{1, -1, 1, -1, 1, \dots\}$$

2. Let $a = 2$ and $r = 5$. Then:

$$\{c_n\} = \{c_0, c_1, c_2, c_3, c_4, \dots\} = \{2, 10, 50, 250, 1250, \dots\}$$

3. Let $a = 6$ and $r = 1/3$. Then:

$$\{d_n\} = \{d_0, d_1, d_2, d_3, d_4, \dots\} = \{6, 2, \frac{2}{3}, \frac{2}{9}, \frac{2}{27}, \dots\}$$



Arithmetic Progression

Discrete
Mathematics

Definition: A *arithmetic progression* (等差数列) is a sequence of the form:

$$a, a + d, a + 2d, \dots, a + nd, \dots$$

where the *initial term* a and the *common difference* d are real numbers.

Examples:

1. Let $a = -1$ and $d = 4$:

-1, 3, 7, 11, 15, ...

2. Let $a = 7$ and $d = -3$:

7, 4, 1, -2, -5, ...



Strings

Discrete
Mathematics

- Sequences of characters or bits are important in computer science.
- The finite sequences are also called strings
- The *empty string* is represented by λ .
- The string *abcde* has *length* 5.



Recurrence Relations

Discrete
Mathematics

Definition: A *recurrence relation* for the sequence $\{a_n\}$ is an equation that expresses a_n in terms of one or more of the previous terms of the sequence, namely, a_0, a_1, \dots, a_{n-1} , for all integers n with $n \geq n_0$, where n_0 is a nonnegative integer.

- A sequence is called a *solution* of a recurrence relation if its terms satisfy the recurrence relation.
- The *initial conditions* for a sequence specify the terms that precede the first term where the recurrence relation takes effect.



Questions about Recurrence Relations

Discrete
Mathematics

Example 1: Let $\{a_n\}$ be a sequence that satisfies the recurrence relation $a_n = a_{n-1} + 3$ for $n = 1, 2, 3, 4, \dots$ and suppose that $a_0 = 2$. What are a_1 , a_2 and a_3 ?

[Here $a_0 = 2$ is the initial condition.]

Solution: We see from the recurrence relation that

$$a_1 = a_0 + 3 = 2 + 3 = 5$$

$$a_2 = 5 + 3 = 8$$

$$a_3 = 8 + 3 = 11$$



Questions about Recurrence Relations

Discrete
Mathematics


Example 2: Let $\{a_n\}$ be a sequence that satisfies the recurrence relation $a_n = a_{n-1} - a_{n-2}$ for $n = 2, 3, 4, \dots$ and suppose that $a_0 = 3$ and $a_1 = 5$. What are a_2 and a_3 ?

[Here the initial conditions are $a_0 = 3$ and $a_1 = 5$.]

Solution: We see from the recurrence relation that

$$a_2 = a_1 - a_0 = 5 - 3 = 2$$

$$a_3 = a_2 - a_1 = 2 - 5 = -3$$



Fibonacci Sequence

Discrete
Mathematics

Definition: Define the *Fibonacci sequence*, f_0, f_1, f_2, \dots , by:

- Initial Conditions: $f_0 = 0, f_1 = 1$
- Recurrence Relation: $f_n = f_{n-1} + f_{n-2}$

Example: Find f_2, f_3, f_4, f_5 and f_6 .

Answer:

$$f_2 = f_1 + f_0 = 1 + 0 = 1,$$

$$f_3 = f_2 + f_1 = 1 + 1 = 2,$$

$$f_4 = f_3 + f_2 = 2 + 1 = 3,$$

$$f_5 = f_4 + f_3 = 3 + 2 = 5,$$

$$f_6 = f_5 + f_4 = 5 + 3 = 8.$$



Solving Recurrence Relations

Discrete
Mathematics

- Finding a formula for the n th term of the sequence generated by a recurrence relation is called *solving the recurrence relation*.
- Such a formula is called a *closed formula*.
- Various methods for solving recurrence relations will be covered in Chapter 8 where recurrence relations will be studied in greater depth.
- Here we illustrate by example the method of iteration



Iterative Solution Example

Discrete
Mathematics

Method 1: Working upward, forward substitution

Let $\{a_n\}$ be a sequence that satisfies the recurrence relation $a_n = a_{n-1} + 3$ for $n = 2, 3, 4, \dots$ and suppose that $a_1 = 2$.

$$a_2 = 2 + 3$$

$$a_3 = (2 + 3) + 3 = 2 + 3 \cdot 2$$

$$a_4 = (2 + 2 \cdot 3) + 3 = 2 + 3 \cdot 3$$

.

.

.

$$1) \quad a_n = a_{n-1} + 3 = (2 + 3 \cdot (n - 2)) + 3 = 2 + 3(n -$$



Iterative Solution Example

Discrete
Mathematics

Method 2: Working downward, backward substitution

Let $\{a_n\}$ be a sequence that satisfies the recurrence relation $a_n = a_{n-1} + 3$ for $n = 2, 3, 4, \dots$ and suppose that $a_1 = 2$.

$$\begin{aligned} a_n &= a_{n-1} + 3 \\ &= (a_{n-2} + 3) + 3 = a_{n-2} + 3 \cdot 2 \\ &= (a_{n-3} + 3) + 3 \cdot 2 = a_{n-3} + 3 \cdot 3 \\ &\quad \cdot \\ &\quad \cdot \\ &\quad \cdot \\ &= a_2 + 3(n-2) = (a_1 + 3) + 3(n-2) = 2 + 3(n-1) \end{aligned}$$



Financial Application

Discrete
Mathematics

Example: Suppose that a person deposits \$10,000.00 in a savings account at a bank yielding 11% per year with interest compounded annually. How much will be in the account after 30 years?

Let P_n denote the amount in the account after n years. P_n satisfies the following recurrence relation:

$$P_n = P_{n-1} + 0.11P_{n-1} = (1.11) P_{n-1}$$

with the initial condition $P_0 = 10,000$

Continued on next slide →



Financial Application

Discrete
Mathematics

$$P_n = P_{n-1} + 0.11P_{n-1} = (1.11) P_{n-1}$$

with the initial condition $P_0 = 10,000$

Solution: Forward Substitution

$$P_1 = (1.11)P_0$$

$$P_2 = (1.11)P_1 = (1.11)^2 P_0$$


$$P_3 = (1.11)P_2 = (1.11)^3 P_0$$

:

$$P_n = (1.11)P_{n-1} = (1.11)^n P_0 = (1.11)^n 10,000$$

$$P_n = (1.11)^n 10,000 \text{ (Can prove by induction, covered in Chapter 5)}$$

$$P_{30} = (1.11)^{30} 10,000 = \$228,992.97$$



Useful Sequences

Discrete
Mathematics

TABLE 1 Some Useful Sequences.

<i>nth Term</i>	<i>First 10 Terms</i>
n^2	1, 4, 9, 16, 25, 36, 49, 64, 81, 100, ...
n^3	1, 8, 27, 64, 125, 216, 343, 512, 729, 1000, ...
n^4	1, 16, 81, 256, 625, 1296, 2401, 4096, 6561, 10000, ...
2^n	2, 4, 8, 16, 32, 64, 128, 256, 512, 1024, ...
3^n	3, 9, 27, 81, 243, 729, 2187, 6561, 19683, 59049, ...
$n!$	1, 2, 6, 24, 120, 720, 5040, 40320, 362880, 3628800, ...
f_n	1, 1, 2, 3, 5, 8, 13, 21, 34, 55, 89, ...



Summations

Discrete
Mathematics


- Sum of the terms a_m, a_{m+1}, \dots, a_n from the sequence $\{a_n\}$
- The notation:

$$\sum_{j=m}^n a_j \quad \sum_{j=m}^n a_j \quad \sum_{m \leq j \leq n} a_j$$

represents

$$a_m + a_{m+1} + \dots + a_n$$

- The variable j is called the *index of summation*. It runs through all the integers starting with its *lower limit* m and ending with its *upper limit* n .



Summations

Discrete
Mathematics

- More generally for a set S :

$$\sum_{j \in S} a_j$$

- **Examples:**

$$r^0 + r^1 + r^2 + r^3 + \dots + r^n = \sum_{j=0}^n r^j$$

$$1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \dots = \sum_{i=1}^{\infty} \frac{1}{i}$$

If $S = \{2, 5, 7, 10\}$ then $\sum_{j \in S} a_j = a_2 + a_5 + a_7 + a_{10}$



Geometric Series

Discrete
Mathematics

Sums of terms of geometric progressions

$$\sum_{j=0}^n ar^j = \begin{cases} \frac{ar^{n+1} - a}{r - 1} & r \neq 1 \\ (n + 1)a & r = 1 \end{cases}$$

Proof:

Let $S_n = \sum_{j=0}^n ar^j$

To compute S_n , first multiply both sides of the equality by r and then manipulate the resulting sum as follows:

$$\begin{aligned} rS_n &= r \sum_{j=0}^n ar^j \\ &= \sum_{j=0}^n ar^{j+1} \end{aligned}$$

Continued on next slide →



Geometric Series

Discrete
Mathematics

$$= \sum_{j=0}^n ar^{j+1} \quad \text{From previous slide.}$$

$$= \sum_{k=1}^{n+1} ar^k \quad \text{Shifting the index of summation with } k = j + 1.$$

$$= \left(\sum_{k=0}^n ar^k \right) + (ar^{n+1} - a) \quad \text{Removing } k = n + 1 \text{ term and adding } k = 0 \text{ term.}$$

$$= S_n + (ar^{n+1} - a) \quad \text{Substituting } S \text{ for summation formula}$$

∴

$$rS_n = S_n + (ar^{n+1} - a)$$

$$S_n = \frac{ar^{n+1} - a}{r - 1} \quad \text{if } r \neq 1$$

$$S_n = \sum_{j=0}^n ar^j = \sum_{j=0}^n a = (n+1)a \quad \text{if } r = 1$$



Some Useful Summation Formulae

Discrete
Mathematics

TABLE 2 Some Useful Summation Formulae.

<i>Sum</i>	<i>Closed Form</i>
$\sum_{k=0}^n ar^k \ (r \neq 0)$	$\frac{ar^{n+1} - a}{r - 1}, r \neq 1$
$\sum_{k=1}^n k$	$\frac{n(n+1)}{2}$
$\sum_{k=1}^n k^2$	$\frac{n(n+1)(2n+1)}{6}$
$\sum_{k=1}^n k^3$	$\frac{n^2(n+1)^2}{4}$
$\sum_{k=0}^{\infty} x^k, x < 1$	$\frac{1}{1-x}$
$\sum_{k=1}^{\infty} kx^{k-1}, x < 1$	$\frac{1}{(1-x)^2}$



西北工业大学
NORTHWESTERN POLYTECHNICAL UNIVERSITY



Cardinality of Sets

Section 2.5



Infinite set


Definition: A set S is infinite if and only if there is a one-to-one correspondence (*i.e.*, a bijection) $f:S \rightarrow S$ and $f(S) \subset S$.

1. \mathbb{N} is infinite. $f(x)=2x$

2. \mathbb{Z} is infinite. $f:\mathbb{Z} \rightarrow \mathbb{N}$, $f(x) = \begin{cases} 2x & x \geq 0 \\ -2x-1 & x < 0 \end{cases}$

3. \mathbb{R} is infinite.

$$f:\mathbb{R} \rightarrow \mathbb{R}, \quad f(x) = \begin{cases} x+1 & x \geq 0 \\ x & x < 0 \end{cases}$$



Cardinality

Definition: The *cardinality* of a set A is equal to the cardinality of a set B , denoted


$$|A| = |B|,$$


if and only if there is a one-to-one correspondence (*i.e.*, a bijection) from A to B .

- If there is a one-to-one function (*i.e.*, an injection) from A to B , the cardinality of A is less than or the same as the cardinality of B and we write $|A| \leq |B|$.
- When $|A| \leq |B|$ and A and B have different cardinality, we say that the cardinality of A is less than the cardinality of B and write $|A| < |B|$.

Example

The same cardinality (red line & blue line)





Cardinality

Discrete
Mathematics

- **Definition:** A set that is either finite or has the same cardinality as the set of positive integers (\mathbf{Z}^+) is called *countable*. A set that is not countable is *uncountable*.
- The set of real numbers \mathbf{R} is an uncountable set.
- When an infinite set is countable (*countably infinite*) its cardinality is \aleph_0 (where \aleph is aleph, the 1st letter of the Hebrew alphabet). We write $|S| = \aleph_0$ and say that S has cardinality “aleph null.”



Showing that a Set is Countable

Discrete
Mathematics

- An infinite set is countable if and only if it is possible to list the elements of the set in a sequence (indexed by the positive integers).
- The reason for this is that a one-to-one correspondence f from the set of positive integers to a set S can be expressed in terms of a sequence $a_1, a_2, \dots, a_n, \dots$ where $a_1 = f(1)$, $a_2 = f(2)$, ..., $a_n = f(n)$, ...




Showing that a Set is Countable

Discrete
Mathematics

Example 1: Show that the set of even positive integers E is countable set.

Solution: Let $f(x) = 2x$.



Then f is a bijection from \mathbf{N} to E since f is both one-to-one and onto. To show that it is one-to-one, suppose that $f(n) = f(m)$. Then $2n = 2m$, and so $n = m$. To see that it is onto, suppose that t is an even positive integer. Then $t = 2k$ for some positive integer k and $f(k) = t$. ◀



Showing that a Set is Countable

Discrete
Mathematics

Example 2: Show that the set of integers **Z** is countable.

Solution: Can list in a sequence:

0, 1, -1, 2, -2, 3, -3,

Or can define a bijection from **N** to **Z**:

- When n is even: $f(n) = n/2$
- When n is odd: $f(n) = -(n-1)/2$





The Positive Rational Numbers are Countable

Discrete
Mathematics

- **Definition:** A *rational number* can be expressed as the ratio of two integers p and q such that $q \neq 0$.
 - $\frac{3}{4}$ is a rational number
 - $\sqrt{2}$ is not a rational number.

Example 3: Show that the positive rational numbers are countable.

Solution: The positive rational numbers are countable since they can be arranged in a sequence:

$$r_1, r_2, r_3, \dots$$

The next slide shows how this is done.





The Positive Rational Numbers are Countable

Discrete
Mathematics

First row $q = 1$.
Second row $q = 2$.
etc.

Constructing the List


First list p/q with $p + q = 2$.

Next list p/q with $p + q = 3$

And so on.

$1, \frac{1}{2}, 2, 3, \frac{1}{3}, \frac{1}{4}, \frac{2}{3}, \dots$


Terms not circled
are not listed
because they
repeat previously
listed terms





The Real Numbers are Uncountable

Georg Cantor
(1845-1918)



crete
ematics

Example: Show that the set of real numbers is uncountable.

Solution: The method is called the Cantor diagonalization argument, and is a proof by contradiction.

1. Suppose \mathbf{R} is countable. Then the real numbers between 0 and 1 are also countable (any subset of a countable set is countable - an exercise in the text).
2. The real numbers between 0 and 1 can be listed in order r_1, r_2, r_3, \dots .
3. Let the decimal representation of this listing be

$$r_1 = 0.d_{11}d_{12}d_{13}d_{14}d_{15}d_{16} \dots$$

$$r_2 = 0.d_{21}d_{22}d_{23}d_{24}d_{25}d_{26} \dots$$

$$r_3 = 0.d_{31}d_{32}d_{33}d_{34}d_{35}d_{36} \dots$$

$$\vdots$$

4. Form a new real number with the decimal expansion $r = .r_1r_2r_3r_4 \dots$
where $r_i = 3$ if $d_{ii} \neq 3$ and $r_i = 4$ if $d_{ii} = 3$

5. r is not equal to any of the r_1, r_2, r_3, \dots . Because it differs from r_i in its i th position after the decimal point. Therefore there is a real number between 0 and 1 that is not on the list since every real number has a unique decimal expansion. Hence, all the real numbers between 0 and 1 cannot be listed, so the set of real numbers between 0 and 1 is uncountable.
6. Since a set with an uncountable subset is uncountable (an exercise), the set of real numbers is uncountable.






conclusions

$$|Z| = \aleph_0 \quad |N| = \aleph_0 \quad |Q| = \aleph_0$$

$$|R| = \aleph_1 = C$$

- \aleph_0 is the smallest infinite number.
- The cardinality of a set is always less than the cardinality of its power set



Homework

Discrete
Mathematics

- 2.4 ----- -3, 12, 16, 32
- 2.5 ----- 2, 10