

Regression Analysis of Dependent Binary Data for Estimating Disease Etiology from Case-Control Studies ("Small-Area Estimation" of Disease Etiology)

Zhenke Wu

Assistant Professor of Biostatistics
Research Assistant Professor of Michigan Institute for Data Science (MIDAS)
University of Michigan, Ann Arbor



2019 EcoSta, Taichung, Taiwan

Background

In large scale disease etiology studies, an important scientific goal is to assess the effect of explanatory variables upon disease etiology

- Some diseases, e.g., childhood pneumonia, may have multiple potential pathogen causes that infected the lung

Background

In large scale disease etiology studies, an important scientific goal is to assess the effect of explanatory variables upon disease etiology

- Some diseases, e.g., childhood pneumonia, may have multiple potential pathogen causes that infected the lung
- Infeasible to directly observe; “latent”

Background

In large scale disease etiology studies, an important scientific goal is to assess the effect of explanatory variables upon disease etiology

- Some diseases, e.g., childhood pneumonia, may have multiple potential pathogen causes that infected the lung
- Infeasible to directly observe; “latent”
- Scientists measure peripheral sites, e.g., nose, for presence or absence of many pathogens

Background

In large scale disease etiology studies, an important scientific goal is to assess the effect of explanatory variables upon disease etiology

- Some diseases, e.g., childhood pneumonia, may have multiple potential pathogen causes that infected the lung
- Infeasible to directly observe; “latent”
- Scientists measure peripheral sites, e.g., nose, for presence or absence of many pathogens
- However, the tests lack sensitivity or specificity

Background

In large scale disease etiology studies, an important scientific goal is to assess the effect of explanatory variables upon disease etiology

- Some diseases, e.g., childhood pneumonia, may have multiple potential pathogen causes that infected the lung
- Infeasible to directly observe; “latent”
- Scientists measure peripheral sites, e.g., nose, for presence or absence of many pathogens
- However, the tests lack sensitivity or specificity
- Data from healthy **controls** provide requisite information about specificity and covariations; They have no lung infection.

Background

In large scale disease etiology studies, an important scientific goal is to assess the effect of explanatory variables upon disease etiology

- Some diseases, e.g., childhood pneumonia, may have multiple potential pathogen causes that infected the lung
- Infeasible to directly observe; “latent”
- Scientists measure peripheral sites, e.g., nose, for presence or absence of many pathogens
- However, the tests lack sensitivity or specificity
- Data from healthy **controls** provide requisite information about specificity and covariations; They have no lung infection.
- **Statistical problem:** estimate cause-specific case fractions, or “population etiologic fractions” (PEFs)

Background

In large scale disease etiology studies, an important scientific goal is to assess the effect of explanatory variables upon disease etiology

- Some diseases, e.g., childhood pneumonia, may have multiple potential pathogen causes that infected the lung
- Infeasible to directly observe; “latent”
- Scientists measure peripheral sites, e.g., nose, for presence or absence of many pathogens
- However, the tests lack sensitivity or specificity
- Data from healthy **controls** provide requisite information about specificity and covariations; They have no lung infection.
- **Statistical problem:** estimate cause-specific case fractions, or “population etiologic fractions” (PEFs)
- **Motivation for this talk:** PEFs may vary by season, a child’s age, HIV status, disease severity

Data (with Covariates)

- $\mathcal{D} = \{(\mathbf{M}_i, Y_i, \mathbf{X}_i, \mathbf{W}_i), i = 1, \dots, N\}$

Data (with Covariates)

- $\mathcal{D} = \{(\mathbf{M}_i, Y_i, \mathbf{X}_i, Y_i, \mathbf{W}_i), i = 1, \dots, N\}$
- $\mathbf{M}_i = (M_{i1}, \dots, M_{iJ})^\top$: binary measurements; Indicate the presence or absence of J pathogens for subject $i = 1, \dots, N$.

Data (with Covariates)

- $\mathcal{D} = \{(\mathbf{M}_i, Y_i, \mathbf{X}_i, Y_i, \mathbf{W}_i), i = 1, \dots, N\}$
- $\mathbf{M}_i = (M_{i1}, \dots, M_{iJ})^\top$: binary measurements; Indicate the presence or absence of J pathogens for subject $i = 1, \dots, N$.
- Y_i : case (1) or a control (0).

Data (with Covariates)

- $\mathcal{D} = \{(\mathbf{M}_i, Y_i, \mathbf{X}_i, \mathbf{W}_i), i = 1, \dots, N\}$
- $\mathbf{M}_i = (M_{i1}, \dots, M_{iJ})^\top$: binary measurements; Indicate the presence or absence of J pathogens for subject $i = 1, \dots, N$.
- Y_i : case (1) or a control (0).
- $\mathbf{X}_i = (X_{i1}, \dots, X_{ip})^\top$: covariates that may influence case i 's etiologic fractions

Data (with Covariates)

- $\mathcal{D} = \{(\mathbf{M}_i, Y_i, \mathbf{X}_i, Y_i, \mathbf{W}_i), i = 1, \dots, N\}$
- $\mathbf{M}_i = (M_{i1}, \dots, M_{iJ})^\top$: binary measurements; Indicate the presence or absence of J pathogens for subject $i = 1, \dots, N$.
- Y_i : case (1) or a control (0).
- $\mathbf{X}_i = (X_{i1}, \dots, X_{ip})^\top$: covariates that may influence case i 's etiologic fractions
- $\mathbf{W}_i = (W_{i1}, \dots, W_{iq})^\top$: possibly different from \mathbf{X}_i ; may influence control distribution of $[\mathbf{M} \mid Y = 0]$. For example, healthy controls **do not** have disease severity information (which can be included in \mathbf{X}_i).

Data (with Covariates)

- $\mathcal{D} = \{(\mathbf{M}_i, Y_i, \mathbf{X}_i, Y_i, \mathbf{W}_i), i = 1, \dots, N\}$
- $\mathbf{M}_i = (M_{i1}, \dots, M_{iJ})^\top$: binary measurements; Indicate the presence or absence of J pathogens for subject $i = 1, \dots, N$.
- Y_i : case (1) or a control (0).
- $\mathbf{X}_i = (X_{i1}, \dots, X_{ip})^\top$: covariates that may influence case i 's etiologic fractions
- $\mathbf{W}_i = (W_{i1}, \dots, W_{iq})^\top$: possibly different from \mathbf{X}_i ; may influence control distribution of $[\mathbf{M} \mid Y = 0]$. For example, healthy controls **do not** have disease severity information (which can be included in \mathbf{X}_i).
- Continuous covariates: the first p_1 and q_1 elements of \mathbf{X}_i and \mathbf{W}_i , respectively.

Real Data: PERCH Study

Data : 494 cases and 944 controls from one site

Real Data: PERCH Study

Data : 494 cases and 944 controls from one site

Goal a. : Estimate PEFs at all covariate values and assign cause-specific probabilities for each case

Real Data: PERCH Study

Data : 494 cases and 944 controls from one site

Goal a. : Estimate PEFs at all covariate values and assign cause-specific probabilities for each case

Goal b. : It is often of policy interest to quantify the overall cause-specific disease burdens in a population, i.e., the overall PEFs $\pi^* = (\pi_1^*, \dots, \pi_L^*)^\top$ as an empirical average of the stratum-specific PEFs (by \mathbf{X}).

Real Data: PERCH Study

Data : 494 cases and 944 controls from one site

Goal a. : Estimate PEFs at all covariate values and assign cause-specific probabilities for each case

Goal b. : It is often of policy interest to quantify the overall cause-specific disease burdens in a population, i.e., the overall PEFs $\pi^* = (\pi_1^*, \dots, \pi_L^*)^\top$ as an empirical average of the stratum-specific PEFs (by \mathbf{X}).

Model : (to be fitted)

- $J = 7$: noisy presence/absence of 2 bacteria and 5 viruses in nasopharyngeal cavities by polymerase chain reaction (PCR)
- Causes: seven **single-pathogen** causes plus an “Not Specified” (NoS) cause; So $L = J + 1$
- \mathbf{X}_i : enrollment date, age ($<$ or $>$ 1 year), disease severity for cases (severe or very severe), HIV status ($+/-$)
- \mathbf{W}_i : \mathbf{X}_i minus “disease severity”.

PERCH Data: Sparsely-Populated Strata

Table: The observed count (frequency) of cases and controls by age, disease severity and HIV status (1: yes; 0: no). The marginal fractions among cases and controls for each covariate are shown at the bottom. Results from the regression analyses are shown later for the first two strata.

age ≥ 1	very severe (VS) (case-only)	HIV positive	# cases (%) total: 524 (100)	# controls (%) total: 964 (100)
→ 0	0	0	208 (39.7)	545 (56.5)
→ 1	0	0	72 (13.7)	278 (28.8)
0	1	0	116 (22.1)	0
1	1	0	33 (6.3)	0
0	0	1	37 (7.1)	85 (8.8)
1	0	1	24 (4.5)	51 (5.3)
0	1	1	25 (4.8)	0
1	1	1	3 (0.6)	0
case: 25.2%	34.5%	17.0%		
control: 34.3%	-	14.1%		

Current Methods (in each covariate strata)

1. Nested partially-latent class models, npLCM (Wu et al., 2017; Wu et al., 2016)

Current Methods (in each covariate strata)

1. Nested partially-latent class models, npLCM (Wu et al., 2017; Wu et al., 2016)
2. A finite mixture model for multivariate binary data, where the control component is observed, and there are L unobserved case mixture components (e.g., each representing a cause of lung infection)

Current Methods (in each covariate strata)

1. Nested partially-latent class models, npLCM (Wu et al., 2017; Wu et al., 2016)
2. A finite mixture model for multivariate binary data, where the control component is observed, and there are L unobserved case mixture components (e.g., each representing a cause of lung infection)
3. The goal is to estimate the mixing distribution/weights for the L case components (we called them PEFs, $\boldsymbol{\pi} = (\pi_1, \dots, \pi_L)^\top$)

Current Methods (in each covariate strata)

1. Nested partially-latent class models, npLCM (Wu et al., 2017; Wu et al., 2016)
2. A finite mixture model for multivariate binary data, where the control component is observed, and there are L unobserved case mixture components (e.g., each representing a cause of lung infection)
3. The goal is to estimate the mixing distribution/weights for the L case components (we called them PEFs, $\boldsymbol{\pi} = (\pi_1, \dots, \pi_L)^\top$)
4. Fitted in a Bayesian framework; can also estimate the posterior probability of the cause of disease for an **individual** case given her measurements.

Current Methods (in each covariate strata)

1. Nested partially-latent class models, npLCM (Wu et al., 2017; Wu et al., 2016)
2. A finite mixture model for multivariate binary data, where the control component is observed, and there are L unobserved case mixture components (e.g., each representing a cause of lung infection)
3. The goal is to estimate the mixing distribution/weights for the L case components (we called them PEFs, $\boldsymbol{\pi} = (\pi_1, \dots, \pi_L)^\top$)
4. Fitted in a Bayesian framework; can also estimate the posterior probability of the cause of disease for an **individual** case given her measurements.
5. Example of restricted latent class models (RLCM, Xu, 2017, AOS) with one (pLCM) or more (nested pLCM) sets of response probability parameters

Current Methods Fall Short

- *Fully-stratified analysis*: fit an npLCM (more later) to the case-control data in each covariate stratum.

Current Methods Fall Short

- *Fully-stratified analysis*: fit an npLCM (more later) to the case-control data in each covariate stratum.

Like pLCM, the npLCM is **partially-identified** in each stratum, **necessitating** multiple sets of *independent* informative priors across multiple strata.

Two primary issues:

Current Methods Fall Short

- *Fully-stratified analysis*: fit an npLCM (more later) to the case-control data in each covariate stratum.

Like pLCM, the npLCM is **partially-identified** in each stratum, **necessitating** multiple sets of *independent* informative priors across multiple strata.

Two primary issues:

Gap 1a Unstable PEF estimates due to sparsely-populated strata.

Current Methods Fall Short

- *Fully-stratified analysis*: fit an npLCM (more later) to the case-control data in each covariate stratum.

Like pLCM, the npLCM is **partially-identified** in each stratum, **necessitating** multiple sets of *independent* informative priors across multiple strata.

Two primary issues:

Gap 1a Unstable PEF estimates due to sparsely-populated strata.

Gap 1b Informative TPR priors are often elicited for a case population and rarely for each stratum; Reusing independent prior distributions of the TPRs across all the strata will lead to **overly-optimistic** posterior uncertainty in π^* , hampering policy decisions.

This talk

Extend the npLCM to perform regression analysis in case-control disease etiology studies that

This talk

Extend the npLCM to perform regression analysis in case-control disease etiology studies that

(a) incorporates **controls** to estimate the PEFs (π),

This talk

Extend the npLCM to perform regression analysis in case-control disease etiology studies that

- (a) incorporates **controls** to estimate the PEFs (π),
- (b) specifies **parsimonious** functional dependence of π upon covariates such as additivity, and

This talk

Extend the npLCM to perform regression analysis in case-control disease etiology studies that

- (a) incorporates **controls** to estimate the PEFs (π),
- (b) specifies **parsimonious** functional dependence of π upon covariates such as additivity, and
- (c) **correctly assesses the posterior uncertainty** of the PEF functions and the overall PEFs π^* by applying the TPR priors just once.

Quick Review: Nested Partially Latent Class Models (npLCM)

For simplicity, we assume “single-pathogen causes”

npLCM Framework (no Covariates)

Three components of an npLCM likelihood function:

npLCM Framework (no Covariates)

Three components of an npLCM likelihood function:

npLCM Framework (no Covariates)

Three components of an npLCM likelihood function:

a. PEFs/cause-specific case fractions: $\boldsymbol{\pi} = (\pi_1, \dots, \pi_L)^\top =$

$$\{\pi_\ell = \mathbb{P}(I = \ell \mid Y = 1), \ell = 1, \dots, L\} \in \mathcal{S}_{L-1};$$

npLCM Framework (no Covariates)

Three components of an npLCM likelihood function:

- a. PEFs/cause-specific case fractions: $\boldsymbol{\pi} = (\pi_1, \dots, \pi_L)^\top =$

$$\{\pi_\ell = \mathbb{P}(I = \ell \mid Y = 1), \ell = 1, \dots, L\} \in \mathcal{S}_{L-1};$$

- b. $\mathbf{P}_{1\ell} = \{\mathbf{P}_{1\ell}(\mathbf{m})\} = \{\mathbb{P}(\mathbf{M} = \mathbf{m} \mid I = \ell, Y = 1)\}$: a table of probabilities of making J binary observations $\mathbf{M} = \mathbf{m}$ in a case class $\ell \neq 0$;

npLCM Framework (no Covariates)

Three components of an npLCM likelihood function:

- a. PEFs/cause-specific case fractions: $\boldsymbol{\pi} = (\pi_1, \dots, \pi_L)^\top =$

$$\{\pi_\ell = \mathbb{P}(I = \ell \mid Y = 1), \ell = 1, \dots, L\} \in \mathcal{S}_{L-1};$$

- b. $\boldsymbol{P}_{1\ell} = \{\boldsymbol{P}_{1\ell}(\boldsymbol{m})\} = \{\mathbb{P}(\boldsymbol{M} = \boldsymbol{m} \mid I = \ell, Y = 1)\}$: a table of probabilities of making J binary observations $\boldsymbol{M} = \boldsymbol{m}$ in a case class $\ell \neq 0$;
- c. $\boldsymbol{P}_0 = \{\boldsymbol{P}_0(\boldsymbol{m})\} = \{\mathbb{P}(\boldsymbol{M} = \boldsymbol{m} \mid I = 0, Y = 0)\}$: the same probability table as above but for controls.

npLCM Framework (no Covariates)

Three components of an npLCM likelihood function:

- a. PEFs/cause-specific case fractions: $\boldsymbol{\pi} = (\pi_1, \dots, \pi_L)^\top =$

$$\{\pi_\ell = \mathbb{P}(I = \ell \mid Y = 1), \ell = 1, \dots, L\} \in \mathcal{S}_{L-1};$$

- b. $\mathbf{P}_{1\ell} = \{\mathbf{P}_{1\ell}(\mathbf{m})\} = \{\mathbb{P}(\mathbf{M} = \mathbf{m} \mid I = \ell, Y = 1)\}$: a table of probabilities of making J binary observations $\mathbf{M} = \mathbf{m}$ in a case class $\ell \neq 0$;
- c. $\mathbf{P}_0 = \{\mathbf{P}_0(\mathbf{m})\} = \{\mathbb{P}(\mathbf{M} = \mathbf{m} \mid I = 0, Y = 0)\}$: the same probability table as above but for controls.

Cases' disease classes are **unobserved**, so the distribution of their measurements is a weighted finite-mixture model: $\mathbf{P}_1 = \sum_{\ell=1}^L \pi_\ell \mathbf{P}_{1\ell}$

npLCM Framework (no Covariates)

Three components of an npLCM likelihood function:

- a. PEFs/cause-specific case fractions: $\boldsymbol{\pi} = (\pi_1, \dots, \pi_L)^\top =$

$$\{\pi_\ell = \mathbb{P}(I = \ell \mid Y = 1), \ell = 1, \dots, L\} \in \mathcal{S}_{L-1};$$

- b. $\mathbf{P}_{1\ell} = \{\mathbf{P}_{1\ell}(\mathbf{m})\} = \{\mathbb{P}(\mathbf{M} = \mathbf{m} \mid I = \ell, Y = 1)\}$: a table of probabilities of making J binary observations $\mathbf{M} = \mathbf{m}$ in a case class $\ell \neq 0$;
- c. $\mathbf{P}_0 = \{\mathbf{P}_0(\mathbf{m})\} = \{\mathbb{P}(\mathbf{M} = \mathbf{m} \mid I = 0, Y = 0)\}$: the same probability table as above but for controls.

Cases' disease classes are **unobserved**, so the distribution of their measurements is a weighted finite-mixture model: $\mathbf{P}_1 = \sum_{\ell=1}^L \pi_\ell \mathbf{P}_{1\ell}$

The likelihood:

$$L = L_1 \cdot L_0 = \left\{ \prod_{i: Y_i=1} \sum_{\ell=1}^L \pi_\ell \cdot \mathbf{P}_{1\ell}(\mathbf{M}_i; \boldsymbol{\Theta}, \boldsymbol{\Psi}, \boldsymbol{\eta}) \right\} \times \prod_{i': Y_{i'}=0} \mathbf{P}_0(\mathbf{M}_{i'}; \boldsymbol{\Psi}, \boldsymbol{\nu})$$

Special Case: pLCM (Wu et al., 2016)

Control model:

1. $P_0(\mathbf{m}) = \prod_{j=1}^J \{\psi_j\}^{m_j} \{1 - \psi_j\}^{1-m_j} = \Pi(\mathbf{m}; \boldsymbol{\psi})$
 - 1a. $\Pi(\mathbf{m}; \mathbf{s}) = \prod_{j=1}^J \{s_j\}^{m_{ij}} \{1 - s_j\}^{1-m_{ij}}$ is the probability mass function for a product Bernoulli distribution given the success probabilities $\mathbf{s} = (s_1, \dots, s_J)^\top$, $0 \leq s_j \leq 1$
 - 1b. Parameters $\boldsymbol{\psi} = (\psi_1, \dots, \psi_J)^\top$ represent the positive rates absent disease, referred to as “false positive rates” (FPRs).

Local Independence: $M_{ij} \perp M_{ij'} \mid I = 0$

Special Case: pLCM (Wu et al., 2016)

Model for case class $\ell \neq 0$

2. $P_{1\ell}(\mathbf{m})$ is a product of the probabilities of measurements made

Special Case: pLCM (Wu et al., 2016)

Model for **case class** $\ell \neq 0$

2. $\mathbf{P}_{1\ell}(\mathbf{m})$ is a product of the probabilities of measurements made

2a. on the *causative* pathogen ℓ ,

$\mathbb{P}(M_\ell \mid I = \ell, Y = 1, \boldsymbol{\theta}) = \{\theta_\ell\}^{M_\ell} \{1 - \theta_\ell\}^{1-M_\ell}$, where

$\boldsymbol{\theta} = (\theta_1, \dots, \theta_L)^\top$ are “true positive rates” (TPRs), larger than FPRs.

Special Case: pLCM (Wu et al., 2016)

Model for **case class** $\ell \neq 0$

2. $P_{1\ell}(\mathbf{m})$ is a product of the probabilities of measurements made

2a. on the *causative* pathogen ℓ ,

$\mathbb{P}(M_\ell \mid I = \ell, Y = 1, \boldsymbol{\theta}) = \{\theta_\ell\}^{M_\ell} \{1 - \theta_\ell\}^{1-M_\ell}$, where

$\boldsymbol{\theta} = (\theta_1, \dots, \theta_L)^\top$ are “true positive rates” (TPRs), larger than FPRs.

2b. on the *non-causative* pathogens

$\mathbb{P}(\mathbf{M}_{i[-\ell]} \mid I_i = \ell, Y_i = 1, \boldsymbol{\psi}_{[-\ell]}) = \prod(\mathbf{M}_{[-\ell]}; \boldsymbol{\psi}_{[-\ell]})$, where $\mathbf{a}_{[-\ell]}$ represents all but the ℓ -th element in a vector \mathbf{a} .

Special Case: pLCM (Wu et al., 2016)

Model for **case class** $\ell \neq 0$

2. $P_{1\ell}(\mathbf{m})$ is a product of the probabilities of measurements made

2a. on the *causative* pathogen ℓ ,

$\mathbb{P}(M_\ell \mid I = \ell, Y = 1, \boldsymbol{\theta}) = \{\theta_\ell\}^{M_\ell} \{1 - \theta_\ell\}^{1-M_\ell}$, where

$\boldsymbol{\theta} = (\theta_1, \dots, \theta_L)^\top$ are “true positive rates” (TPRs), larger than FPRs.

2b. on the *non-causative* pathogens

$\mathbb{P}(\mathbf{M}_{i[-\ell]} \mid I_i = \ell, Y_i = 1, \boldsymbol{\psi}_{[-\ell]}) = \prod (\mathbf{M}_{[-\ell]}; \boldsymbol{\psi}_{[-\ell]})$, where $\mathbf{a}_{[-\ell]}$ represents all but the ℓ -th element in a vector \mathbf{a} .

2c. Under the single-pathogen-cause assumption, pLCM uses J TPRs $\boldsymbol{\theta}$ for $L = J$ causes and J FPRs $\boldsymbol{\psi}$.

Special Case: pLCM (Wu et al., 2016)

Model for **case class** $\ell \neq 0$

2. $\mathbf{P}_{1\ell}(\mathbf{m})$ is a product of the probabilities of measurements made

2a. on the **causative** pathogen ℓ ,

$\mathbb{P}(M_\ell \mid I = \ell, Y = 1, \boldsymbol{\theta}) = \{\theta_\ell\}^{M_\ell} \{1 - \theta_\ell\}^{1-M_\ell}$, where
 $\boldsymbol{\theta} = (\theta_1, \dots, \theta_L)^\top$ are “true positive rates” (TPRs), larger than FPRs.

2b. on the **non-causative** pathogens

$\mathbb{P}(\mathbf{M}_{i[-\ell]} \mid I_i = \ell, Y_i = 1, \boldsymbol{\psi}_{[-\ell]}) = \prod (\mathbf{M}_{i[-\ell]}; \boldsymbol{\psi}_{[-\ell]})$, where $\mathbf{a}_{[-\ell]}$ represents all but the ℓ -th element in a vector \mathbf{a} .

2c. Under the single-pathogen-cause assumption, pLCM uses J TPRs $\boldsymbol{\theta}$ for $L = J$ causes and J FPRs $\boldsymbol{\psi}$.

2a-2b: **Local Independence (LI)**: $M_{ij} \perp M_{ij'} \mid I = \ell \neq 0$

Special Case: pLCM (Wu et al., 2016)

Model for **case class** $\ell \neq 0$

2. $\mathbf{P}_{1\ell}(\mathbf{m})$ is a product of the probabilities of measurements made

2a. on the **causative** pathogen ℓ ,

$\mathbb{P}(M_\ell \mid I = \ell, Y = 1, \boldsymbol{\theta}) = \{\theta_\ell\}^{M_\ell} \{1 - \theta_\ell\}^{1-M_\ell}$, where $\boldsymbol{\theta} = (\theta_1, \dots, \theta_L)^\top$ are “true positive rates” (TPRs), larger than FPRs.

2b. on the **non-causative** pathogens

$\mathbb{P}(\mathbf{M}_{i[-\ell]} \mid I_i = \ell, Y_i = 1, \boldsymbol{\psi}_{[-\ell]}) = \prod(\mathbf{M}_{i[-\ell]}; \boldsymbol{\psi}_{[-\ell]})$, where $\mathbf{a}_{[-\ell]}$ represents all but the ℓ -th element in a vector \mathbf{a} .

2c. Under the single-pathogen-cause assumption, pLCM uses J TPRs $\boldsymbol{\theta}$ for $L = J$ causes and J FPRs $\boldsymbol{\psi}$.

2a-2b: **Local Independence (LI)**: $M_{ij} \perp M_{ij'} \mid I = \ell \neq 0$

2a-2b. **Non-interference**: disease-causing pathogen(s) are more frequently detected among cases than controls ($\theta_\ell > \psi_\ell$) and the non-causative pathogens are observed with the same rates among cases as in controls

“nested” pLCM

Relax the LI and Non-interference Assumption

- Direct evidence: control measurements $(M_{i1}, \dots, M_{iJ})'$

“nested” pLCM

Relax the LI and Non-interference Assumption

- **Direct evidence:** control measurements $(M_{i1}, \dots, M_{iJ})'$
 - test cross-reactions (prevented in PERCH assays)
 - lab technicians effect
 - heterogeneity in subjects' immunity level

“nested” pLCM

Relax the LI and Non-interference Assumption

- **Direct evidence:** control measurements $(M_{i1}, \dots, M_{iJ})'$
 - test cross-reactions (prevented in PERCH assays)
 - lab technicians effect
 - heterogeneity in subjects' immunity level
- **Deviations from independence impacts inference** (Cf. Pepe and Janes, 2007, *Biostatistics*; Albert et al., 2001, *Biometrics*)

“nested” pLCM

Relax the LI and Non-interference Assumption

- **Direct evidence:** control measurements $(M_{i1}, \dots, M_{iJ})'$
 - test cross-reactions (prevented in PERCH assays)
 - lab technicians effect
 - heterogeneity in subjects' immunity level
- **Deviations from independence impacts inference** (Cf. Pepe and Janes, 2007, *Biostatistics*; Albert et al., 2001, *Biometrics*)
- **Modeling Deviation from LI** Modeling a cross-classified probability contingency table

$$\mathbb{P}[M_{i1} = m_1, \dots, M_{iJ} = m_J], \forall \mathbf{m} = (m_1, \dots, m_J)'$$

“nested” pLCM

Relax the LI and Non-interference Assumption

- **Direct evidence:** control measurements $(M_{i1}, \dots, M_{iJ})'$
 - test cross-reactions (prevented in PERCH assays)
 - lab technicians effect
 - heterogeneity in subjects' immunity level
- **Deviations from independence impacts inference** (Cf. Pepe and Janes, 2007, *Biostatistics*; Albert et al., 2001, *Biometrics*)
- **Modeling Deviation from LI** Modeling a cross-classified probability contingency table

$$\mathbb{P}[M_{i1} = m_1, \dots, M_{iJ} = m_J], \forall \mathbf{m} = (m_1, \dots, m_J)'$$

- Log-linear parametrization

“nested” pLCM

Relax the LI and Non-interference Assumption

- **Direct evidence:** control measurements $(M_{i1}, \dots, M_{iJ})'$
 - test cross-reactions (prevented in PERCH assays)
 - lab technicians effect
 - heterogeneity in subjects' immunity level
- **Deviations from independence impacts inference** (Cf. Pepe and Janes, 2007, *Biostatistics*; Albert et al., 2001, *Biometrics*)
- **Modeling Deviation from LI** Modeling a cross-classified probability contingency table

$$\mathbb{P}[M_{i1} = m_1, \dots, M_{iJ} = m_J], \forall \mathbf{m} = (m_1, \dots, m_J)'$$

- Log-linear parametrization
- Generalized linear mixed-effect models (GLMM)

“nested” pLCM

Relax the LI and Non-interference Assumption

- **Direct evidence:** control measurements $(M_{i1}, \dots, M_{iJ})'$
 - test cross-reactions (prevented in PERCH assays)
 - lab technicians effect
 - heterogeneity in subjects' immunity level
- **Deviations from independence impacts inference** (Cf. Pepe and Janes, 2007, *Biostatistics*; Albert et al., 2001, *Biometrics*)
- **Modeling Deviation from LI** Modeling a cross-classified probability contingency table

$$\mathbb{P}[M_{i1} = m_1, \dots, M_{iJ} = m_J], \forall \mathbf{m} = (m_1, \dots, m_J)'$$

- Log-linear parametrization
- Generalized linear mixed-effect models (GLMM)
- Simplex factor model; similar to mixed-membership model (Cf. Bhattacharya and Dunson, 2012, *JASA*)

“nested” pLCM

Relax the LI and Non-interference Assumption


- **Direct evidence:** control measurements $(M_{i1}, \dots, M_{iJ})'$
 - test cross-reactions (prevented in PERCH assays)
 - lab technicians effect
 - heterogeneity in subjects' immunity level
- **Deviations from independence impacts inference** (Cf. Pepe and Janes, 2007, *Biostatistics*; Albert et al., 2001, *Biometrics*)
- **Modeling Deviation from LI** Modeling a cross-classified probability contingency table

$$\mathbb{P}[M_{i1} = m_1, \dots, M_{iJ} = m_J], \forall \mathbf{m} = (m_1, \dots, m_J)'$$

- Log-linear parametrization
- Generalized linear mixed-effect models (GLMM)
- Simplex factor model; similar to mixed-membership model (Cf. Bhattacharya and Dunson, 2012, *JASA*)
- **PARAFAC decomposition** (Cf. Dunson and Xing, 2009, *JASA*)


Nested Partially-Latent Class Models (npLCM; Wu and Zeger, 2016)

Example: 5 Pathogens, 2 Subclasses; BrS Data Only



Nested Partially-Latent Class Models (npLCM; Wu and Zeger, 2016)



Example: 5 Pathogens, 3 Subclasses; BrS Data Only



Encourage Few Subclasses: Stick-Breaking Prior

$$V_j \sim \text{Beta}(1, \alpha); \text{ Example: } K = 10, \alpha = 1$$

Length = 1



- On average, the first several segments receive most weights

npLCM: Likelihood and Prior

BrS Data Only

- Likelihood

$$P_0(\mathbf{M}_i = \mathbf{m}) = \sum_{k=1}^K \nu_k \prod_{j=1}^J \left\{ \psi_k^{(j)} \right\}^{m_j} \left\{ 1 - \psi_k^{(j)} \right\}^{1-m_j},$$

$$P_1(\mathbf{M}_i = \mathbf{m}) = \sum_{j=1}^J \pi_j \sum_{k=1}^K \left[\eta_k \left\{ \theta_k^{(j)} \right\}^{m_j} \left\{ 1 - \theta_k^{(j)} \right\}^{1-m_j} \prod_{\ell \neq j} \left\{ \psi_k^{(\ell)} \right\}^{m_\ell} \left\{ 1 - \psi_k^{(\ell)} \right\}^{1-m_\ell} \right],$$

- Prior:

$$\boldsymbol{\pi} \sim \text{Dirichlet}(.5, \dots, .5),$$

$$\psi_k^{(j)} \sim \text{Beta}(1, 1), \quad \theta_k \sim \text{Beta}(c_{1kj}, c_{2kj}), j = 1, \dots, J; k = 1, \dots, \infty,$$

$$Z_{i'} \mid I_{i'}^L = j \sim \sum_{k=1}^{\infty} U_k \prod_{\ell < k} [1 - U_\ell] \delta_k, \quad U_k \sim \text{Beta}(1, \alpha_0), \text{ for all cases,}$$

$$Z_i \sim \sum_{k=1}^{\infty} V_k \prod_{\ell < k} [1 - V_\ell] \delta_k, \quad V_k \sim \text{Beta}(1, \alpha_0), \text{ for all controls,}$$

$$\alpha_0 \sim \text{Gamma}(0.25, 0.25),$$

Regression Extension for P_0
and P_1 :

letting π_ℓ, ν_k, η_k depend on
covariates

Roadmap

Let three sets of parameters in an npLCM (pg.17) depend on the observed covariates

- 1x. Etiology regression function among cases, $\{\pi_\ell(\mathbf{x}), \ell \neq 0\}$, which is of primary scientific interest
- 2x. Conditional probability of measurements \mathbf{m} given covariates \mathbf{w} in controls: $P_0(\mathbf{m}; \mathbf{w}) = [\mathbf{M} = \mathbf{m} \mid \mathbf{W} = \mathbf{w}, l = 0]$,
- 3x. 2x above, but in the case class ℓ :
 $P_{1\ell}(\mathbf{m}; \mathbf{w}) = [\mathbf{M} = \mathbf{m} \mid \mathbf{W} = \mathbf{w}, l = \ell], \ell = 1, \dots, L$

note Keep the specifications for the TPRs and FPRs (Θ, Ψ) as in the original npLCM.

Etiology Regression $\pi_\ell(\mathbf{X})$

$\pi_\ell(\mathbf{X})$ is the primary target of inference.

Etiology Regression $\pi_\ell(\mathbf{X})$

$\pi_\ell(\mathbf{X})$ is the primary target of inference.

1. Recall that $I_i = \ell$ represents case i 's disease being caused by pathogen ℓ .

Etiology Regression $\pi_\ell(\mathbf{X})$

$\pi_\ell(\mathbf{X})$ is the primary target of inference.

1. Recall that $I_i = \ell$ represents case i 's disease being caused by pathogen ℓ .
2. Occurs with probability $\pi_{i\ell}$ that depends upon covariates.

Etiology Regression $\pi_\ell(\mathbf{X})$

$\pi_\ell(\mathbf{X})$ is the primary target of inference.

1. Recall that $I_i = \ell$ represents case i 's disease being caused by pathogen ℓ .
2. Occurs with probability $\pi_{i\ell}$ that depends upon covariates.

3. Over-parameterized multinomial logistic regression:

$\pi_{i\ell} = \pi_\ell(\mathbf{X}_i) = \exp\{\phi_\ell(\mathbf{X}_i)\} / \sum_{\ell'=1}^L \exp\{\phi_{\ell'}(\mathbf{X}_i)\}$, $\ell = 1, \dots, L$,
where $\phi_\ell(\mathbf{X}_i) - \phi_L(\mathbf{X}_i)$ is the log odds of case i in disease class ℓ relative to L : $\log \pi_{i\ell} / \pi_{iL}$.

Etiology Regression $\pi_\ell(\mathbf{X})$

$\pi_\ell(\mathbf{X})$ is the primary target of inference.

1. Recall that $I_i = \ell$ represents case i 's disease being caused by pathogen ℓ .
2. Occurs with probability $\pi_{i\ell}$ that depends upon covariates.
3. Over-parameterized multinomial logistic regression:
 $\pi_{i\ell} = \pi_\ell(\mathbf{X}_i) = \exp\{\phi_\ell(\mathbf{X}_i)\} / \sum_{\ell'=1}^L \exp\{\phi_{\ell'}(\mathbf{X}_i)\}$, $\ell = 1, \dots, L$,
where $\phi_\ell(\mathbf{X}_i) - \phi_L(\mathbf{X}_i)$ is the log odds of case i in disease class ℓ relative to L : $\log \pi_{i\ell} / \pi_{iL}$.
4. Without specifying a baseline category, we treat all the disease classes symmetrically which simplifies prior specification.

Etiology Regression $\pi_\ell(\mathbf{X})$

$\pi_\ell(\mathbf{X})$ is the primary target of inference.

1. Recall that $I_i = \ell$ represents case i 's disease being caused by pathogen ℓ .
2. Occurs with probability $\pi_{i\ell}$ that depends upon covariates.
3. Over-parameterized multinomial logistic regression:
$$\pi_{i\ell} = \pi_\ell(\mathbf{X}_i) = \exp\{\phi_\ell(\mathbf{X}_i)\} / \sum_{\ell'=1}^L \exp\{\phi_{\ell'}(\mathbf{X}_i)\}, \ell = 1, \dots, L,$$
where $\phi_\ell(\mathbf{X}_i) - \phi_L(\mathbf{X}_i)$ is the log odds of case i in disease class ℓ relative to L : $\log \pi_{i\ell} / \pi_{iL}$.
4. Without specifying a baseline category, we treat all the disease classes symmetrically which simplifies prior specification.
5. Additive models for $\phi_\ell(\mathbf{x}; \mathbf{\Gamma}_\ell^\pi) = \sum_{j=1}^{p_1} f_{\ell j}^\pi(x_j; \beta_{\ell j}^\pi) + \tilde{\mathbf{x}}^\top \boldsymbol{\gamma}_\ell^\pi$

Etiology Regression $\pi_\ell(\mathbf{X})$

$\pi_\ell(\mathbf{X})$ is the primary target of inference.

1. Recall that $I_i = \ell$ represents case i 's disease being caused by pathogen ℓ .
2. Occurs with probability $\pi_{i\ell}$ that depends upon covariates.
3. Over-parameterized multinomial logistic regression:
$$\pi_{i\ell} = \pi_\ell(\mathbf{X}_i) = \exp\{\phi_\ell(\mathbf{X}_i)\} / \sum_{\ell'=1}^L \exp\{\phi_{\ell'}(\mathbf{X}_i)\}, \ell = 1, \dots, L,$$
where $\phi_\ell(\mathbf{X}_i) - \phi_L(\mathbf{X}_i)$ is the log odds of case i in disease class ℓ relative to L : $\log \pi_{i\ell} / \pi_{iL}$.
4. Without specifying a baseline category, we treat all the disease classes symmetrically which simplifies prior specification.
5. Additive models for $\phi_\ell(\mathbf{x}; \mathbf{\Gamma}_\ell^\pi) = \sum_{j=1}^{P_1} f_{\ell j}^\pi(x_j; \beta_{\ell j}^\pi) + \tilde{\mathbf{x}}^\top \boldsymbol{\gamma}_\ell^\pi$
- 5a. Use B-spline basis expansion to approximate $f_{\ell j}^\pi(\cdot)$ and use P-spline for estimating smooth functions.
- 5b. $\tilde{\mathbf{x}}$ is the subvector of the predictors \mathbf{x} ; $\mathbf{\Gamma}_\ell^\pi = (\beta_{\ell j}^\pi, \boldsymbol{\gamma}_\ell^\pi)$.

P_0 : Multivariate binary regression for controls

Desirable properties

Model Specification:

- Model space large enough for complex conditional dependence of \mathbf{M} given covariates \mathbf{W}
- Upward compatibility, or reproducibility (**invariant parameter interpretation** with increasing dimensions or complex patterns of missing responses)

Estimation:

- Consistency and Efficiency
- **Adaptivity: regularization to adapt to the difficulty of the problem**, e.g., model residual dependence $[\mathbf{M} \mid \mathbf{W}, I = 0]$ only if necessary; model the effect of covariates only if necessary

Regression Specifications: For Controls

- The pmf for controls' measurements:

$$Pr(\mathbf{M}_i = \mathbf{m} \mid \mathbf{W}_i, l_i = 0) = \sum_{k=1}^K \nu_k(\mathbf{W}_i) \Pi(\mathbf{m}; \Psi_k),$$

$$\Psi_k = (\psi_k^{(1)}, \dots, \psi_k^{(J)})'$$

Regression Specifications: For Controls

- The pmf for controls' measurements:

$$Pr(\mathbf{M}_i = \mathbf{m} \mid \mathbf{W}_i, l_i = 0) = \sum_{k=1}^K \nu_k(\mathbf{W}_i) \Pi(\mathbf{m}; \Psi_k),$$

$$\Psi_k = (\psi_k^{(1)}, \dots, \psi_k^{(J)})'$$

- The vector $(\nu_1(\mathbf{W}_i), \dots, \nu_K(\mathbf{W}_i))$ lies in a $(K - 1)$ -simplex

Regression Specifications: For Controls

- The pmf for controls' measurements:

$$Pr(\mathbf{M}_i = \mathbf{m} \mid \mathbf{W}_i, l_i = 0) = \sum_{k=1}^K \nu_k(\mathbf{W}_i) \Pi(\mathbf{m}; \Psi_k),$$

$$\Psi_k = (\psi_k^{(1)}, \dots, \psi_k^{(J)})'$$

- The vector $(\nu_1(\mathbf{W}_i), \dots, \nu_K(\mathbf{W}_i))$ lies in a $(K - 1)$ -simplex
- $\Pi(\mathbf{m}; s) = \prod_{j=1}^J \{s_j\}^{m_{ij}} (1 - s_j)^{1 - m_{ij}}$

Regression Specifications: For Controls

- The pmf for controls' measurements:
$$Pr(\mathbf{M}_i = \mathbf{m} \mid \mathbf{W}_i, l_i = 0) = \sum_{k=1}^K \nu_k(\mathbf{W}_i) \Pi(\mathbf{m}; \Psi_k),$$
$$\Psi_k = (\psi_k^{(1)}, \dots, \psi_k^{(J)})'$$
 - The vector $(\nu_1(\mathbf{W}_i), \dots, \nu_K(\mathbf{W}_i))$ lies in a $(K - 1)$ -simplex
 - $\Pi(\mathbf{m}; s) = \prod_{j=1}^J \{s_j\}^{m_{ij}} (1 - s_j)^{1 - m_{ij}}$
- An equivalent generative process:

Regression Specifications: For Controls

- The pmf for controls' measurements:

$$Pr(\mathbf{M}_i = \mathbf{m} \mid \mathbf{W}_i, l_i = 0) = \sum_{k=1}^K \nu_k(\mathbf{W}_i) \Pi(\mathbf{m}; \Psi_k),$$

$$\Psi_k = (\psi_k^{(1)}, \dots, \psi_k^{(J)})'$$

- The vector $(\nu_1(\mathbf{W}_i), \dots, \nu_K(\mathbf{W}_i))$ lies in a $(K - 1)$ -simplex
- $\Pi(\mathbf{m}; s) = \prod_{j=1}^J \{s_j\}^{m_{ij}} (1 - s_j)^{1-m_{ij}}$

- An equivalent generative process:

sample subclass indicator : $Z_i \mid \mathbf{W}_i \sim \text{Categorical}_K(\boldsymbol{\nu}(\mathbf{W}_i))$

generate measurements : $M_{ij} \mid Z_i = k \sim \text{Bernoulli}(\psi_k^{(j)}),$
independently for $j = 1, \dots, J.$

Regression Specifications: Covariate Dependent Weights for Controls

Stick-breaking parametrization of weight functions

$\nu_k(\mathbf{W}_i) = P(Z_i = k \mid \mathbf{W}_i)$ by

$$h_k(\mathbf{W}_i; \Gamma_k^\nu) = \begin{cases} g(\alpha_{ik}^\nu) \prod_{s < k} \{1 - g(\alpha_{is}^\nu)\}, & \text{if } k < K, \\ \prod_{s < k} \{1 - g(\alpha_{is}^\nu)\}, & \text{if } k = K, \end{cases}$$

Regression Specifications: Covariate Dependent Weights for Controls

Stick-breaking parametrization of weight functions

$\nu_k(\mathbf{W}_i) = P(Z_i = k \mid \mathbf{W}_i)$ by

$$h_k(\mathbf{W}_i; \Gamma_k^\nu) = \begin{cases} g(\alpha_{ik}^\nu) \prod_{s < k} \{1 - g(\alpha_{is}^\nu)\}, & \text{if } k < K, \\ \prod_{s < k} \{1 - g(\alpha_{is}^\nu)\}, & \text{if } k = K, \end{cases}$$

We specify α_{ik}^ν via additive models at g^{-1} scale ($g(\cdot) = 1/(1 + \exp\{\cdot\})$):

$$\alpha_{ik}^\nu = \mu_{k0} + \sum_{j=1}^{q_1} f_{kj}(\mathbf{W}_{ij}; \beta_{kj}^\nu) + \widetilde{\mathbf{W}}_i^\top \gamma_k^\nu, \quad k = 1, \dots, K-1.$$

Regression Specifications: Covariate Dependent Weights for Controls

Stick-breaking parametrization of weight functions

$\nu_k(\mathbf{W}_i) = P(Z_i = k \mid \mathbf{W}_i)$ by

$$h_k(\mathbf{W}_i; \Gamma_k^\nu) = \begin{cases} g(\alpha_{ik}^\nu) \prod_{s < k} \{1 - g(\alpha_{is}^\nu)\}, & \text{if } k < K, \\ \prod_{s < k} \{1 - g(\alpha_{is}^\nu)\}, & \text{if } k = K, \end{cases}$$

We specify α_{ik}^ν via additive models at g^{-1} scale ($g(\cdot) = 1/(1 + \exp\{\cdot\})$):

$$\alpha_{ik}^\nu = \mu_{k0} + \sum_{j=1}^{q_1} f_{kj}(\mathbf{W}_{ij}; \beta_{kj}^\nu) + \widetilde{\mathbf{W}}_i^\top \gamma_k^\nu, \quad k = 1, \dots, K-1.$$

Expanded the smooth functions by B-spline bases with coefficients

β_{kj}^ν ; $\widetilde{\mathbf{w}}$ is a subvector of covariates \mathbf{w}

Adaptivity Considerations

Proposed Model

- Prevent overfitting when the regression is easy, and improve interpretability
- We *a priori* place substantial probabilities on models with the following two features:
 - a) Smooth weight regression curves $\nu_k(\cdot)$ in by Bayesian Penalized-Splines (P-Splines) combined with mixture priors on spline coefficients to sensitively distinguish constant $\alpha_k^\nu(\cdot)$ from flexible smooth curves, and,
 - b) Few subclasses with effective weights (in the sense that $\nu_k(\cdot)$ is bounded away from 0 and 1) using a novel additive half-Cauchy prior for μ_{k0} .

On Consideration b) Selective Stopping, or “Uniform Shrinkage over Simplex”

Proposed Model

On Consideration b) Selective Stopping, or “Uniform Shrinkage over Simplex”

Proposed Model

- We let $\mu_{k0} = \sum_{j=1}^k u_{kj} \mu_{k0}^*$ where $u_{kj}, 1 \leq j \leq k$ are pre-specified triangular array of positive values.

On Consideration b) Selective Stopping, or “Uniform Shrinkage over Simplex”

Proposed Model

- We let $\mu_{k0} = \sum_{j=1}^k u_{kj} \mu_{k0}^*$ where $u_{kj}, 1 \leq j \leq k$ are pre-specified triangular array of positive values.
- $u_{kj} = 1, j = 1, \dots, k$; other choices: $u_{kj} = \mathbb{I}\{k = j\}$ or $u_{kj} = 1/k$ may be useful in other settings.

On Consideration b) Selective Stopping, or “Uniform Shrinkage over Simplex”

Proposed Model

- We let $\mu_{k0} = \sum_{j=1}^k u_{kj} \mu_{k0}^*$ where $u_{kj}, 1 \leq j \leq k$ are pre-specified triangular array of positive values.
- $u_{kj} = 1, j = 1, \dots, k$; other choices: $u_{kj} = \mathbb{I}\{k = j\}$ or $u_{kj} = 1/k$ may be useful in other settings.
- We specify the prior distributions for μ_{k0}^* to be heavy-tailed:

$$\mu_{k0}^* \sim \text{Cauchy}^+(0, s_k), \quad k = 1, \dots, K,$$

On Consideration b) Selective Stopping, or “Uniform Shrinkage over Simplex”

Proposed Model

- We let $\mu_{k0} = \sum_{j=1}^k u_{kj} \mu_{k0}^*$ where $u_{kj}, 1 \leq j \leq k$ are pre-specified triangular array of positive values.
- $u_{kj} = 1, j = 1, \dots, k$; other choices: $u_{kj} = \mathbb{I}\{k = j\}$ or $u_{kj} = 1/k$ may be useful in other settings.
- We specify the prior distributions for μ_{k0}^* to be heavy-tailed:

$$\mu_{k0}^* \sim \text{Cauchy}^+(0, s_k), \quad k = 1, \dots, K,$$

- Large values of s_k produce large μ_{k0}^* and help stop stick-breaking at class k , while small values let the stick-breaking continue to Step $k + 1$.

On Consideration b) Selective Stopping, or “Uniform Shrinkage over Simplex”

Proposed Model


- We let $\mu_{k0} = \sum_{j=1}^k u_{kj} \mu_{k0}^*$ where $u_{kj}, 1 \leq j \leq k$ are pre-specified triangular array of positive values.
- $u_{kj} = 1, j = 1, \dots, k$; other choices: $u_{kj} = \mathbb{I}\{k = j\}$ or $u_{kj} = 1/k$ may be useful in other settings.
- We specify the prior distributions for μ_{k0}^* to be heavy-tailed:

$$\mu_{k0}^* \sim \text{Cauchy}^+(0, s_k), \quad k = 1, \dots, K,$$

- Large values of s_k produce large μ_{k0}^* and help stop stick-breaking at class k , while small values let the stick-breaking continue to Step $k + 1$.
- Encourages using a small number of classes (K) to approximate the observed 2^J probability contingency table in finite samples.

Estimation Performance

Simulation: With Covariate



X-axis: covariate values

Y-axis: weight; 0 to 1.

Subclass Weight Regression: For Cases

Subclass Weight Regression: For Cases

The pmf for cases' measurements:

$$Pr(\mathbf{M}_i = \mathbf{m}) = \sum_{\ell=1}^L \pi_{i\ell} \sum_{k=1}^K \eta_{ik} \Pi(\mathbf{M}_i; \mathbf{p}_{k\ell})$$

Subclass Weight Regression: For Cases

The pmf for cases' measurements:

$$Pr(\mathbf{M}_i = \mathbf{m}) = \sum_{\ell=1}^L \pi_{i\ell} \sum_{k=1}^K \eta_{ik} \Pi(\mathbf{M}_i; \mathbf{p}_{k\ell})$$

- $\mathbf{p}_{k\ell} = \{p_{k\ell}^{(j)}, j = 1, \dots, J\}$ are positive rates for J measurements in subclass k of disease class ℓ :

$$p_{k\ell}^{(j)} = \left\{ \theta_k^{(j)} \right\}^{\mathbb{I}\{j=\ell\}} \cdot \left\{ \psi_k^{(j)} \right\}^{1-\mathbb{I}\{j=\ell\}}$$

Subclass Weight Regression: For Cases

The pmf for cases' measurements:

$$Pr(\mathbf{M}_i = \mathbf{m}) = \sum_{\ell=1}^L \pi_{i\ell} \sum_{k=1}^K \eta_{ik} \Pi(\mathbf{M}_i; \mathbf{p}_{k\ell})$$

- $\mathbf{p}_{k\ell} = \{p_{k\ell}^{(j)}, j = 1, \dots, J\}$ are positive rates for J measurements in subclass k of disease class ℓ :

$$p_{k\ell}^{(j)} = \left\{ \theta_k^{(j)} \right\}^{\mathbb{I}\{j=\ell\}} \cdot \left\{ \psi_k^{(j)} \right\}^{1-\mathbb{I}\{j=\ell\}}$$
- Equals the TPR $\theta_k^{(j)}$ for a causative pathogen and the FPR $\psi_k^{(j)}$ otherwise

Subclass Weight Regression: For Cases

The pmf for cases' measurements:

$$Pr(\mathbf{M}_i = \mathbf{m}) = \sum_{\ell=1}^L \pi_{i\ell} \sum_{k=1}^K \eta_{ik} \Pi(\mathbf{M}_i; \mathbf{p}_{k\ell})$$

- $\mathbf{p}_{k\ell} = \{p_{k\ell}^{(j)}, j = 1, \dots, J\}$ are positive rates for J measurements in subclass k of disease class ℓ :

$$p_{k\ell}^{(j)} = \left\{ \theta_k^{(j)} \right\}^{\mathbb{I}\{j=\ell\}} \cdot \left\{ \psi_k^{(j)} \right\}^{1-\mathbb{I}\{j=\ell\}}$$

- Equals the TPR $\theta_k^{(j)}$ for a causative pathogen and the FPR $\psi_k^{(j)}$ otherwise
- Subclass weight regression $\eta_k(\mathbf{W})$ is also specified via stick-breaking: $\eta_{ik} = h_k(\mathbf{W}_i; \mathbf{\Gamma}_k^\eta)$, $k = 1, \dots, K - 1$

Subclass Weight Regression: For Cases

The pmf for cases' measurements:

$$Pr(\mathbf{M}_i = \mathbf{m}) = \sum_{\ell=1}^L \pi_{i\ell} \sum_{k=1}^K \eta_{ik} \Pi(\mathbf{M}_i; \mathbf{p}_{k\ell})$$

- $\mathbf{p}_{k\ell} = \{p_{k\ell}^{(j)}, j = 1, \dots, J\}$ are positive rates for J measurements in subclass k of disease class ℓ :

$$p_{k\ell}^{(j)} = \left\{ \theta_k^{(j)} \right\}^{\mathbb{I}\{j=\ell\}} \cdot \left\{ \psi_k^{(j)} \right\}^{1-\mathbb{I}\{j=\ell\}}$$
- Equals the TPR $\theta_k^{(j)}$ for a causative pathogen and the FPR $\psi_k^{(j)}$ otherwise
- Subclass weight regression $\eta_k(\mathbf{W})$ is also specified via stick-breaking: $\eta_{ik} = h_k(\mathbf{W}_i; \mathbf{\Gamma}_k^{\eta})$, $k = 1, \dots, K - 1$
- α_{ik}^{η} : GAMs

Subclass Weight Regression: For Cases

The pmf for cases' measurements:

$$Pr(\mathbf{M}_i = \mathbf{m}) = \sum_{\ell=1}^L \pi_{i\ell} \sum_{k=1}^K \eta_{ik} \Pi(\mathbf{M}_i; \mathbf{p}_{k\ell})$$

- $\mathbf{p}_{k\ell} = \{p_{k\ell}^{(j)}, j = 1, \dots, J\}$ are positive rates for J measurements in subclass k of disease class ℓ :

$$p_{k\ell}^{(j)} = \left\{ \theta_k^{(j)} \right\}^{\mathbb{I}\{j=\ell\}} \cdot \left\{ \psi_k^{(j)} \right\}^{1-\mathbb{I}\{j=\ell\}}$$
- Equals the TPR $\theta_k^{(j)}$ for a causative pathogen and the FPR $\psi_k^{(j)}$ otherwise
- Subclass weight regression $\eta_k(\mathbf{W})$ is also specified via stick-breaking: $\eta_{ik} = h_k(\mathbf{W}_i; \mathbf{\Gamma}_k^\eta)$, $k = 1, \dots, K - 1$
- α_{ik}^η : GAMs
- $\alpha_{ik}^\eta = \alpha_k^\eta(\mathbf{W}_i; \mathbf{\Gamma}_k^\eta) = \mu_{k0} + \sum_{j=1}^{q_1} f_{kj}(W_{ij}; \beta_{kj}^\eta) + \widetilde{\mathbf{W}}_i^\top \boldsymbol{\gamma}_k^\eta$, where $\mathbf{\Gamma}_k^\eta = \{\mu_{k0}, \{\beta_{kj}^\eta\}, \boldsymbol{\gamma}_k^\eta\}$ are the regression parameters.

Subclass Weight Regression: For Cases

The pmf for cases' measurements:

$$Pr(\mathbf{M}_i = \mathbf{m}) = \sum_{\ell=1}^L \pi_{i\ell} \sum_{k=1}^K \eta_{ik} \Pi(\mathbf{M}_i; \mathbf{p}_{k\ell})$$

- $\mathbf{p}_{k\ell} = \{p_{k\ell}^{(j)}, j = 1, \dots, J\}$ are positive rates for J measurements in subclass k of disease class ℓ :

$$p_{k\ell}^{(j)} = \left\{ \theta_k^{(j)} \right\}^{\mathbb{I}\{j=\ell\}} \cdot \left\{ \psi_k^{(j)} \right\}^{1-\mathbb{I}\{j=\ell\}}$$
- Equals the TPR $\theta_k^{(j)}$ for a causative pathogen and the FPR $\psi_k^{(j)}$ otherwise
- Subclass weight regression $\eta_k(\mathbf{W})$ is also specified via stick-breaking: $\eta_{ik} = h_k(\mathbf{W}_i; \mathbf{\Gamma}_k^\eta)$, $k = 1, \dots, K - 1$
- α_{ik}^η : GAMs
- $\alpha_{ik}^\eta = \alpha_k^\eta(\mathbf{W}_i; \mathbf{\Gamma}_k^\eta) = \mu_{k0} + \sum_{j=1}^{q_1} f_{kj}(W_{ij}; \beta_{kj}^\eta) + \widetilde{\mathbf{W}}_i^\top \boldsymbol{\gamma}_k^\eta$, where $\mathbf{\Gamma}_k^\eta = \{\mu_{k0}, \{\beta_{kj}^\eta\}, \boldsymbol{\gamma}_k^\eta\}$ are the regression parameters.
- we use μ_{k0} from the control regression to ensure important subclasses in the cases

npLCM Regression Framework

The npLCM regression framework is then obtained as:

npLCM Regression Framework

The npLCM regression framework is then obtained as:

- Control likelihood with covariates:

$$L_0^{\text{reg}} = \prod_{i: Y_i=0} \sum_{k=1}^K h_k(\mathbf{W}_i; \Gamma_k^\nu) \Pi(\mathbf{M}_i; \Psi_k).$$

npLCM Regression Framework

The npLCM regression framework is then obtained as:

- Control likelihood with covariates:

$$L_0^{\text{reg}} = \prod_{i: Y_i=0} \sum_{k=1}^K h_k(\mathbf{W}_i; \mathbf{\Gamma}_k^\nu) \Pi(\mathbf{M}_i; \mathbf{\Psi}_k).$$

- Cases likelihood with covariates:

$$L_1^{\text{reg}} = \prod_{i: Y_i=1} \left\{ \sum_{\ell=1}^L \left[\pi_{\ell}(\mathbf{X}_i; \mathbf{\Gamma}_{\ell}^{\pi}) \sum_{k=1}^K \{ \eta_{ik} \cdot \Pi(\mathbf{M}_i; \mathbf{p}_{k\ell}) \} \right] \right\}, \quad (2)$$

npLCM Regression Framework

The npLCM regression framework is then obtained as:

- Control likelihood with covariates:

$$L_0^{\text{reg}} = \prod_{i: Y_i=0} \sum_{k=1}^K h_k(\mathbf{W}_i; \Gamma_k^\nu) \Pi(\mathbf{M}_i; \Psi_k).$$

- Cases likelihood with covariates:

$$L_1^{\text{reg}} = \prod_{i: Y_i=1} \left\{ \sum_{\ell=1}^L \left[\pi_{\ell}(\mathbf{X}_i; \Gamma_{\ell}^{\pi}) \sum_{k=1}^K \{ \eta_{ik} \cdot \Pi(\mathbf{M}_i; \mathbf{p}_{k\ell}) \} \right] \right\}, \quad (2)$$

The joint likelihood for the regression model can be written as:

$$L^{\text{reg}} = L_1^{\text{reg}} \times L_0^{\text{reg}}.$$

Prior Specifications

Unknown parameters:

- etiology regression coefficients ($\{\Gamma_\ell^\pi\}$),
- subclass mixing weight parameters for cases ($\{\Gamma_k^\eta\}$) and controls ($\{\Gamma_k^\nu\}$),
- true and false positive rates ($\Theta = \{\theta_k^{(j)}\}$, $\Psi = \{\psi_k^{(j)}\}$).

Prior Specifications

Unknown parameters:

- etiology regression coefficients ($\{\Gamma_\ell^\pi\}$),
- subclass mixing weight parameters for cases ($\{\Gamma_k^\eta\}$) and controls ($\{\Gamma_k^\nu\}$),
- true and false positive rates ($\Theta = \{\theta_k^{(j)}\}$, $\Psi = \{\psi_k^{(j)}\}$).

To avoid potential overfitting, we *a priori* introduce:

- (a) few non-trivial subclasses via novel additive half-Cauchy prior for the intercepts $\{\mu_{k0}\}$
- (b) for continuous variable: smooth regression curves $\pi_\ell(\cdot)$, $\nu_k(\cdot)$ and $\eta_k(\cdot)$ by Bayesian Penalized-splines (Lang, 2004) combined with shrinkage priors on spline coefficients (Ni et.al, 2015) (to encourage towards constant values)

Posterior Inference

Use Markov chain Monte Carlo (MCMC) algorithm to approximate joint posterior distribution

Posterior Inference

Use Markov chain Monte Carlo (MCMC) algorithm to approximate joint posterior distribution

- Posterior inference is flexible and can be obtained from any functions of model parameters and individual latent variables

Posterior Inference

Use Markov chain Monte Carlo (MCMC) algorithm to approximate joint posterior distribution

- Posterior inference is flexible and can be obtained from any functions of model parameters and individual latent variables

Fit npLCMs (w/ or w/out covariates using R package baker
(<https://github.com/zhenkewu/baker>)

Posterior Inference

Use Markov chain Monte Carlo (MCMC) algorithm to approximate joint posterior distribution

- Posterior inference is flexible and can be obtained from any functions of model parameters and individual latent variables

Fit npLCMs (w/ or w/out covariates using R package baker (<https://github.com/zhenkewu/baker>))

- calls Bayesian model fitting software JAGS 4.2.0 (Plummer et al., 2003) from within R
- provides functions to visualize the posterior distributions of the unknowns
- also performs posterior predictive model checking

Simulation Results

Simulation Results

- Simulation I: demonstrate flexible statistical inferences about the PEF functions $\{\pi_\ell(\cdot)\}$


Simulation Results

- Simulation I: demonstrate flexible statistical inferences about the PEF functions $\{\pi_\ell(\cdot)\}$
- Simulation II: determining overall PEF π_ℓ^* (empirical average) to quantify disease burdens in a population (potential policy interest)


Simulation I Results

- $N_d = 500$ cases and $N_u = 500$ controls for each of two levels of S (discrete covariate); Uniformly sample the subjects' enrollment dates over a period of 300 days.


Etiology Regression Curves: Seasonality




Simulation I: Recovery of Truth $\pi_l^0(t, S = s)$




Simulation II: Regression Model Reduces the Percent Relative Bias in Recovering the Overall PEFs π_l^*




Simulation II: Regression Model Reduces the Percent Relative Bias in Recovering the Overall PEFs π_ℓ^*




Simulation II: Regression Model Produces More Valid 95% Crls in Recovering the Overall PEFs π_ℓ^*



Simulation II: Regression Model Produces More Valid 95% Crls in Recovering the Overall PEFs π_ℓ^*





Discussion

Limitations: Current npLCM methods do not describe the relationship between covariates and PEFs

Contribution: We address this by including covariate regression in the framework to estimate disease etiology

Three features of our approach:

Discussion

Limitations: Current npLCM methods do not describe the relationship between covariates and PEFs

Contribution: We address this by including covariate regression in the framework to estimate disease etiology

Three features of our approach:

- 1) allows analysts to specify a model that links important covariates to PEFs

Discussion

Limitations: Current npLCM methods do not describe the relationship between covariates and PEFs

Contribution: We address this by including covariate regression in the framework to estimate disease etiology

Three features of our approach:

- 1) allows analysts to specify a model that links important covariates to PEFs
- 2) produces covariate-dependent reference distribution for controls, which is critical for assigning cause-specific probabilities to a given case
 - because we can compare control measurements to case measurements with similar covariate values

Discussion

Limitations: Current npLCM methods do not describe the relationship between covariates and PEFs

Contribution: We address this by including covariate regression in the framework to estimate disease etiology

Three features of our approach:

- 1) allows analysts to specify a model that links important covariates to PEFs
- 2) produces covariate-dependent reference distribution for controls, which is critical for assigning cause-specific probabilities to a given case
 - because we can compare control measurements to case measurements with similar covariate values
- 3) TPR priors are only used once; avoids overly-optimistic etiology uncertainty estimates.

Future Directions

Future disease etiology studies may benefit from four improvements:

- Bayesian additive regression tree with variable selection (rather than GAM) may provide an alternative for characterizing interactions
- Class-specific predictor selection methods could be useful in the presence of many predictors in stabilizing and improving PEF estimates
- When the subset of population pathogen-cause combinations is unknown, combining the proposed method with subset selection procedures could be fruitful
- Scalable posterior inference for regression parameters will likely improve computational speed when number of disease classes/covariates is larger

Thank You!

Student
Irena Chen

Collaborators
Scott Zeger
Katherine O'Brien
Maria Deloria-Knoll
Laura Hammitt


Funding
Patient-Centered Outcome Research Institute
[PCORI ME-1408-20318]
Bill & Melinda Gates Foundation [48968]
Michigan Precision Health Investigator Award
National Cancer Institute (P30CA046592,
U01CA229437)

Some References (More at: zhenkewu.com)


1. **Wu Z** and Chen I (2019+).
Regression Analysis of Dependent Binary Data: Estimating Disease Etiology from Case-Control Studies.
Submitted. <https://arxiv.org/abs/1906.08436>
2. **Wu Z**, Deloria-Knoll M and Zeger SL (2019+).
A Bayesian Approach to Restricted Latent Class Models for Scientifically-Structured Clustering of Multivariate Binary Outcomes.
Submitted. <https://doi.org/10.1101/400192>
3. **Wu Z**, Deloria-Knoll M and Zeger SL (2017).
Nested Partially-Latent Class Models for Estimating Disease Etiology from Case-Control Data.
Biostatistics. 18 (2): 200-213.
4. **Wu Z**, Deloria-Knoll M, Hammitt LL, and Zeger SL, for the PERCH Core Team (2015).
Partially Latent Class Models (pLCM) for Case-Control Studies of Childhood Pneumonia Etiology.
Journal of the Royal Statistical Society: Series C (Applied Statistics). 65:97-114.

Simulation I: Recovery of $\nu_k(t)$ and $\eta_k(t)$

True $K^0 = 2$; Model fitted using a working number $K = 7$



(a) case



Appendix: Simulation II Setup

- npLCM regression analysis with $K^* = 3$, $R = 200$ replication data sets simulated under 48 different scenarios
- $L = J = 3, 6, 9$ causes, under single-pathogen-cause assumption, BrS measurements made on N_d cases and N_u controls for each level of X where $N_d = N_u = 250$ or 500 .
- $\phi_\ell(X) = \beta_{0\ell} + \beta_{1\ell} \mathbb{I}\{X = 2\}$ take two sets of values to reflect PEF variability across X : i) $\beta_0^i = (0, 0, 0, 0, 0, 0)$, $\beta_1^i = (-1.5, 0, -1.5, -1.5, 0, -1.5)$; ii) $\beta_0^{ii} = (1, 0, 1, 1, 0, 1)$ and $\beta_1^{ii} = (-1.5, 1, -1.5, -1.5, 1, -1.5)$
- TPRs $\theta_k^{(j)} = 0.95$ or 0.8 and FPRs $(\psi_1^{(j)}, \psi_2^{(j)}) \in \{(0.5, 0.05), (0.5, 0.15)\}$, for $j = 1, \dots, J$.
- $\nu_k(W) = \eta_k(W) = \text{logit}^{-1}(\gamma_{k0} + \gamma_{k1} \mathbb{I}\{W = 2\})$ where $(\gamma_{10}, \gamma_{11}) = (-0.5, 1.5)$ and $(\gamma_{20}, \gamma_{21}) = (1, -1.5)$.

Appendix




Figure: Posterior distributions of the stratum-specific (Row 1 and 2) and the overall (Bottom Row) PEFs based on a simulation with a two-level discrete covariate and $L = J = 6$ causes. The vertical gray lines indicate the 2.5% and 97.5% posterior quantiles, respectively; The truths are indicated by vertical blue dashed lines. *Row 1-2*) PEFs by stratum (level = 1,2) and cause (A-F); *Bottom*) π_ℓ^* : overall population etiologic fraction for cause A-F (empirical average of the two PEFs above).

Appendix




Figure: NPLCM analyses with or without regression perform similarly in terms of percent relative bias (top) and empirical coverage rates (bottom) over $R = 100$ replications in simulations where the case and control subclass weights *do not* vary by covariates. Each panel corresponds to one of 16 combinations of true

Appendix






Figure: NPLCM analyses with or without regression perform similarly in terms of percent relative bias (top) and empirical coverage rates (bottom) over $R = 100$ replications in simulations where the case and control subclass weights *do not* vary by covariates. Each panel corresponds to one of 16 combinations of true

Appendix



(a) Cause: RSV



(b) Cause: NoS

Figure: Individual etiology fraction estimates for RSV (left) and NoS (right) differ by age and season among HIV negative and severe pneumonia cases for whom the seven pathogens were *all tested negative* in the nasopharyngeal specimens.