

Phenylspirodrimane Derivatives From Cultures of the Fungus *Stachybotrys chartarum* YIM DT 10079

Natural Product Communications
Volume 14(10): 1–4
© The Author(s) 2019
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/1934578X19878906
journals.sagepub.com/home/npx



Zhang-Gui Ding^{1*}, Jian-Hai Ding^{2*}, Jiang-Yuan Zhao^{1*}, Ming-Gang Li¹, Dong-Bao Hu³, Xian-Jun Jiang⁴, Shao-Jie Gu⁴, Fei Wang⁴, and Meng-Liang Wen¹

Abstract

A new phenylspirodrimane derivative, stachartin F (**1**), and 2 known secondary metabolites stachybonoid E (**2**) and stachybonoid F (**3**) were isolated from cultures of the tin mine tailings-associated fungus *Stachybotrys chartarum* YIM DT 10079. Their structures were determined with the help of extensive spectroscopic analyses and absolute configuration of compound **1** was rationalized by quantum chemical calculations of the electronic circular dichroism spectra.

Keywords

Stachybotrys chartarum, fungus, secondary metabolite, phenylspirodrimane derivative

Received: February 28th, 2019; Accepted: July 24th, 2019.

The phenylspirodrimanes, structurally characterized by the fusion of a drimane-based sesquiterpene with a phenyl moiety, are known as characteristic secondary metabolites of the fungus *Stachybotrys chartarum*. These compounds have exhibited diverse pharmacological activities, such as antihyperlipidemic activities,¹ anti-HIV activity,² osteoclast differentiation inhibitor,³ and antimalarial activity.⁴ As part of our ongoing search for novel molecules from extremophiles,^{5,6} *S. chartarum* YIM DT 10079 was isolated from a soil sample collected from the Datun tin mine tailings area, Yunnan, China. An EtOAc extract of cultures of *S. chartarum* YIM DT 10079 was subjected to investigation, which resulted in the isolation of a new phenylspirodrimane derivative, stachartin F (**1**), and the known stachybonoid E (**2**) and stachybonoid F (**3**)⁷ (Figure 1). Herein, we report the isolation and structure elucidation of these compounds, and stachartin F (**1**) was tested for its cytotoxicity against 5 human cancer cell lines.

Compound **1** was isolated as a white powder. Its molecular formula was established as C₂₈H₃₉NO₆ by high-resolution electrospray ionization mass spectrometry (HR-ESI-MS) at *m/z* 508.2671 [M+Na]⁺ (calcd 508.2670), indicating 10 degrees of unsaturation. Infrared Spectroscopy (IR) absorption bands at 3429, 1725, and 1658 cm⁻¹ implied the presence of hydroxy and carbonyl functionalities. Comparison of the ¹H and ¹³C Nuclear magnetic resonance spectroscopy (NMR) data (Table 1) with those of chartarlactam H¹ showed the presence of the same phenylspirodrimane skeleton, except the side chain moiety on the nitrogen was replaced by a methyl butyrate

group in **1**, which was further confirmed by the Heteronuclear Multiple Bond Correlation (HMBC) correlations (Figure 2) from H₃-5" (δ_{H} 3.54) and H₂-2" (δ_{H} 1.79) to C-4" (δ_{C} 173.1) and from H₂-1" (δ_{H} 3.35) to C-7' (δ_{C} 49.3) and C-8' (δ_{C} 166.1), and ¹H-¹H correlation spectroscopy (COSY) correlations of H-2"

¹Key Laboratory for Microbial Resources, Ministry of Education, Yunnan Institute of Microbiology, Yunnan University, Kunming, P.R. China

²Engineering and Technology Research Center of Liupanshan Resources, College of Chemistry and Chemical Engineering, Ningxia Normal University, Guyuan, P. R. China

³School of Chemical Biology and Environment, Yuxi Normal University, P.R. China

⁴BioBioPha Co., Ltd., Kunming, P.R. China

*The authors Zhang-Gui Ding, Jian-Hai Ding and Jiang-Yuan Zhao contributed equally to the work.

Corresponding Authors:

Jian-Hai Ding, Engineering and Technology Research Center of Liupanshan Resources, College of Chemistry and Chemical Engineering, Ningxia Normal University, Guyuan 756000, P. R. China.

Email: dingjh_nx@126.com

Fei Wang, BioBioPha Co., Ltd., Kunming 650201, P.R. China.

Email: f.wang@mail.biobioph.com

Dong-Bao Hu, School of Chemical Biology and Environment, Yuxi Normal University, Yuxi 653100, P.R. China.

Email: hudongbao@126.com

Meng-Liang Wen, Key Laboratory for Microbial Resources, Ministry of Education, Yunnan Institute of Microbiology, Yunnan University, Kunming 650091, P.R. China.

Email: mlwen@ynu.edu.cn




Table 1. ^1H and ^{13}C NMR Data for Compound **1**.

No.	δ_{H}	δ_{C}	No.	δ_{H}	δ_{C}
1a	1.69 (brt)	23.9 t	1'		112.5 s
1b	0.89 (brd)				
2a	1.79 (m)	25.1 t	2'		156.5 s
2b	1.36 (m)				
3	3.15 (m)	73.4 d	3'	6.31 (s)	101.4 d
4		37.4 s	4'		144.2 s
5	2.05 (dd, 12.2, 1.5)	39.3 d	5'		105.9 s
6a	1.45 (m)	20.6 d	6'		158.2 s
6b	1.38 (m)				
7a	1.47 (m)	30.5 t	7'	4.22	49.3 t
8	1.75 (m)	36.7 d	8'		166.1 s
9		98.3 s	1''	3.35 (m)	40.8 t
10		41.8 s	2''	1.79 (m)	23.3 t
11a	3.00 (d, 16.1)	30.9 t	3''	2.31 (m)	30.9 t
11b	2.63 (d, 16.1)				
12	0.59 (d, 6.6)	15.5 q	4''		173.1 s
13	0.87 (s)	28.7 q	5''	3.54 (s)	51.4 q
14	0.78 (s)	22.6 q	3-OH	4.09 (s)	
15	0.92 (s)	16.0 q	3'-OH	9.94 (s)	

with H-1'' and H-3'' (δ_{H} 2.31). Its relative configuration was determined by the Rotating frame overhauser enhancement spectroscopy (ROESY) correlations of H-3/H₃-14, H₃-14/H₃-15, H₃-15/H-8, and H₃-13/H-5, H-5/H α -7, H α -7/H₃-12 determined that H-3, H₃-14, and H₃-15 as being in β -orientations, while H₃-13, H-5, and H₃-12 as being in α -orientations (Figure 2). Consequently, compound **1** was established as stachartin F. The absolute configuration of **1** was determined by means of electronic circular dichroism (ECD). As shown in Figure 3, the calculated ECD spectrum of **1** matched well with the experimental one, suggesting the absolute configuration of **1** to be 3R, 5S, 8R, 9R, 10S.

Compound **1** was evaluated for its cytotoxicity against 5 human cancer cell lines using the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide (MTT) method as reported previously.⁸ Cisplatin (Sigma, United States) was used as the positive control. The compound was inactive (IC_{50} values $>40 \mu\text{M}$).

**Figure 1.** Chemical structures of **1** to **3** from *Stachybotrys chartarum* YIM DT 10079.


Experimental

General

Optical rotations, a Horiba SEPA-300 polarimeter; NMR, Avance III 600, Bruker DRX-500, and Bruker AM-400 spectrometers; HR-ESI-MS, an API-Qstar-Pulsar-1 spectrometer.

Fungal Material and Cultivation Conditions

Stachybotrys chartarum was isolated from a soil sample collected from the Datun tin mine tailings area, Yunnan, P.R. China. A voucher specimen (No. YIM DT 10079) was deposited at Yunnan Institute of Microbiology, Yunnan University. The culture medium consisted of glucose (1.0%), peptone from porcine meat (0.5%), yeast powder (0.5%), KH_2PO_4 (0.1%), and $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ (0.02%). Fermentation was carried out on a shaker at 200 rpm for 15 days.

**Figure 2.** Selected 2D NMR correlations of **1**.




Figure 3. Experimental and calculated electronic circular dichroism for compound **1**.

Extraction and Isolation

The culture broth (100 L) of *S. chartarum* YIM DT 10079 was filtered, and the filtrate was extracted 3 times with EtOAc, while the mycelium was extracted 3 times with CHCl₃/MeOH (1:1). The EtOAc layer together with the mycelium extraction was concentrated under reduced pressure to give a crude extract. The extract was subjected to column chromatography over silica gel (200–300 mesh) eluted with a gradient of CHCl₃/MeOH (1:0→0:1) to obtain 6 fractions (1–6). Fraction 3 eluted with CHCl₃/MeOH (80:1) was separated repeatedly by Sephadex LH-20 (CHCl₃/MeOH 1:1) to afford **1** (16.0 mg), **2** (25.0 mg), and **3** (200.0 mg).

Cytotoxicity Assay

Five human cancer cell lines, breast cancer SK-BR-3, hepatocellular carcinoma SMMC-7721, human myeloid leukemia HL-60, pancreatic cancer PANC-1, and lung cancer A-549 cells, were used in the cytotoxic assay. Cells were cultured in RPMI-1640 or in DMEM medium (Hyclone, United States), supplemented with 10% fetal bovine serum (Hyclone, United States) in 5% CO₂ at 37°C. The cytotoxicity assay was performed according to the MTT method in 96-well microplates.⁸ Briefly, 100 µL of adherent cells were seeded into each well of 96-well cell culture plates and allowed to adhere for 12 hours before addition of test compounds, while suspended cells were seeded just before drug addition with initial density of 1×10^5 cells/mL. Each tumor cell line was exposed to the test compound at concentrations of 0.0625, 0.32, 1.6, and 8 µM in triplicates for 48 hours, with cisplatin (Sigma, United States) as positive control. After compound treatment, cell viability was detected and a cell growth curve

was graphed. IC₅₀ values were calculated by Reed and Muench's method.⁹

Computational Methods

All Discrete fourier transformation (DFT) and Time dependent density functional theory (TD-DFT) calculations were carried out at 298 K in the gas phase with Gaussian 09. Conformational searches were carried out at the molecular mechanics level of theory employing MMFF force fields.^{10,11} The conformers with relative energy within 10 kcal/mol of the lowest-energy conformer were selected and further geometry optimized at the B3LYP/6-311++G(2d,p) level. All the lowest-energy conformers, which correspond to 99% of the total Boltzmann distribution, were selected for ECD spectra calculation. The Boltzmann factor for each conformer was calculated based on Gibbs free energy. Vibrational analysis at the B3LYP/6-311++G(2d,p) level of theory resulted in no imaginary frequencies, confirming the considered conformers as real minima. TD-DFT was employed to calculate excitation energy (in nm) and rotatory strength *R* in dipole velocity form, at the B3LYP/6-311++G(2d,p) level.

Stachartin F (1)

White powder.

$[\alpha]_D^{26}$: −6.0 (*c* 0.35, MeOH).

¹H and ¹³C NMR: Table 1.

HR-ESI-MS: *m/z* 508.2671 [M+Na]⁺ (calcd for C₂₈H₃₉NNaO₆, 508.2670)

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This project was supported by the Scientific Research Foundation of the Higher Education Institutions of Ningxia, China (NGY2018-107), the Ningxia Normal University Research Grant (Grant number NXSFDZDA1904), the First-rate Discipline (Education) Construction Project of the Higher Education Institutions of Ningxia (NXYLXK2017B110), China, the National Natural Sciences Foundation of China (31760090, 31760017), the Program for Innovation Talents of Science and Technology in Ningxia, China (KJT2015005), Youth talent cultivation project in Ningxia, China (NXRS2017(279)), the "312 Domestic Talents Project" of Ningxia, China (NXRS2018(139)), the Natural Science Foundation of Ningxia, China (2019AAC03245), the Key Research Foundation

of Ningxia, China (2019BDE03009), and the Guyuan Science and Technology Support Program (2018GYYQ0007).

Supplemental Material

Supplemental material for this article is available online.

References

1. Li Y, Wu C, Liu D, Proksch P, Guo P, Lin W. Chartarlactams A-P, phenylspirodrimanes from the sponge-associated fungus *Stachybotrys chartarum* with antihyperlipidemic activities. *J Nat Prod*. 2014;77(1):138-147.
2. Ma X, Li L, Zhu T, XH M, et al. Phenylspirodrimanes with anti-HIV activity from the sponge-derived fungus *stachybotrys chartarum* MXH-X73. *J Nat Prod*. 2013;76(12):2298-2306.
3. Kim JW, Ko S-K, Kim H-M, et al. Stachybotrysin, an osteoclast differentiation inhibitor from the marine-derived fungus *Stachybotrys* sp. KCB13F013. *J Nat Prod*. 2016;79(10):2703-2708.
4. Hemtasin C, Kanokmedhakul S, Kanokmedhakul K, et al. Cytotoxic pentacyclic and tetracyclic aromatic sesquiterpenes from *Phomopsis archeri*. *J Nat Prod*. 2011;74(4):609-613.
5. Chunyu W-X, Ding Z-G, Li M-G, et al. Stachartins A - E, phenylspirodrimanes from the tin mine tailings-associated fungus *Stachybotrys chartarum*. *Helv Chim Acta*. 2016;99(8):583-587.
6. Ding Z-G, Ding J-H, Zhao J-Y, et al. A new phenylspirodrimane dimer from the fungus *Stachybotrys chartarum*. *Fitoterapia*. 2018;125:94-97.
7. Zhang P, Li Y, Jia C, et al. Antiviral and anti-inflammatory meroterpenoids: stachybonoids A-F from the crinoid-derived fungus *stachybotrys chartarum* 952. *RSC Adv*. 2017;7(79):49910-49916.
8. Mosmann T. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J Immunol Methods*. 1983;65(1-2):55-63.
9. Reed LJ, Muench H. A simple method of estimating fifty percent endpoints. *Am J Hyg*. 1938;27:493-497.
10. Qm L, Ren J, Zhou BD, et al. Determining the absolute configuration of benzopyrenomycin by optical rotation, electronic circular dichroism, and population analysis of different conformations via DFT methods and experiments. *Tetrahedron*. 2013;69:3067-3074.
11. Db H, Wx L, Zhao ZZ, et al. Highly unsaturated pyranone derivatives from the basidiomycete *Junghuhnianitida*. *Tetrahedron Lett*. 2014;55:6530-6533.