

# Chapter 4

## Motion in a Plane

### Solutions

#### SECTION - A

##### Objective Type Questions

(Scalars and Vectors, Multiplication of Vectors by Real Numbers, Addition and Subtraction of Vectors - Graphical Methods, Resolution of Vectors, Vector Addition - Analytical Method)

1. Which of the following is a vector?

- (1) Current                      (2) Time                      (3) Acceleration                      (4) Volume

**Sol.** Answer (3)

Acceleration is a vector quantity.

2. The change in a vector may occur due to

- (1) Rotation of frame of reference                      (2) Translation of frame of reference  
(3) Rotation of vector                      (4) Both (1) & (3)

**Sol.** Answer (3)

Change in a vector may occur due to rotation of vector and not due to rotation of frame of reference.

3. Which one of the following pair cannot be the rectangular components of force vector of 10 N?

- (1) 6 N & 8 N                      (2) 7 N &  $\sqrt{51}$  N                      (3)  $6\sqrt{2}$  N &  $2\sqrt{7}$  N                      (4) 9 N & 1 N

**Sol.** Answer (4)

$$\text{The vector magnitude} = \sqrt{A_x^2 + A_y^2}$$

$$\text{Vector magnitude} = 10$$

But (4) option gives the magnitude

$$\Rightarrow \sqrt{9^2 + 1^2} = \sqrt{82} \neq 10 \quad [\text{by trial method check options}]$$

4. The resultant of two vectors at an angle  $150^\circ$  is 10 units and is perpendicular to one vector. The magnitude of the smaller vector is

- (1) 10 units                      (2)  $10\sqrt{3}$  units                      (3)  $10\sqrt{2}$  units                      (4)  $5\sqrt{3}$  units

**Sol.** Answer (2)

$$\Rightarrow R^2 + A^2 = B^2 \quad \dots(1)$$

$$R = 10$$


$$\text{Also } \tan 30^\circ = \frac{\text{Perpendicular}}{\text{Base}}$$

$$\frac{1}{\sqrt{3}} = \frac{R}{A}$$

$$\text{From equation (1) } A = 10\sqrt{3}$$

$$(10)^2 + (10\sqrt{3})^2 = B^2$$

$$B = 20$$



5. Two vectors, each of magnitude  $A$  have a resultant of same magnitude  $A$ . The angle between the two vectors is

(1)  $30^\circ$

(2)  $60^\circ$

(3)  $120^\circ$

(4)  $150^\circ$

**Sol.** Answer (3)

$$|\vec{A}| = |\vec{B}| = |\vec{R}|$$




$$R = \sqrt{A^2 + B^2 + 2AB\cos\theta}$$

$$A^2 = A^2 + A^2 + 2A^2\cos\theta$$

$$-A^2 = 2A^2\cos\theta$$

$$\cos\theta = -\frac{1}{2} \Rightarrow \theta = 120^\circ$$

6. Let  $\theta$  be the angle between vectors  $\vec{A}$  and  $\vec{B}$ . Which of the following figures correctly represents the angle  $\theta$ ?



**Sol.** Answer (3)

To find angle between vectors, they will be joined either head to head or tail to tail.

7.  $\vec{A}$  is a vector of magnitude 2.7 units due east. What is the magnitude and direction of vector  $4\vec{A}$ ?

(1) 4 units due east

(2) 4 units due west

(3) 2.7 units due east

(4) 10.8 units due east

**Sol.** Answer (4)

$$\vec{A} = 2.7\hat{i}$$

Vector  $4\vec{A}$

$$\Rightarrow 4(2.7\hat{i}) = 10.8\hat{i} \text{ or } 10.8 \text{ units due east.}$$

8. Two forces of magnitude 8 N and 15 N respectively act at a point. If the resultant force is 17 N, the angle between the forces has to be

(1)  $60^\circ$  (2)  $45^\circ$  (3)  $90^\circ$  (4)  $30^\circ$

**Sol.** Answer (3)

$$R = \sqrt{A^2 + B^2 + 2AB\cos\theta}$$

$$A = 8, B = 15, R = 17$$

$$17^2 = 8^2 + 15^2 + 2 \times 8 \times 15 \times \cos \theta$$

$$289 = 64 + 225 + 240 \cos \theta$$

$$\Rightarrow 289 = 289 + 24 \cos \theta$$

$$24 \cos \theta = 0$$

$$\cos \theta = 0 \Rightarrow \theta = 90^\circ$$

9. Two forces of 10 N and 6 N act upon a body. The direction of the forces are unknown. The resultant force on the body may be

(1) 15 N (2) 3 N (3) 17 N (4) 2 N

**Sol.** Answer (1)

The resultant of two vectors always lie between  $(A + B)$  &  $(A - B)$ .

So the resultant of 10 N & 6 N should lie between 16 N & 4 N.

So answer is 15 N.

10. The vector  $\vec{OA}$  where O is origin is given by  $\vec{OA} = 2\hat{i} + 2\hat{j}$ . Now it is rotated by  $45^\circ$  anticlockwise about O. What will be the new vector?

(1)  $2\sqrt{2}\hat{j}$  (2)  $2\hat{j}$  (3)  $2\hat{i}$  (4)  $2\sqrt{2}\hat{i}$


**Sol.** Answer (1)

$$\vec{OA} = 2\hat{i} + 2\hat{j}$$

$$|\vec{OA}| = \sqrt{4+4} \Rightarrow 2\sqrt{2}$$

On rotating by an angle of  $45^\circ$  anticlockwise it will lie along y-axis.

$$\text{So } \vec{A} = 2\sqrt{2}\hat{j}$$



11. If the sum of two unit vectors is also a unit vector, then magnitude of their difference and angle between the two given unit vectors is

(1)  $\sqrt{3}, 60^\circ$  (2)  $\sqrt{3}, 120^\circ$  (3)  $\sqrt{2}, 60^\circ$  (4)  $\sqrt{2}, 120^\circ$

**Sol.** Answer (2)

$$|\vec{R}| = |\vec{A} + \vec{B}| = \sqrt{A^2 + B^2 + 2AB\cos\theta}$$

$$|\vec{A}| = |\vec{B}| = |\vec{R}| = 1$$

$$1 = 1 + 1 + 2 \times 1 \times 1 \times \cos \theta$$

$$\cos \theta = -\frac{1}{2} \Rightarrow \theta = 120^\circ$$

$$|\vec{R}| = |\vec{A} - \vec{B}| = \sqrt{A^2 + B^2 - 2AB\cos 120^\circ}$$



$$= \sqrt{1^2 + 1^2 - 2 \times 1 \times 1 \times \left(-\frac{1}{2}\right)} = \sqrt{3} = |\vec{A} - \vec{B}|$$

**(Motion in a Plane, Motion in a Plane with Constant Acceleration, Relative Velocity in Two Dimensions)**

12. A car moves towards north at a speed of 54 km/h for 1 h. Then it moves eastward with same speed for same duration. The average speed and velocity of car for complete journey is

- (1) 54 km/h, 0                      (2)  $15 \text{ m/s}, \frac{15}{\sqrt{2}} \text{ m/s}$                       (3) 0, 0                      (4)  $0, \frac{54}{\sqrt{2}} \text{ km/h}$

**Sol.** Answer (2)



$$\text{Displacement} = \frac{54\sqrt{2}}{\text{Km}}$$

$$\text{Distance} = 2 \times 54 = 108 \text{ Km}$$



$$\text{Average speed} = \frac{108}{2} = 54 \text{ Km h}^{-1} \times \frac{5}{18} = 15 \text{ ms}^{-1}$$

$$\text{Average velocity} = \frac{\text{disp.}}{\text{time}} = \frac{54\sqrt{2}}{2} \Rightarrow 27\sqrt{2} \times \frac{5}{18} \Rightarrow \frac{15}{\sqrt{2}} \text{ m/s}$$

13. Ram moves in east direction at a speed of 6 m/s and Shyam moves  $30^\circ$  east of north at a speed of 6 m/s. The magnitude of their relative velocity is

- (1) 3 m/s                      (2) 6 m/s                      (3)  $6\sqrt{3} \text{ m/s}$                       (4)  $6\sqrt{2} \text{ m/s}$


**Sol.** Answer (2)



$$|\vec{V}_{RS}| = \sqrt{V_R^2 + V_S^2 - 2V_R V_S \cos \theta}$$

$$= \sqrt{6^2 + 6^2 - 2 \times 6^2 \times \frac{1}{2}}$$

$$= 6 \text{ ms}^{-1}$$



14. A train is running at a constant speed of 90 km/h on a straight track. A person standing at the top of a boggy moves in the direction of motion of the train such that he covers 1 meters on the train each second. The speed of the person with respect to ground is

- (1) 25 m/s                      (2) 91 km/h                      (3) 26 km/h                      (4) 26 m/s

**Sol.** Answer (4)

$$V_T = 90 \text{ Km h}^{-1} = 90 \times \frac{5}{18} = 25 \text{ ms}^{-1}$$

$$V_m = ?$$


$$d = \text{speed} \times \text{time}$$

$$d_{\text{net}} = V_{\text{net}} \times t$$

$$1 = (V_m - 25) \times 1$$

$$V_m = 26 \text{ ms}^{-1}$$

15. Figure shows two ships moving in x-y plane with velocities  $V_A$  and  $V_B$ . The ships move such that B always remains north of A. The ratio  $\frac{V_A}{V_B}$  is equal to



- (1)  $\cos\theta$                       (2)  $\sin\theta$                       (3)  $\sec\theta$                       (4)  $\csc\theta$

**Sol.** Answer (1)

If ship B is always north of ship A then, their horizontal component should be equal, so,

$$V_A = V_B \cos \theta$$

$$\Rightarrow \frac{V_A}{V_B} = \cos \theta$$

16. Four persons P, Q, R and S are initially at the four corners of a square of side  $d$ . Each person now moves with a constant speed  $v$  in such a way that P always moves directly towards Q, Q towards R, R towards S, and S towards P. The four persons will meet after time

- (1)  $\frac{d}{2v}$                       (2)  $\frac{d}{v}$                       (3)  $\frac{3d}{2v}$                       (4) They will never meet

**Sol.** Answer (2)


$$T = \frac{d}{v_{rel}}$$

$$v_{rel} = v - v \cos 90^\circ$$

$$= v - 0$$

$$= v$$

$$T = \frac{d}{v}$$



17. A person, reaches a point directly opposite on the other bank of a flowing river, while swimming at a speed of 5 m/s at an angle of  $120^\circ$  with the flow. The speed of the flow must be

- (1) 2.5 m/s                      (2) 3 m/s                      (3) 4 m/s                      (4) 1.5 m/s


**Sol.** Answer (1)

For drift to be zero

$$u = v \sin 30^\circ$$

$$= 5 \times \frac{1}{2}$$

$$= 2.5 \text{ ms}^{-1}$$



18. A car with a vertical windshield moves in a rain storm at a speed of 40 km/hr. The rain drops fall vertically with constant speed of 20 m/s. The angle at which rain drops strike the windshield is

(1)  $\tan^{-1} \frac{5}{9}$

(2)  $\tan^{-1} \frac{9}{5}$


(3)  $\tan^{-1} \frac{3}{2}$

(4)  $\tan^{-1} \frac{2}{3}$

**Sol.** Answer (1)

$$\tan \theta = \frac{v_m}{v_r} = \frac{20 \times 5}{20}$$

$$\theta = \tan^{-1} \left( \frac{5}{9} \right)$$



### (Projectile Motion)

19. A particle projected from origin moves in x-y plane with a velocity  $\vec{v} = 3\hat{i} + 6x\hat{j}$ , where  $\hat{i}$  and  $\hat{j}$  are the unit vectors along x and y axis. Find the equation of path followed by the particle

(1)  $y = x^2$

(2)  $y = \frac{1}{x^2}$

(3)  $y = 2x^2$

(4)  $y = \frac{1}{x}$

**Sol.** Answer (1)

**Method 1:**

$$\vec{V} = 3\hat{i} + 6x\hat{j}$$

$$\text{also } \vec{V} = \frac{dx}{dt}\hat{i} + \frac{dy}{dt}\hat{j}$$

$$\Rightarrow \frac{dx}{dt} = 3,$$

$$\int dx = \int 3dt$$

$$x = 3t$$

$$\frac{dy}{dt} = 6x$$

$$dy = 6x \times dt$$

$$\int dy = \int 6 \times 3t dt$$

$$= 18 \int t dt \Rightarrow 18 \times \frac{t^2}{2}$$

$$y = 9t^2$$

$$= 9 \times \frac{x^2}{9}$$

$$y = x^2$$

**Method 2:**

$$V_x \hat{i} + V_y \hat{j} = \vec{V}$$

$$V_x = 3$$

$$V_y = 6x$$

We know

$$\frac{dy}{dx} = \tan \theta = \frac{V_y}{V_x}$$

$$\frac{dy}{dx} = \frac{6x}{3}$$

$$\int_0^y dy = \int_0^x 2x dx$$

$$y = x^2$$

20. A body of mass 1 kg is projected from ground at an angle  $30^\circ$  with horizontal on a level ground at a speed 50 m/s. The magnitude of change in momentum of the body during its flight is ( $g = 10 \text{ m/s}^2$ )

(1)  $50 \text{ kg ms}^{-1}$  (2)  $100 \text{ kg ms}^{-1}$  (3)  $25 \text{ kg ms}^{-1}$  (4) Zero


**Sol.** Answer (1)

$$\Rightarrow \text{The change in momentum} = -2mu \sin \theta \hat{j}$$

$$|\Delta \vec{p}| = 2mu \sin \theta$$

$$= 2 \times 1 \times 50 \times \sin 30^\circ$$

$$|\Delta \vec{p}| = 50 \text{ Kg ms}^{-1}$$



21. Two projectiles are projected at angles  $\left(\frac{\pi}{4} + \theta\right)$  and  $\left(\frac{\pi}{4} - \theta\right)$  with the horizontal, where  $\theta < \frac{\pi}{4}$ , with same speed.

The ratio of horizontal ranges described by them is

(1)  $\tan \theta : 1$  (2)  $1 : \tan^2 \theta$  (3)  $1 : 1$  (4)  $1 : \sqrt{3}$

**Sol.** Answer (3)

The horizontal range is same when the angles of projection are complimentary to each other.

22. A shell is fired vertically upwards with a velocity  $v_1$  from a trolley moving horizontally with velocity  $v_2$ . A person on the ground observes the motion of the shell as a parabola, whose horizontal range is

(1)  $\frac{2v_1^2 v_2}{g}$  (2)  $\frac{2v_1^2}{g}$  (3)  $\frac{2v_2^2}{g}$  (4)  $\frac{2v_1 v_2}{g}$

**Sol.** Answer (4)

There is no acceleration in the horizontal direction.

$$S_x = U_x T + \frac{1}{2} a_0 \times T^2$$

$$R = U_x T$$

$$S_y = U_y T + \frac{1}{2} g_y T^2$$

$$0 = V_1 T - \frac{1}{2} g T^2$$

$$\Rightarrow V_1 T = \frac{1}{2} g T$$


$$T = \frac{2V_1}{g}$$

We know,

$$(R) \text{ range} = (\text{Horizontal velocity } 4x) \times \text{flight + time } (T)$$

$$\text{i.e., } R = 4x \times T$$

$$R = V_2 \times \frac{2V_1}{g} \Rightarrow \frac{2V_1 V_2}{g}$$



23. The position coordinates of a projectile projected from ground on a certain planet (with no atmosphere) are given by  $y = (4t - 2t^2)$  m and  $x = (3t)$  metre, where  $t$  is in second and point of projection is taken as origin. The angle of projection of projectile with vertical is

- (1)  $30^\circ$  (2)  $37^\circ$  (3)  $45^\circ$  (4)  $60^\circ$

**Sol.** Answer (2)

$$y = 4t - 2t^2$$

$$x = 3t$$

$$\mathbf{V} = V_x \hat{i} + V_y \hat{j}$$

$$V_x = \frac{dx}{dt}, V_y = \frac{dy}{dt}$$

$$V_x = 3, V_y = 4 - 4t$$

$$\text{for } t = 0, V_y = 4$$

$$\tan \theta = \frac{V_y}{V_x} = \frac{4}{3}$$

$$\theta = 53^\circ \text{ with horizontal}$$

With vertical

$$\theta = 37^\circ$$


24. A particle is projected from ground with speed 80 m/s at an angle  $30^\circ$  with horizontal from ground. The magnitude of average velocity of particle in time interval  $t = 2$  s to  $t = 6$  s is [Take  $g = 10 \text{ m/s}^2$ ]

- (1)  $40\sqrt{2}$  m/s (2) 40 m/s (3) Zero (4)  $40\sqrt{3}$  m/s

**Sol.** Answer (4)

Average velocity of the projectile when it is at the same vertical height is :  $u \cos \theta$ .

$$\Rightarrow 80 \times \cos 30^\circ \Rightarrow 40\sqrt{3} \text{ m/s}.$$



25. A stone projected from ground with certain speed at an angle  $\theta$  with horizontal attains maximum height  $h_1$ . When it is projected with same speed at an angle  $\theta$  with vertical attains height  $h_2$ . The horizontal range of projectile is

- (1)  $\frac{h_1 + h_2}{2}$  (2)  $2h_1 h_2$  (3)  $4\sqrt{h_1 h_2}$  (4)  $h_1 + h_2$



**Sol.** Answer (3)

When the angles are complimentary the range is same,

$$h_1 = \frac{u^2 \sin^2 \theta}{2g},$$

$$h_2 = \frac{u^2 \sin^2 (90 - \theta)}{2g}$$

$$h_1 = \frac{u^2 \sin^2 \theta}{2g}$$





$$h_2 = \frac{u^2 \cos^2 \theta}{2g}$$

$$h_1 h_2 = \frac{u^4 \sin^2 \theta \cos^2 \theta}{4g^2} \Rightarrow \left( \frac{2u \sin \theta \cos \theta}{g} \right)^2 \times \frac{1}{4g} \times \frac{1}{4}$$

$$h_1 h_2 = R^2 \frac{1}{16} \Rightarrow R^2 = 16 h_1 h_2$$

$$R = 4(\sqrt{h_1 h_2})$$

26. Two objects are thrown up at angles of  $45^\circ$  and  $60^\circ$  respectively, with the horizontal. If both objects attain same vertical height, then the ratio of magnitude of velocities with which these are projected is

(1)  $\sqrt{\frac{5}{3}}$

(2)  $\sqrt{\frac{3}{5}}$

(3)  $\sqrt{\frac{2}{3}}$

(4)  $\sqrt{\frac{3}{2}}$

**Sol.** Answer (4)

$$h_1 = h_2$$

$$\frac{u_1^2 \sin^2 45^\circ}{2g} = \frac{v_2^2 \sin^2 60^\circ}{2g}$$

$$\frac{u_1^2}{v_2^2} = \frac{\frac{\sqrt{3}}{2} \times \frac{\sqrt{3}}{2}}{\frac{1}{2}} = \frac{3}{2}$$

$$\frac{v_1}{v_2} = \sqrt{\frac{3}{2}}$$

27. For an object projected from ground with speed  $u$  horizontal range is two times the maximum height attained by it. The horizontal range of object is

(1)  $\frac{2u^2}{3g}$

(2)  $\frac{3u^2}{4g}$

(3)  $\frac{3u^2}{2g}$

(4)  $\frac{4u^2}{5g}$

**Sol.** Answer (4)

$$R = 2H \text{ also, } \frac{H}{R} = \frac{1}{4} \tan \theta$$


$$\frac{H}{R} = \frac{1}{2} \Rightarrow \frac{1}{2} = \frac{1}{4} \tan \theta$$

$$\tan \theta = 2 = \frac{P}{B}$$

$$R = \frac{2u^2 \sin \theta \cos \theta}{g}$$

$$R = \frac{2u^2}{g} \cdot \frac{2}{\sqrt{5}} \times \frac{1}{\sqrt{5}}$$

$$R = \frac{4u^2}{5g}$$



28. The velocity at the maximum height of a projectile is  $\frac{\sqrt{3}}{2}$  times its initial velocity of projection ( $u$ ). Its range on the horizontal plane is

(1)  $\frac{\sqrt{3}u^2}{2g}$

(2)  $\frac{3u^2}{2g}$

(3)  $\frac{3u^2}{g}$

(4)  $\frac{u^2}{2g}$

**Sol.** Answer (1)

$$u_h = u \cos \theta$$


$$\frac{\sqrt{3}}{2} u = u \cos \theta$$

$$\Rightarrow \cos \theta = \frac{\sqrt{3}}{2}$$

$$\theta = 30^\circ$$

$$R = \frac{u^2 \sin 2\theta}{g}$$

$$= \frac{u^2 \sin 60^\circ}{g} \Rightarrow \frac{\sqrt{3} u^2}{2g} = R$$



29. A projectile is thrown into space so as to have a maximum possible horizontal range of 400 metres. Taking the point of projection as the origin, the co-ordinates of the point where the velocity of the projectile is minimum are

(1) (400, 100)

(2) (200, 100)

(3) (400, 200)

(4) (200, 200)

**Sol.** Answer (2)

$$R_{\max} = 400 \text{ m}$$


The velocity is minimum at the highest point

$$\Rightarrow H \rightarrow \frac{R}{2}$$

$$R = 4H$$

$$400 = 4 \times H$$

$$H = 100 \text{ m}$$



30. If the time of flight of a bullet over a horizontal range  $R$  is  $T$ , then the angle of projection with horizontal is

(1)  $\tan^{-1}\left(\frac{gT^2}{2R}\right)$

(2)  $\tan^{-1}\left(\frac{2R^2}{gT}\right)$

(3)  $\tan^{-1}\left(\frac{2R}{g^2T}\right)$

(4)  $\tan^{-1}\left(\frac{2R}{gT}\right)$

**Sol.** Answer (1)

$$T = \frac{2u \sin \theta}{g} \Rightarrow u = \frac{gT}{2 \sin \theta}$$

$$R = \frac{2u^2 \sin \theta \cos \theta}{g}$$

$$R = \frac{2u \sin \theta}{g} \times u \cos \theta$$

$$R = T \times u \cos \theta$$


$$R = T \times \frac{gT \cos \theta}{2 \sin \theta}$$

$$R = \frac{gT^2}{2} \frac{1}{\tan \theta}$$

$$\tan \theta = \frac{gT^2}{2R}$$

$$\theta = \tan^{-1}\left(\frac{gT^2}{2R}\right)$$


31. In the graph shown in figure, which quantity associated with projectile motion is plotted along y-axis?



- (1) Kinetic energy      (2) Momentum      (3) Horizontal velocity      (4) None of these

**Sol.** Answer (3)

It is the horizontal component of velocity that remains constant throughout the motion as there is no acceleration in that direction  $a_x = 0$ ,  $u_x = \text{constant}$



32. The equation of a projectile is  $y = ax - bx^2$ . Its horizontal range is

- (1)  $\frac{a}{b}$       (2)  $\frac{b}{a}$       (3)  $a + b$       (4)  $b - a$


**Sol.** Answer (1)

$$y = ax - bx^2$$

When the body lands then  $y = 0$ ,  $x = R$ ,  $0 = aR - bR^2$


$$aR = bR^2$$

$$R = \frac{a}{b}$$



33. Figure shows a projectile thrown with speed  $u = 20 \text{ m/s}$  at an angle  $30^\circ$  with horizontal from the top of a building  $40 \text{ m}$  high. Then the horizontal range of projectile is

- (1)  $20\sqrt{3} \text{ m}$       (2)  $40\sqrt{3} \text{ m}$   
(3)  $40 \text{ m}$       (4)  $20 \text{ m}$



**Sol.** Answer (2)

$$S_y = u_y T + \frac{1}{2} g_y T^2$$

$$-40 = 4 \sin 30^\circ T - \frac{1}{2} g T^2$$

$$-40 = 20 \times \frac{1}{2} T - 5T^2$$

$$-8 = 2T - T^2$$


$$T^2 - 2T - 8 = 0$$

$$T^2 - 4T + 2T - 8 = 0$$

$$T = -2, 4$$

$$R = u \cos \theta T = 20 \times \frac{\sqrt{3}}{2} \times 4$$

$$R = 40\sqrt{3} \text{ m}$$



34. When a particle is projected at some angle to the horizontal, it has a range  $R$  and time of flight  $t_1$ . If the same particle is projected with the same speed at some other angle to have the same range, its time of flight is  $t_2$ , then

$$(1) \quad t_1 + t_2 = \frac{2R}{g}$$

$$(2) \quad t_1 - t_2 = \frac{R}{g}$$

$$(3) \quad t_1 t_2 = \frac{2R}{g}$$

$$(4) \quad t_1 t_2 = \frac{R}{g}$$

**Sol.** Answer (3)

The angles has to be complimentary i.e., if  $\theta_1 \rightarrow \theta$ ,  $\theta_2 \rightarrow (90 - \theta)$

$$t_1 = \frac{2u \sin \theta}{g}, \quad t_2 = \frac{2u \sin(90 - \theta)}{g}$$

$$t_2 = \frac{2u \cos \theta}{g}$$

$$t_1 t_2 = \frac{2u \sin \theta}{g} \times \frac{2u \cos \theta}{g}$$

$$t_1 t_2 = \frac{2R}{g}$$

35. A projectile is thrown with velocity  $v$  at an angle  $\theta$  with horizontal. When the projectile is at a height equal to half of the maximum height, the vertical component of the velocity of projectile is

$$(1) \quad v \sin \theta \times 3$$

$$(2) \quad \frac{v \sin \theta}{3}$$

$$(3) \quad \frac{v \sin \theta}{\sqrt{2}}$$


$$(4) \quad \frac{v \sin \theta}{\sqrt{3}}$$

**Sol.** Answer (3)


$$v_B^2 = v^2 \sin^2 \theta - \frac{2g}{2} \left( \frac{u^2 \sin^2 \theta}{2g} \right)$$

$$v_B^2 = \frac{v^2 \sin^2 \theta}{2}$$

$$v_B = \frac{v \sin \theta}{\sqrt{2}}$$



36. In the given figure for a projectile



$$(1) \quad y = \left[ \frac{x_1 x_2}{x_1 - x_2} \right] \tan \theta$$

$$(2) \quad y = \left[ \frac{x_1 x_2}{x_1 + x_2} \right] \tan \theta$$

$$(3) \quad y = \left[ \frac{2x_1 x_2}{x_1 + x_2} \right] \cos \theta$$


$$(4) \quad y = \left[ \frac{2x_1 x_2}{x_1 + x_2} \right] \tan \theta$$

**Sol.** Answer (2)

The equation of trajectory for point 'P' can be written as :

$$y = x \tan \theta \left( 1 - \frac{x}{R} \right) = x_1 \tan \theta \left( 1 - \frac{x_1}{x_1 + x_2} \right) = x_1 \tan \theta \left( \frac{x_1 + x_2 - x_1}{x_1 + x_2} \right)$$

$$y = \frac{x_1 x_2}{x_1 + x_2} \tan \theta$$



37. Two paper screens  $A$  and  $B$  are separated by distance  $100\text{ m}$ . A bullet penetrates  $A$  and  $B$ , at points  $P$  and  $Q$  respectively, where  $Q$  is  $10\text{ cm}$  below  $P$ . If bullet is travelling horizontally at the time of hitting  $A$ , the velocity of bullet at  $A$  is nearly

- (1)  $100\text{ m/s}$  (2)  $200\text{ m/s}$  (3)  $600\text{ m/s}$  (4)  $700\text{ m/s}$

**Sol.** Answer (4)

$$10\text{ cm} \Rightarrow 10 \times 10^{-2}\text{ m} \Rightarrow 10^{-1} \Rightarrow 0.1\text{ m}$$

It is a case of horizontal projectile.


$$\text{So, } a_x = 0, u_x = 4, u_y = 0, a_y = -g$$

$$R = 100\text{ m}, T = \sqrt{\frac{2H}{g}} \Rightarrow \text{Time of flight}$$

$$R = u_x T$$

$$100 = u \sqrt{\frac{2 \times 0.1}{100}} \Rightarrow \frac{u\sqrt{2}}{10} = 100$$

$$u = \frac{1000}{\sqrt{2}} \approx 707\text{ ms}^{-1}$$



38. An object is projected from ground with speed  $u$  at angle  $\theta$  with horizontal. the radius of curvature of its trajectory at maximum height from ground is


- (1)  $\frac{u^2 \sin 2\theta}{g}$  (2)  $\frac{u^2 \cos^2 \theta}{g}$  (3)  $\frac{u^2 \sin^2 \theta}{g}$  (4)  $\frac{u^2 \sin^2 \theta}{2g}$

**Sol.** Answer (2)

$$a_c = \frac{v^2}{r}$$


$$r = \frac{v^2}{a_c}, \frac{u^2 \cos^2 \theta}{g}$$

$$r = \frac{u^2 \cos^2 \theta}{g}$$



**(Uniform Circular Motion)**

39. A particle is moving in a circle of radius  $r$  having centre at  $O$ , with a constant speed  $v$ . The magnitude of change in velocity in moving from  $A$  to  $B$  is



- (1)  $2v$  (2)  $0$  (3)  $\sqrt{3}v$  (4)  $v$

**Sol.** Answer (4)

$$|\Delta \vec{v}| = 2V \sin \frac{\theta}{2} = 2 \times V \times \sin \left( \frac{60^\circ}{2} \right) = 2 \times V \times \frac{1}{2} \Rightarrow V = |\Delta \vec{v}|$$

40. A car is going round a circle of radius  $R_1$  with constant speed. Another car is going round a circle of radius  $R_2$  with constant speed. If both of them take same time to complete the circles, the ratio of their angular speeds and linear speeds will be

(1)  $\sqrt{\frac{R_1}{R_2}}, \frac{R_1}{R_2}$  (2) 1, 1 (3)  $1, \frac{R_1}{R_2}$  (4)  $\frac{R_1}{R_2}, 1$

**Sol.** Answer (3)

The angular speed is given by

$$\omega = \frac{2\pi}{T}$$

$$\omega \propto \frac{1}{T} \Rightarrow \frac{\omega_1}{\omega_2} = \frac{T_2}{T_1}$$

$$\text{if } T_1 = T_2 \Rightarrow \omega_1 = \omega_2$$

$$\text{So, ratio} \Rightarrow 1 : 1$$

$$\text{and linear speed } v = R\omega$$

$$V \propto R$$

$$\frac{V_1}{V_2} = \frac{R_1}{R_2}$$

41. A body revolves with constant speed  $v$  in a circular path of radius  $r$ . The magnitude of its average acceleration during motion between two points in diametrically opposite direction is

(1) Zero (2)  $\frac{v^2}{r}$  (3)  $\frac{2v^2}{\pi r}$  (4)  $\frac{v^2}{2r}$

**Sol.** Answer (3)


$$a_{\text{avg}} = \frac{2v \sin\left(\frac{\theta}{2}\right)}{\left(\frac{r\theta}{v}\right)}$$

$$a_{\text{avg}} = \frac{2v^2 \sin\left(\frac{\theta}{2}\right)}{r\theta}$$

$$\text{Here, } \theta = \pi \text{ rad}$$

$$a_{\text{avg}} = \frac{2v^2 \sin\left(\frac{\pi}{2}\right)}{r \times \pi}$$

$$a_{\text{avg}} = \frac{2v^2}{\pi r}$$



42. An object of mass  $m$  moves with constant speed in a circular path of radius  $R$  under the action of a force of constant magnitude  $F$ . The kinetic energy of object is

(1)  $\frac{1}{2}FR$  (2)  $FR$  (3)  $2FR$  (4)  $\frac{1}{4}FR$

**Sol.** Answer (1)

$$\text{KE} = \frac{1}{2}mv^2 = \frac{1}{2} \frac{F}{a} \times v^2 = \frac{1}{2} \frac{F \times v^2}{\left(\frac{v^2}{R}\right)} = \frac{1}{2}FR$$

43. The angular speed of earth around its own axis is

- (1)  $\frac{\pi}{43200}$  rad/s      (2)  $\frac{\pi}{3600}$  rad/s      (3)  $\frac{\pi}{86400}$  rad/s      (4)  $\frac{\pi}{1800}$  rad/s

**Sol.** Answer (1)

$$\text{Angular speed} = \frac{2\pi}{T}$$

$T \rightarrow$  Time period of earth = 24 h

$$\omega = \frac{2\pi}{24 \times 60 \times 60} = \frac{\pi}{43200} \text{ rad s}^{-1}$$

44. A particle moves in a circle of radius 25 cm at two revolutions per second. The acceleration of the particle is (in  $\text{m/s}^2$ )

- (1)  $\pi^2$       (2)  $8\pi^2$       (3)  $4\pi^2$       (4)  $2\pi^2$

**Sol.** Answer (3)

$$a = r\omega^2$$

$$a = \frac{25}{100} (2 \times 2\pi)^2$$

$$a = 4\pi^2 \text{ m/s}^2$$

45. A particle is revolving in a circular path of radius 25 m with constant angular speed 12 rev/min. Then the angular acceleration of particle is

- (1)  $2\pi^2 \text{ rad/s}^2$       (2)  $4\pi^2 \text{ rad/s}^2$       (3)  $\pi^2 \text{ rad/s}^2$       (4) Zero

**Sol.** Answer (4)

Angular acceleration is the rate of change of angular speed or angular velocity if  $\omega$  remains constant then  $\alpha = 0$

46. Two particles are moving in circular paths of radii  $r_1$  and  $r_2$  with same angular speeds. Then the ratio of their centripetal acceleration is

- (1) 1 : 1      (2)  $r_1 : r_2$       (3)  $r_2 : r_1$       (4)  $r_2^2 : r_1^2$

**Sol.** Answer (2)

Centripetal acceleration is given by

$$a = \frac{v^2}{r} = r\omega^2$$

For same ' $\omega$ '

$$a_c \propto r \Rightarrow \frac{a_1}{a_2} = \frac{r_1}{r_2}$$

47. A particle  $P$  is moving in a circle of radius  $r$  with uniform speed  $v$ .  $C$  is the centre of the circle and  $AB$  is diameter. The angular velocity of  $P$  about  $A$  and  $C$  is in the ratio

- (1) 4 : 1      (2) 2 : 1      (3) 1 : 2      (4) 1 : 1


**Sol.** Answer (3)

$$\omega_{P/C} = \frac{d\theta}{dt}$$

$$\omega_{P/A} = \frac{1}{2} \frac{d\theta}{dt}$$

$$\omega_{P/A} = \frac{1}{2} \omega_{P/C}$$

$$\frac{\omega_{P/A}}{\omega_{P/C}} = \frac{1}{2} = 1:2$$



48. A car is moving at a speed of 40 m/s on a circular track of radius 400 m. This speed is increasing at the rate of 3 m/s<sup>2</sup>. The acceleration of car is

- (1) 4 m/s<sup>2</sup>                      (2) 7 m/s<sup>2</sup>                      (3) 5 m/s<sup>2</sup>                      (4) 3 m/s<sup>2</sup>

**Sol.** Answer (3)

$$v = 40 \text{ ms}^{-1}$$

$$r = 400 \text{ m}$$

$$a_T = 3 \text{ ms}^{-2}$$







$$a_c = \frac{v^2}{r} = \frac{40 \times 40}{400} = 4 \text{ ms}^{-2}$$

$$a = \sqrt{a_c^2 + a_T^2}$$


$$a = \sqrt{4^2 + 3^2} = 5 \text{ ms}^{-2}$$


$$a = 5 \text{ ms}^{-2}$$


49. Four particles A, B, C and D are moving with constant speed  $v$  each. At the instant shown relative velocity of A with respect to B, C and D are in directions


**Sol.** Answer (1)



(ii)   $\Rightarrow \vec{v}_{AB} = \vec{v}_A - \vec{v}_B \Rightarrow \vec{v}_A + (-\vec{v}_B)$

  $\vec{v}_A + (-\vec{v}_B)$

(iii)   $\Rightarrow \vec{v}_{AD} = \vec{v}_A - \vec{v}_D = \vec{v}_A + (-\vec{v}_D)$

  $\vec{v}_A - \vec{v}_D$

50. The ratio of angular speeds of minute hand and hour hand of a watch is

(1) 6 : 1

(2) 12 : 1

(3) 60 : 1

(4) 1 : 60

**Sol.** Answer (2)

$\omega_{mh}$  = Angular speed of minute hand

$\omega_{hh}$  = Angular speed of hour hand

$$\omega_{mh} = \frac{2\pi}{60 \text{ m}} = \frac{2\pi}{60 \times 60} \text{ rad s}^{-1}$$

$$\omega_{hh} = \frac{2\pi}{12 \text{ h}} = \frac{2\pi}{12 \times 60 \times 60} \text{ rad s}^{-1}$$

$$\frac{\omega_{mh}}{\omega_{hh}} = \frac{\frac{2\pi}{60 \times 60}}{\frac{2\pi}{12 \times 60 \times 60}} = \frac{1}{1} \times \frac{12}{1}$$

$$\omega_{mh} : \omega_{hh} \Rightarrow 12 : 1$$

51. If  $\theta$  is angle between the velocity and acceleration of a particle moving on a circular path with decreasing speed, then


(1)  $\theta = 90^\circ$

(2)  $0^\circ < \theta < 90^\circ$

(3)  $90^\circ < \theta < 180^\circ$

(4)  $0^\circ \leq \theta \leq 180^\circ$

**Sol.** Answer (3)



$\theta$  between  $v$  &  $Q$  is

$$90^\circ < \theta < 180^\circ$$

52. If speed of an object revolving in a circular path is doubled and angular speed is reduced to half of original value, then centripetal acceleration will become/remains

- (1) Same (2) Double (3) Half (4) Quadruple

**Sol.** Answer (1)

$$a_c = r\omega^2 = (r\omega)(\omega)$$

$$a_c = v\omega$$

$$a_c = (2v)\left(\frac{\omega}{2}\right) = v\omega = a_c$$

## SECTION - B

### Objective Type Questions

**(Motion in a Plane, Motion in a Plane with Constant Acceleration, Relative Velocity in Two Dimensions)**

1. Two particles A and B start moving with velocities 20 m/s and  $30\sqrt{2}$  m/s along x-axis and at an angle  $45^\circ$  with x-axis respectively in xy-plane from origin. The relative velocity of B w.r.t. A

- (1)  $(10\hat{i} + 30\hat{j})$  m/s (2)  $(30\hat{i} + 10\hat{j})$  m/s  
(3)  $(30\hat{i} - 20\sqrt{2}\hat{j})$  m/s (4)  $(30\sqrt{2}\hat{i} + 10\sqrt{2}\hat{j})$  m/s

**Sol.** Answer (1)


$$v_A = 20 \text{ m/s}$$

$$v_B = 30\sqrt{2} \text{ m/s along } 45^\circ \text{ with x-axis}$$

$$\vec{v}_B = v_B \cos 45^\circ \hat{i} + v_B \sin 45^\circ \hat{j} = 30\hat{i} + 30\hat{j}$$

$$\vec{v}_{BA} = \vec{v}_B - \vec{v}_A = 30\hat{i} + 30\hat{j} - 20\hat{i}$$

$$\boxed{\vec{v}_{BA} = 10\hat{i} + 30\hat{j}}$$



2. A particle is moving with constant speed  $v$  in xy plane as shown in figure. The magnitude of its angular velocity about point O is

- (1)  $\frac{v}{\sqrt{a^2 + b^2}}$  (2)  $\frac{v}{b}$  (3)  $\frac{vb}{(a^2 + b^2)}$  (4)  $\frac{v}{a}$


**Sol.** Answer (3)

$$v \sin \theta = r\omega$$

$$v \sin \theta = \sqrt{a^2 + b^2} \times \omega$$


$$\frac{v}{\sqrt{a^2 + b^2}} \frac{b}{\sqrt{a^2 + b^2}} = \omega$$

$$\boxed{\frac{vb}{(a^2 + b^2)} = \omega}$$



$$v_{\perp} = r\omega$$

3. A particle is moving eastwards with a speed of 6 m/s. After 6 s, the particle is found to be moving with same speed in a direction  $60^\circ$  north of east. The magnitude of average acceleration in this interval of time is

(1)  $6 \text{ m/s}^2$ (2)  $3 \text{ m/s}^2$ (3)  $1 \text{ m/s}^2$ 

(4) Zero


**Sol.** Answer (3)

$$|\Delta \vec{v}| = 2v \sin \frac{\theta}{2} = 2v \sin \left( \frac{60}{2} \right) = 2v \sin 30^\circ = v$$

$$|\Delta \vec{v}| = 6 \text{ ms}^{-1}$$


$$\Delta t = 6 \text{ s}$$

$$\text{so, } a_{av} = \frac{6}{6} = 1 \text{ ms}^{-2}$$



$$a_{av} = \frac{(\Delta \vec{v})}{\Delta t}$$

4. When a force  $F$  acts on a particle of mass  $m$ , the acceleration of particle becomes  $a$ . Now if two forces of magnitude  $3F$  and  $4F$  acts on the particle simultaneously as shown in figure, then the acceleration of the particle is

(1)  $a$ (2)  $2a$ (3)  $5a$ (4)  $8a$ **Sol.** Answer (3)

$$F_{\text{net}} = \sqrt{3^2 + 4^2} = 5F$$


$$\text{So, } F_{\text{net}} = ma$$

$$5F = ma'$$

$$\Rightarrow a' = \frac{5F}{m}$$

$$\boxed{a' = 5a}$$

$$\frac{F}{m} = a$$




5. Out of the two cars  $A$  and  $B$ , car  $A$  is moving towards east with a velocity of  $10 \text{ m/s}$  whereas  $B$  is moving towards north with a velocity  $20 \text{ m/s}$ , then velocity of  $A$  w.r.t.  $B$  is (nearly)

(1)  $30 \text{ m/s}$ (2)  $10 \text{ m/s}$ (3)  $22 \text{ m/s}$ (4)  $42 \text{ m/s}$ **Sol.** Answer (3)

$$\vec{v}_{AB} = \vec{v}_A - \vec{v}_B$$

$$v_{AB} = \sqrt{v_A^2 + v_B^2}$$

$$|\vec{v}_{AB}| = \sqrt{10^2 + 20^2} = \sqrt{100 + 400} = \sqrt{500} \approx 22 \text{ ms}^{-1}$$



6. A man moves in an open field such that after moving 10 m on a straight line, he makes a sharp turn of  $60^\circ$  to his left. The total displacement just at the start of 8<sup>th</sup> turn is equal to
- (1) 12 m (2) 15 m (3) 17.32 m (4) 14.14 m

**Sol.** Answer (3)



After 8 such turns object is at 'B'.

Displacement = AB

Two vectors are at  $60^\circ$





$$\sqrt{10^2 + 10^2 + 2 \times 10^2 \times \frac{1}{2}} = 10\sqrt{3} \text{ m}$$

$$\Rightarrow \boxed{17.32 \text{ m} = AB}$$



### (Projectile Motion)

7. A particle is projected at angle  $\theta$  with horizontal from ground. The slope ( $m$ ) of the trajectory of the particle varies with time ( $t$ ) as



**Sol.** Answer (1)

Slope of trajectory


$$\tan \phi = \frac{u \sin \theta - gt}{u \cos \theta}$$

$$\text{So, } m = \frac{u \sin \theta}{u \cos \theta} - \frac{gt}{u \cos \theta}$$

$$m = \tan \theta - \frac{g}{u \cos \theta} t$$

$$\Rightarrow y = a - bx$$

Therefore,



8. If  $H_1$  and  $H_2$  be the greatest heights of a projectile in two paths for a given value of range, then the horizontal range of projectile is given by

- (1)  $\frac{H_1 + H_2}{2}$  (2)  $\frac{H_1 + H_2}{4}$  (3)  $4\sqrt{H_1 H_2}$  (4)  $4[H_1 + H_2]$


**Sol.** Answer (3)

$$\theta_1 + \theta_2 = 90^\circ$$

$$H_1 = \frac{u^2 \sin^2 \theta_1}{2g}$$

$$H_2 = \frac{u^2 \sin^2 (90^\circ - \theta_1)}{2g}$$

$$H_1 H_2 = \frac{R^2}{16} \quad \therefore R = \frac{u^2 \sin^2 \theta_1}{g}$$



$$R = 4\sqrt{H_1 H_2}$$

9. If  $R$  and  $H$  are the horizontal range and maximum height attained by a projectile, then its speed of projection is

(1)  $\sqrt{2gR + \frac{4R^2}{gH}}$

(2)  $\sqrt{2gH + \frac{R^2 g}{8H}}$

(3)  $\sqrt{2gH + \frac{8H}{Rg}}$

(4)  $\sqrt{2gH + \frac{R^2}{H}}$

**Sol.** Answer (2)

$$H = \frac{u^2 \sin^2 \theta}{2g} \Rightarrow \sin \theta = \sqrt{\frac{2gH}{u^2}}$$

$$R = \frac{2u^2 \sin \theta \cos \theta}{g}$$

$$R = \frac{2u^2}{g} \sqrt{\frac{2gH}{u^2}} \times \sqrt{1 - \frac{2gH}{u^2}}$$

$$R = \frac{2u^2}{g} \sqrt{\frac{2gH}{u^2}} \times \sqrt{\frac{u^2 - 2gH}{u^2}}$$

$$\frac{gR}{2\sqrt{2gH}} = \sqrt{u^2 - 2gH}$$

Squaring both the sides,

$$\frac{gR^2}{4 \times 2gH} = u^2 - 2gH$$

$$\Rightarrow u^2 = 2gH + \frac{gR^2}{8H}$$

$$u = \sqrt{2gH + \frac{gR^2}{8H}}$$

10. A particle projected from ground moves at angle  $45^\circ$  with horizontal one second after projection and speed is minimum two seconds after the projection. The angle of projection of particle is [Neglect the effect of air resistance]

- (1)  $\tan^{-1}(3)$  (2)  $\tan^{-1}(2)$  (3)  $\tan^{-1}(\sqrt{2})$  (4)  $\tan^{-1}(4)$

**Sol.** Answer (2)

$$\theta = 45^\circ, t = 1 \text{ s}$$

$$\tan \phi = \frac{V_y}{U_y} = \frac{u \sin \theta - gt}{u \cos \theta}$$

$$\tan 45^\circ = \frac{u \sin \theta - g \times 1}{u \cos \theta} \Rightarrow u \cos \theta = u \sin \theta - g$$

also,  $V_y = 0$ , after 1<sup>st</sup> (as speed is minimum)

$$u \sin \theta - g \times 2 = 0 \Rightarrow \boxed{u \sin \theta = 2g} \quad \dots(i)$$

$$\text{so, } u \cos \theta = 2g - g$$

$$\boxed{u \cos \theta = g} \quad \dots(ii)$$

$$\text{so, } \frac{(i)}{(ii)} = \frac{u \sin \theta}{u \cos \theta} = \frac{2g}{g}$$


$$\Rightarrow \tan \theta = 2$$

$$\boxed{\theta = \tan^{-1}(2)}$$

11. A ball is projected from ground at an angle  $45^\circ$  with horizontal from distance  $d_1$  from the foot of a pole and just after touching the top of pole it falls on ground at distance  $d_2$  from pole on other side, the height of pole is

- (1)  $2\sqrt{d_1 d_2}$  (2)  $\frac{d_1 + d_2}{4}$  (3)  $\frac{2d_1 d_2}{d_1 + d_2}$  (4)  $\frac{d_1 d_2}{d_1 + d_2}$

**Sol.** Answer (4)



$$\tan \theta + \tan \beta = \tan \theta$$

$$\frac{y}{d_1} + \frac{y}{d_2} = \tan 45^\circ$$

$$y = \left( \frac{d_1 d_2}{d_1 + d_2} \right)$$

12. A particle is projected with speed  $u$  at angle  $\theta$  with horizontal from ground. If it is at same height from ground at time  $t_1$  and  $t_2$ , then its average velocity in time interval  $t_1$  to  $t_2$  is

- (1) Zero (2)  $u \sin \theta$  (3)  $u \cos \theta$  (4)  $\frac{1}{2}[u \cos \theta]$

**Sol.** Answer (3)

When projectile is at same height, average velocity =  $u \cos \theta$ .

13. A particle is projected from ground at an angle  $\theta$  with horizontal with speed  $u$ . The ratio of radius of curvature of its trajectory at point of projection to radius of curvature at maximum height is


- (1)  $\frac{1}{\sin^2 \theta \cos \theta}$  (2)  $\cos^2 \theta$  (3)  $\frac{1}{\sin^3 \theta}$  (4)  $\frac{1}{\cos^3 \theta}$

**Sol.** Answer (4)

At the point of projection

$$r_A = \frac{u^2}{g \cos \theta}$$

$$r_H = \frac{u^2 \cos^2 \theta}{g}$$



$$\text{Ratio, } \frac{r_A}{r_H} = \frac{\frac{u^2}{g \cos \theta}}{\frac{u^2 \cos^2 \theta}{g}} = \frac{1}{\cos^3 \theta} = \frac{r_A}{r_H}$$

14. An object of mass 10 kg is projected from ground with speed 40 m/s at an angle  $60^\circ$  with horizontal. The rate of change of momentum of object one second after projection in SI unit is [Take  $g = 9.8 \text{ m/s}^2$ ]

- (1) 73 (2) 98 (3) 176 (4) 140

**Sol.** Answer (2)

$$\text{Force} = \frac{\Delta p}{\Delta t}, \text{ force remains constant} = mg$$

$$\Rightarrow 10 \times 9.8 \Rightarrow 98 \text{ N}$$

At  $t = 1$ , particle is at its maximum height.

15. An object is projected from ground with speed 20 m/s at angle  $30^\circ$  with horizontal. Its centripetal acceleration one second after the projection is [Take  $g = 10 \text{ m/s}^2$ ]

- (1)  $10 \text{ m/s}^2$  (2) Zero (3)  $5 \text{ m/s}^2$  (4)  $12 \text{ m/s}^2$

**Sol.** Answer (1)

$$\text{Centripetal acceleration} = \frac{v^2}{r} = g = 10 \text{ ms}^2$$

16. A particle is moving on a circular path with constant speed  $v$ . It moves between two points A and B, which subtends an angle  $60^\circ$  at the centre of circle. The magnitude of change in its velocity and change in magnitude of its velocity during motion from A to B are respectively

- (1) Zero, Zero (2)  $v, 0$  (3)  $0, v$  (4)  $2v, v$

**Sol.** Answer (2)

$$\Delta v = 2v \sin \frac{\theta}{2}$$

$$= 2v \times \sin \left( \frac{60}{2} \right)$$

$$|\Delta v| = v$$

Change in magnitude of velocity = 0

17. What is the path followed by a moving body, on which a constant force acts in a direction other than initial velocity (i.e. excluding parallel and antiparallel direction)?

- (1) Straight line                      (2) Parabolic                      (3) Circular                      (4) Elliptical

**Sol.** Answer (2)

The path will be parabolic.

18. Two stones are thrown with same speed  $u$  at different angles from ground in air. If both stones have same range and height attained by them are  $h_1$  and  $h_2$ , then  $h_1 + h_2$  is equal to

- (1)  $\frac{u^2}{g}$                       (2)  $\frac{u^2}{2g}$                       (3)  $\frac{u^2}{3g}$                       (4)  $\frac{u^2}{4g}$

**Sol.** Answer (2)

If range is same then, one angle is  $\theta$  and other angle is  $(90 - \theta)$

$$\Rightarrow h_1 = \frac{u^2 \sin^2 \theta}{2g}, \quad h_2 = \frac{u^2 \sin^2 (90 - \theta)}{2g}$$

$$h_1 = \frac{u^2 \sin^2 \theta}{2g}, \quad h_2 = \frac{u^2 \cos^2 \theta}{2g}$$

$$\text{So, } h_1 + h_2 \Rightarrow \frac{u^2 \sin^2 \theta}{2g} + \frac{u^2 \cos^2 \theta}{2g} = \frac{u^2}{2g} (\sin^2 \theta + \cos^2 \theta)$$

$$\boxed{h_1 + h_2 = \frac{u^2}{2g}}$$

19. A projectile is projected with speed  $u$  at an angle  $\theta$  with the horizontal. The average velocity of the projectile between the instants it crosses the same level is

- (1)  $u \cos \theta$                       (2)  $u \sin \theta$                       (3)  $u \cot \theta$                       (4)  $u \tan \theta$

**Sol.** Answer (1)

20. A ball is thrown at an angle  $\theta$  with the horizontal. Its horizontal range is equal to its maximum height. This is possible only when the value of  $\tan \theta$  is

- (1) 4                      (2) 2                      (3) 1                      (4) 0.5

**Sol.** Answer (1)

$$\frac{H}{R} = \frac{1}{4} \tan \theta$$


$$\Rightarrow H = R, \text{ given,}$$

$$\boxed{\tan \theta = 4}$$

$$\Rightarrow \boxed{\theta = \tan^{-1}(4)}$$



21. A ball is projected from a point O as shown in figure. It will strike the ground after ( $g = 10 \text{ m/s}^2$ )



(1) 4 s

(2) 3 s

(3) 2 s

(4) 5 s

**Sol.** Answer (1)


$$s_y = u_y T + \frac{1}{2} a_y T^2$$

$$-60 = 10 \sin 30^\circ \cdot T - \frac{1}{2} g T^2$$


$$-60 = 5T - 5T^2$$

$$T^2 - T - 2 = 0$$

$$\boxed{T = 4 \text{ s}}$$



22. A particle is thrown with a velocity of  $u \text{ m/s}$ . It passes A and B as shown in figure at time  $t_1 = 1 \text{ s}$  and  $t_2 = 3 \text{ s}$ . The value of  $u$  is ( $g = 10 \text{ m/s}^2$ )



(1) 20 m/s

(2) 10 m/s

(3) 40 m/s

(4) 5 m/s

**Sol.** Answer (3)

$$t_1 + t_2 = \frac{2u \sin \theta}{g}$$

$$1 + 3 = \frac{2u \times \sin 30^\circ}{10}$$

$$20 \times 2 = u$$

$$\Rightarrow \boxed{u = 40 \text{ ms}^{-1}}$$

23. Which one of the following statements is not true about the motion of a projectile?

- (1) The time of flight of a projectile is proportional to the speed with which it is projected at a given angle of projection
- (2) The horizontal range of a projectile is proportional to the square root of the speed with which it is projected
- (3) For a given speed of projection, the angle of projection for maximum range is  $45^\circ$
- (4) At maximum height, the acceleration due to gravity is perpendicular to the velocity of the projectile

**Sol.** Answer (2)

$$R = \frac{u^2 \sin 2\theta}{g} \Rightarrow \boxed{R \propto u^2}$$

24. A projectile is thrown with speed  $40 \text{ ms}^{-1}$  at angle  $\theta$  from horizontal. It is found that projectile is at same height at 1 s and 3 s. What is the angle of projection?

(1)  $\tan^{-1}\left(\frac{1}{\sqrt{2}}\right)$       (2)  $\tan^{-1}\left(\frac{1}{\sqrt{3}}\right)$       (3)  $\tan^{-1}(\sqrt{3})$       (4)  $\tan^{-1}(\sqrt{2})$

**Sol.** Answer (2)

$$\tan \theta = \frac{v_y}{v_x}$$

$$\text{Also, } t_1 + t_2 = \frac{2u \sin \theta}{g}$$

$$4 = \frac{2 \times 40 \times \sin \theta}{10}$$

$$\sin \theta = \frac{1}{2} \Rightarrow \theta = 30^\circ$$

$$\text{So, } \tan \theta = \tan 30^\circ \Rightarrow \frac{1}{\sqrt{3}}$$

$$\theta = \tan^{-1}\left(\frac{1}{\sqrt{3}}\right)$$

**(Uniform Circular Motion)**

25. A particle is moving in xy-plane in a circular path with centre at origin. If at an instant the position of particle is given by  $\frac{1}{\sqrt{2}}(\hat{i} + \hat{j})$ , then velocity of particle is along

(1)  $\frac{1}{\sqrt{2}}(\hat{i} - \hat{j})$       (2)  $\frac{1}{\sqrt{2}}(\hat{j} - \hat{i})$       (3)  $\frac{1}{\sqrt{2}}(\hat{i} + \hat{j})$       (4) Either (1) or (2)

**Sol.** Answer (4)

$$\vec{r} = \frac{1}{2}\hat{i} + \frac{1}{\sqrt{2}}\hat{j}$$

$\vec{v} \cdot \vec{r} = 0$  as velocity is always tangential to the path

$$(v_x \hat{i} + v_y \hat{j}) \cdot \frac{1}{2}(\hat{i} + \hat{j}) = 0$$


$$v_x + v_y = 0 \quad \Rightarrow \quad \begin{cases} v_x = -v_y \\ \text{or} \\ v_y = -v_x \end{cases}$$

$$v = \sqrt{v_x^2 + v_y^2} \quad \Rightarrow \quad \sqrt{2}v_x = v \quad \Rightarrow \quad v_x = \frac{v}{\sqrt{2}}$$

$$v_y = -\frac{v}{\sqrt{2}}$$

$$\text{or } v_x = -\frac{v}{\sqrt{2}}, v_y = \frac{v}{\sqrt{2}}$$

$$\text{So, possible value of } v \Rightarrow v_x \hat{i} + v_y \hat{j} \Rightarrow \frac{v}{\sqrt{2}}\hat{i} - \frac{v}{\sqrt{2}}\hat{j} \text{ or } -\frac{v}{\sqrt{2}}\hat{i} + \frac{v}{\sqrt{2}}\hat{j}$$




26. Consider the two statements related to circular motion in usual notations

- A. In uniform circular motion  $\vec{\omega}$ ,  $\vec{v}$  and  $\vec{a}$  are always mutually perpendicular  
 B. In non-uniform circular motion,  $\vec{\omega}$ ,  $\vec{v}$  and  $\vec{a}$  are always mutually perpendicular

- (1) Both A and B are true  
 (2) Both A and B are false  
 (3) A is true but B is false  
 (4) A is false but B is true

**Sol.** Answer (3)



Only first statement is correct.

$\omega \rightarrow$  mutually perpendicular to  $v$  and  $a$ .

27. Which of the following quantities remains constant during uniform circular motion?

- (1) Centripetal acceleration                      (2) Velocity  
 (3) Momentum                                        (4) Speed

**Sol.** Answer (4)

Speed remains constant.

## SECTION - C

### Previous Years Questions

1. The  $x$  and  $y$  coordinates of the particle at any time are  $x = 5t - 2t^2$  and  $y = 10t$  respectively, where  $x$  and  $y$  are in meters and  $t$  in seconds. The acceleration of the particle at  $t = 2$  s is [NEET-2017]

- (1) 0  
 (2)  $5 \text{ m/s}^2$   
 (3)  $-4 \text{ m/s}^2$   
 (4)  $-8 \text{ m/s}^2$

**Sol.** Answer (3)

$$x = 5t - 2t^2 \qquad y = 10t$$

$$\frac{dx}{dt} = 5 - 4t \qquad \frac{dy}{dt} = 10$$


$$v_x = 5 - 4t \qquad v_y = 10$$

$$\frac{dv}{dt} x = -4 \qquad \frac{dv}{dt} y = 10$$

$$a_x = -4 \qquad a_y = 0$$

Acceleration of particle at  $t = 2$  s is  $= -4 \text{ m/s}^2$

2. In the given figure,  $a = 15 \text{ m/s}^2$  represents the total acceleration of a particle moving in the clockwise direction in a circle of radius  $R = 2.5 \text{ m}$  at a given instant of time. The speed of the particle is [NEET (Phase-2) 2016]



- (1) 4.5 m/s                      (2) 5.0 m/s                      (3) 5.7 m/s                      (4) 6.2 m/s

**Sol.** Answer (3)

$$a \cos 30^\circ = \frac{v^2}{r} \Rightarrow 15 \frac{\sqrt{3}}{2} = \frac{v^2}{2.5} \Rightarrow v = 5.7 \text{ m/s}$$

3. If the magnitude of sum of two vectors is equal to the magnitude of difference of the two vectors, the angle between these vectors is [NEET-2016]

- (1)  $180^\circ$                       (2)  $0^\circ$                       (3)  $90^\circ$                       (4)  $45^\circ$

**Sol.** Answer (3)

$$|\vec{A} + \vec{B}| = |\vec{A} - \vec{B}| \Rightarrow \cos \theta = 0 \Rightarrow \theta = 90^\circ$$

4. A particle moves so that its position vector is given by  $\vec{r} = \cos \omega t \hat{x} + \sin \omega t \hat{y}$ , where  $\omega$  is a constant. Which of the following is true? [NEET-2016]

- (1) Velocity is perpendicular to  $\vec{r}$  and acceleration is directed away from the origin  
 (2) Velocity and acceleration both are perpendicular to  $\vec{r}$   
 (3) Velocity and acceleration both are parallel to  $\vec{r}$   
 (4) Velocity is perpendicular to  $\vec{r}$  and acceleration is directed towards the origin

**Sol.** Answer (4)

$$\begin{aligned} \vec{r} &= \cos \omega t \hat{x} + \sin \omega t \hat{y}, \\ \vec{v} &= \frac{d\vec{r}}{dt} = -\omega \sin \omega t \hat{x} + \omega \cos \omega t \hat{y} \\ \vec{a} &= -\omega^2 \cos \omega t \hat{x} - \omega^2 \sin \omega t \hat{y} = -\omega^2 \vec{r} \end{aligned} \quad \Rightarrow \vec{v} \cdot \vec{r} = 0$$

5. A particle of mass 10 g moves along a circle of radius 6.4 cm with a constant tangential acceleration. What is the magnitude of this acceleration if the kinetic energy of the particle becomes equal to  $8 \times 10^{-4} \text{ J}$  by the end of the second revolution after the beginning of the motion? [NEET-2016]

- (1)  $0.2 \text{ m/s}^2$                       (2)  $0.1 \text{ m/s}^2$                       (3)  $0.15 \text{ m/s}^2$                       (4)  $0.18 \text{ m/s}^2$

**Sol.** Answer (2)

$$m = 0.01 \text{ kg}, \quad r = 6.4 \text{ cm}$$

$$\frac{1}{2}mv^2 = 8 \times 10^{-4} \text{ J}$$

$$v^2 = \frac{16 \times 10^{-4}}{0.01} = 16 \times 10^{-2}$$

$$\text{Speed } v^2 = 2a_t s$$

$$v^2 = 2a_t 4\pi r \Rightarrow a_t = \frac{v^2}{8\pi r} = \frac{16 \times 10^{-2}}{8 \times 3.14 \times 6.4 \times 10^{-2}} = 0.1 \text{ m/s}^2$$

6. The position vector of a particle  $\vec{R}$  as a function of time is given by  $\vec{R} = 4\sin(2\pi t)\hat{i} + 4\cos(2\pi t)\hat{j}$ , where  $R$  is in meters,  $t$  is in seconds and  $\hat{i}$  and  $\hat{j}$  denote unit vectors along  $x$ - and  $y$ -directions, respectively. Which one of the following statements is wrong for the motion of particle? [Re-AIPMT-2015]

- (1) Path of the particle is a circle of radius 4 meter
- (2) Acceleration vector is along  $-\vec{R}$
- (3) Magnitude of acceleration vector is  $\frac{v^2}{R}$ , where  $v$  is the velocity of particle
- (4) Magnitude of the velocity of particle is 8 meter/second

**Sol.** Answer (4)

$$\vec{R} = 4\sin(2\pi t)\hat{i} + 4\cos(2\pi t)\hat{j} = x\hat{i} + y\hat{j}$$

Now,  $x^2 + y^2 = 4^2$  which is equation of circle of radius  $R$ .

So, the motion is UCM with speed

$$V = 8\pi\sqrt{2} \text{ m/s}$$

7. Two particles  $A$  and  $B$ , move with constant velocities  $\vec{v}_1$  and  $\vec{v}_2$ . At the initial moment their position vectors are  $\vec{r}_1$  and  $\vec{r}_2$  respectively. The condition for particles  $A$  and  $B$  for their collision is [Re-AIPMT-2015]

- (1)  $\vec{r}_1 - \vec{r}_2 = \vec{v}_1 - \vec{v}_2$
- (2)  $\frac{\vec{r}_1 - \vec{r}_2}{|\vec{r}_1 - \vec{r}_2|} = \frac{\vec{v}_2 - \vec{v}_1}{|\vec{v}_2 - \vec{v}_1|}$
- (3)  $\vec{r}_1 \cdot \vec{v}_1 = \vec{r}_2 \cdot \vec{v}_2$
- (4)  $\vec{r}_1 \times \vec{v}_1 = \vec{r}_2 \times \vec{v}_2$

**Sol.** Answer (2)

For collision final positions should be equal

$$\Rightarrow \vec{r}_1 + \vec{v}_1 t = \vec{r}_2 + \vec{v}_2 t$$

$$\Rightarrow \vec{r}_1 - \vec{r}_2 = (\vec{v}_2 - \vec{v}_1)t$$

$$\Rightarrow \vec{r}_1 - \vec{r}_2 = (\vec{v}_2 - \vec{v}_1) \frac{|\vec{r}_1 - \vec{r}_2|}{|\vec{v}_2 - \vec{v}_1|}$$

8. A ship  $A$  is moving Westwards with a speed of  $10 \text{ km h}^{-1}$  and a ship  $B$   $100 \text{ km}$  South of  $A$ , is moving Northwards with a speed of  $10 \text{ km h}^{-1}$ . The time after which the distance between them becomes shortest, is [AIPMT-2015]

- (1)  $10\sqrt{2} \text{ h}$
- (2)  $0 \text{ h}$
- (3)  $5 \text{ h}$
- (4)  $5\sqrt{2} \text{ h}$

**Sol.** Answer (3)

9. A projectile is fired from the surface of the earth with a velocity of  $5 \text{ ms}^{-1}$  and angle  $\theta$  with the horizontal. Another projectile fired from another planet with a velocity of  $3 \text{ ms}^{-1}$  at the same angle follows a trajectory which is identical with the trajectory of the projectile fired from the earth. The value of the acceleration due to gravity on the planet is (in  $\text{ms}^{-2}$ ) is (Given  $g = 9.8 \text{ ms}^{-2}$ ) [AIPMT-2014]

- (1) 3.5
- (2) 5.9
- (3) 16.3
- (4) 110.8

**Sol.** Answer (1)

Since trajectory is same, so range and maximum height both will be identical from earth and planet. So equating maximum height (Answer can be obtained by equating range also)

$$\frac{u_e^2 \sin^2 \theta}{2g_e} = \frac{u_p^2 \sin^2 \theta}{2g_p}$$

$$\frac{2.5}{9.8} = \frac{9}{g_p} \Rightarrow g_p = 3.5 \text{ m/s}^2$$

10. A particle is moving such that its position coordinates  $(x, y)$  are

$(2 \text{ m}, 3 \text{ m})$  at time  $t = 0$ ,

$(6 \text{ m}, 7 \text{ m})$  at time  $t = 2 \text{ s}$  and

$(13 \text{ m}, 14 \text{ m})$  at time  $t = 5 \text{ s}$ .

Average velocity vector  $(\vec{v}_{av})$  from  $t = 0$  to  $t = 5 \text{ s}$  is

[AIPMT-2014]

(1)  $\frac{1}{5}(13\hat{i} + 14\hat{j})$

(2)  $\frac{7}{3}(\hat{i} + \hat{j})$


(3)  $2(\hat{i} + \hat{j})$

(4)  $\frac{11}{5}(\hat{i} + \hat{j})$

**Sol.** Answer (4)

11. The velocity of a projectile at the initial point  $A$  is  $(2\hat{i} + 3\hat{j}) \text{ m/s}$ . Its velocity (in  $\text{m/s}$ ) at point  $B$  is

[NEET-2013]



(1)  $-2\hat{i} + 3\hat{j}$

(2)  $2\hat{i} - 3\hat{j}$

(3)  $2\hat{i} + 3\hat{j}$

(4)  $-2\hat{i} - 3\hat{j}$

**Sol.** Answer (2)

The change is only in the  $y$ -component

So,  $\boxed{v_f = 2\hat{i} - 3\hat{j}}$

$\therefore a_x = 0$

12. The horizontal range and the maximum height of a projectile are equal. The angle of projection of the projectile is

[AIPMT (Prelims)-2012]

(1)  $\theta = \tan^{-1}(2)$

(2)  $\theta = 45^\circ$

(3)  $\theta = \tan^{-1}\left(\frac{1}{4}\right)$

(4)  $\theta = \tan^{-1}(4)$

**Sol.** Answer (4)

$H = R$

$\tan \theta = 4 \quad \left( \because \frac{H}{R} = \frac{1}{4} \tan \theta \right)$

$\boxed{\theta = \tan^{-1}(4)}$

13. A particle has initial velocity  $(2\vec{i} + 3\vec{j})$  and acceleration  $(0.3\vec{i} + 0.2\vec{j})$ . The magnitude of velocity after 10 s will be [AIPMT (Prelims)-2012]

(1) 5 units (2) 9 units (3)  $9\sqrt{2}$  units (4)  $5\sqrt{2}$  units

**Sol.** Answer (4)

$$\vec{u} = 3\hat{i} + 3\hat{j}, \quad \vec{a} = 0.3\hat{i} + 0.2\hat{j}$$

$$\Delta t = 10 \text{ s}$$

$$\vec{v} = \vec{u} + \vec{a}t$$

$$\begin{aligned}\vec{v} &= 2\hat{i} + 3\hat{j} + (0.3\hat{i} + 0.2\hat{j}) \times 10 \\ &= 2\hat{i} + 3\hat{j} + 3\hat{i} + 2\hat{j}\end{aligned}$$

$$\vec{v} = 5\hat{i} + 5\hat{j}$$

$$v = \sqrt{5^2 + 5^2} = \sqrt{50}$$

$$v \Rightarrow 5\sqrt{2} \text{ ms}^{-1}$$

14. A particle moves in a circle of radius 5 cm with constant speed and time period  $0.2\pi$  s. The acceleration of the particle is [AIPMT (Prelims)-2011]

(1)  $5 \text{ m/s}^2$  (2)  $15 \text{ m/s}^2$  (3)  $25 \text{ m/s}^2$  (4)  $36 \text{ m/s}^2$

**Sol.** Answer (1)

$$r = 5 \text{ cm}, v = ?, T = 0.2\pi \text{ s}$$

$$T = \frac{2\pi}{\omega} \Rightarrow \omega = \frac{20\pi}{0.2\pi} = 10 \text{ rad s}^{-1}$$

$$a = r\omega^2 = 5 \times 10^{-2} \times 100$$

$$a = 5 \text{ ms}^{-2}$$

15. A body is moving with velocity  $30 \text{ m/s}$  towards east. After  $10 \text{ s}$  its velocity becomes  $40 \text{ m/s}$  towards north. The average acceleration of the body is [AIPMT (Prelims)-2011]

(1)  $5 \text{ m/s}^2$  (2)  $1 \text{ m/s}^2$   
(3)  $7 \text{ m/s}^2$  (4)  $\sqrt{7} \text{ m/s}^2$


**Sol.** Answer (1)

$$a_{av} = \frac{|\Delta \vec{v}|}{\Delta t}$$

$$\Delta v = (\vec{v}_2 - \vec{v}_1) = \sqrt{v_2^2 + v_1^2} \quad (\because \theta = 90^\circ)$$

$$= 5\sqrt{30^2 + 40^2} = 50 \text{ ms}^{-1}$$

$$\text{so, } a_{av} = \frac{50}{10} \Rightarrow 5 \text{ ms}^{-2} = a_{av}$$



16. A missile is fired for maximum range with an initial velocity of 20 m/s. If  $g = 10 \text{ m/s}^2$ , the range of the missile is [AIPMT (Prelims)-2011]

(1) 20 m (2) 40 m (3) 50 m (4) 60 m

**Sol.** Answer (2)


For maximum range  $\theta = 45^\circ$

$$v = 20 \text{ ms}^{-1}$$

$$R = \frac{u^2}{g} = \frac{20 \times 20}{10} \quad [\because \theta = 45^\circ]$$

$$\boxed{R = 40 \text{ m}}$$

17. A particle of mass  $m$  is released from rest and follows a parabolic path as shown. Assuming that the displacement of the mass from the origin is small, which graph correctly depicts the position of the particle as a function of time? [AIPMT (Prelims)-2011]



**Sol.** Answer (2)

18. A projectile is fired at an angle of  $45^\circ$  with the horizontal. Elevation angle of the projectile at its highest point as seen from the point of projection is [AIPMT (Mains)-2011]

(1)  $\tan^{-1}\left(\frac{\sqrt{3}}{2}\right)$

(2)  $45^\circ$

(3)  $60^\circ$

(4)  $\tan^{-1}\frac{1}{2}$


**Sol.** Answer (4)

$$\frac{1}{4}\tan\theta = \frac{H}{R}$$

$$4H = R \quad \dots(i)$$


$$\text{In } \triangle ABC, \tan\phi = \frac{H}{\frac{R}{2}} = \frac{2H}{R}$$

$$\tan\phi = \frac{2H}{4H} = \frac{1}{2} \Rightarrow \phi = \tan^{-1}\left(\frac{1}{2}\right)$$






19. Six vectors,  $\vec{a}$  through  $\vec{f}$  have the magnitudes and directions indicated in the figure. Which of the following statements is true? [AIPMT (Prelims)-2010]



- (1)  $\vec{b} + \vec{c} = \vec{f}$       (2)  $\vec{d} + \vec{c} = \vec{f}$       (3)  $\vec{d} + \vec{e} = \vec{f}$       (4)  $\vec{b} + \vec{e} = \vec{f}$

**Sol.** Answer (3)



$$\vec{d} + \vec{e} = \vec{f}$$

20. The speed of a projectile at its maximum height is half of its initial speed. The angle of projection is

- (1)  $60^\circ$       (2)  $15^\circ$       (3)  $30^\circ$       (4)  $45^\circ$  [AIPMT (Mains)-2010]

**Sol.** Answer (1)

$$u_H = u \cos \theta$$

$$\frac{u}{2} = u \cdot \cos \theta \Rightarrow \cos \theta = \frac{1}{2}$$

$$\theta = 60^\circ$$

21. A particle moves in x-y plane according to rule  $x = a \sin \omega t$  and  $y = a \cos \omega t$ . The particle follows

- (1) An elliptical path  
(2) A circular path  
(3) A parabolic path  
(4) A straight line path inclined equally to x and y-axes [AIPMT (Mains)-2010]

**Sol.** Answer (2)

$$x = a \sin \omega t \Rightarrow x^2 = a^2 \sin^2 \omega t$$

$$y = a \cos \omega t \Rightarrow y^2 = a^2 \cos^2 \omega t$$

$$x^2 + y^2 = a^2 (\sin^2 \omega t + \cos^2 \omega t)$$

$$x^2 + y^2 = a^2 \rightarrow \text{equation of circle.}$$

22. A particle has initial velocity  $(3\hat{i} + 4\hat{j})$  and has acceleration  $(0.4\hat{i} + 0.3\hat{j})$ . Its speed after 10 s is

[AIPMT (Prelims)-2010]

- (1) 7 units                      (2)  $7\sqrt{2}$  units                      (3) 8.5 units                      (4) 10 units

Sol. Answer (2)

23. A particle of mass  $m$  is projected with velocity  $v$  making an angle of  $45^\circ$  with the horizontal. When the particle lands on the level ground the magnitude of the change in its momentum will be [AIPMT (Prelims)-2008]

- (1) Zero                      (2)  $2mv$                       (3)  $\frac{mv}{\sqrt{2}}$                       (4)  $mv\sqrt{2}$

Sol. Answer (4)

$$\Delta p = -2mv \sin \theta \hat{j}$$

$$|\Delta \vec{p}| = 2mv \sin \theta = 2mv \times \frac{1}{\sqrt{2}}$$

$$|\Delta \vec{p}| = \sqrt{2} mv$$


24. A particle starting from the origin  $(0, 0)$  moves in a straight line in the  $(x, y)$  plane. Its coordinates at a later time are  $(\sqrt{3}, 3)$ . The path of the particle makes with the  $x$ -axis an angle of [AIPMT (Prelims)-2007]

- (1)  $0^\circ$                       (2)  $30^\circ$                       (3)  $45^\circ$                       (4)  $60^\circ$

Sol. Answer (4)

$$\tan \theta = \frac{P}{B} = \frac{3}{\sqrt{3}}$$

$$\theta = 60^\circ$$



25.  $\vec{A}$  and  $\vec{B}$  are two vectors and  $\theta$  is the angle between them, if  $|\vec{A} \times \vec{B}| = \sqrt{3}(\vec{A} \cdot \vec{B})$ , the value of  $\theta$  is

[AIPMT (Prelims)-2007]

- (1)  $90^\circ$                       (2)  $60^\circ$                       (3)  $45^\circ$                       (4)  $30^\circ$

Sol. Answer (2)

26. For angles of projection of a projectile at angles  $(45^\circ - \theta)$  and  $(45^\circ + \theta)$ , the horizontal ranges described by the projectile are in the ratio of [AIPMT (Prelims)-2006]

- (1) 1 : 1  
(2) 2 : 3  
(3) 1 : 2  
(4) 2 : 1

Sol. Answer (1)

27. A car runs at a constant speed on a circular track of radius 100 m, taking 62.8 s for every circular lap. The average velocity and average speed for each circular lap respectively is [AIPMT (Prelims)-2006]

- (1) 0, 0 (2) 0, 10 m/s  
(3) 10 m/s, 10 m/s (4) 10 m/s, 0

**Sol.** Answer (2)

$$T = 62.8 \text{ s}$$

$$r = 100 \text{ m}$$

$$T = \frac{2\pi}{\omega}$$

$$\omega = \frac{2\pi}{T} = \frac{2 \times 3.14 \times 10}{62.8 \times 100}$$

$$\boxed{\omega = 0.1 \text{ rad s}^{-1}}$$


$$v = r\omega$$

$$v = 100 \times 0.1$$

$$\boxed{v = 10 \text{ ms}^{-1}}$$

$$\text{Average velocity} = 0$$

$$\text{Average speed} = 10 \text{ ms}^{-1}$$



28. The vectors  $\vec{A}$  and  $\vec{B}$  are such that:  $|\vec{A} + \vec{B}| = |\vec{A} - \vec{B}|$ . The angle between the two vectors is [AIPMT (Prelims)-2006]

- (1)  $90^\circ$  (2)  $60^\circ$   
(3)  $75^\circ$  (4)  $45^\circ$

**Sol.** Answer (1)

$$|\vec{A} + \vec{B}| = |\vec{A} - \vec{B}|$$

$$\sqrt{A^2 + B^2 + 2AB\cos\theta} = \sqrt{A^2 + B^2 - 2AB\cos\theta}$$

Squaring both the sides,

$$A^2 + B^2 + 2AB\cos\theta = A^2 + B^2 - 2AB\cos\theta$$

$$4AB\cos\theta = 0 \Rightarrow \cos\theta = 0 \Rightarrow \boxed{\theta = 90^\circ}$$

29. If a vector  $2\hat{i} + 3\hat{j} + 8\hat{k}$  is perpendicular to the vector  $4\hat{j} - 4\hat{i} + \alpha\hat{k}$ , then the value of  $\alpha$  is [AIPMT (Prelims)-2005]

- (1) -1 (2)  $\frac{1}{2}$   
(3)  $-\frac{1}{2}$  (4) 1

**Sol.** Answer (3)

30. A stone tied to the end of a string of 1 m long is whirled in a horizontal circle with a constant speed. If the stone makes 22 revolutions in 44 s, what is the magnitude and direction of acceleration of the stone?

[AIPMT (Prelims)-2005]

- (1)  $\frac{\pi^2}{4} \text{ ms}^{-2}$  and direction along the radius towards the centre  
 (2)  $\pi^2 \text{ ms}^{-2}$  and direction along the radius away from centre  
 (3)  $\pi^2 \text{ ms}^{-2}$  and direction along the radius towards the centre  
 (4)  $\pi^2 \text{ ms}^{-2}$  and direction along the tangent to the circle

Sol. Answer (3)

$$\omega = \frac{22 \times 2\pi}{44 \text{ s}} \Rightarrow \pi \text{ rad s}^{-1}$$

Centripetal acceleration,  $a = r\omega^2$ 

$$a = 1 \times \pi^2 \text{ ms}^{-2} \text{ along the radius towards the centre}$$

31. Two boys are standing at the ends A and B of a ground, where  $AB = a$ . The boy at B starts running in a direction perpendicular to AB with velocity  $v_1$ . The boy at A starts running simultaneously with velocity  $v$  and catches the other boy in a time  $t$ , where  $t$  is

[AIPMT (Prelims)-2005]

- (1)  $\frac{a}{\sqrt{v^2 + v_1^2}}$  (2)  $\frac{a}{\sqrt{v^2 - v_1^2}}$  (3)  $\frac{a}{(v - v_1)}$  (4)  $\frac{a}{(v + v_1)}$

Sol. Answer (2)

The distance travelled by body at B,

$$= \text{Speed} \times t$$

$$= v_1 t$$


So,  $BC = v_1 t$ , similarly,  $AC = vt$ Applying pythagoras in  $\triangle ABC$ ,

$$v^2 t^2 = v_1^2 t^2 + a^2$$

$$(v^2 - v_1^2) t^2 = a^2$$

$$t^2 = \frac{a^2}{v^2 - v_1^2}$$

$$t = \frac{a}{\sqrt{v^2 - v_1^2}}$$



32. If the angle between the vectors  $\vec{A}$  and  $\vec{B}$  is  $\theta$ , the value of the product  $(\vec{B} \times \vec{A}) \cdot \vec{A}$  is equal to

[AIPMT (Prelims)-2005]


- (1)  $BA^2 \cos \theta$  (2)  $BA^2 \sin \theta$   
 (3)  $BA^2 \sin \theta \cos \theta$  (4) Zero

Sol. Answer (4)

33. A boat is sent across a river with a velocity of 8 km/h. If the resultant velocity of the boat is 10 km/h, then velocity of the river is

(1) 8 km/h (2) 10 km/h (3) 12.8 km/h (4) 6 km/h

**Sol.** Answer (4)



$$v_r = \sqrt{v^2 + u^2}$$

$$v_r = 10 \text{ kmh}^{-1}, v = 8 \text{ kmh}^{-1}$$

$$u = ?$$

$$100 = 8^2 + u_R^2$$

$$\Rightarrow u_R^2 = 36 \Rightarrow \boxed{u_R = 6 \text{ km/h}}$$

34. Which of the following is correct relation between an arbitrary vector  $\vec{A}$  and null vector  $\vec{0}$ ?

(1)  $\vec{A} + \vec{0} + \vec{A} \times \vec{0} = \vec{A}$  (2)  $\vec{A} + \vec{0} + \vec{A} \times \vec{0} \neq \vec{A}$  (3)  $\vec{A} + \vec{0} + \vec{A} \times \vec{0} = \vec{0}$  (4) None of these

**Sol.** Answer (1)

Knowledge based.

35. An object is being thrown at a speed of 20 m/s in a direction  $45^\circ$  above the horizontal. The time taken by the object to return to the same level is

(1)  $20/g$  (2)  $20/g$  (3)  $20\sqrt{2}/g$  (4)  $20\sqrt{2}g$

**Sol.** Answer (3)

$$u = 20 \text{ ms}^{-1}$$

$$\theta = 45^\circ$$

$$T = \frac{2u \sin \theta}{g}$$

$$T = \frac{2u}{g} \times \frac{1}{\sqrt{2}} = \frac{\sqrt{2}}{10} u$$

$$T = \frac{2 \times 20}{g} \frac{1}{\sqrt{2}} \Rightarrow \boxed{\frac{20\sqrt{2}}{g} = T}$$

36. A body is whirled in a horizontal circle of radius 20 cm. It has an angular velocity of 10 rad/s. What is its linear velocity at any point on circular path?

(1) 20 m/s (2)  $\sqrt{2}$  m/s (3) 10 m/s (4) 2 m/s

**Sol.** Answer (4)

$$v = r\omega = 20 \times 10^{-2} \times 10$$

$$\boxed{v = 2 \text{ ms}^{-1}}$$


37. Identify the vector quantity among the following.

- (1) Distance (2) Angular momentum  
(3) Heat (4) Energy

**Sol.** Answer (2)

Angular momentum is an axial vector.

38. Two particles A and B are connected by a rigid rod AB. The rod slides along perpendicular rails as shown here. The velocity of A to the left is 10 m/s. What is the velocity of B when angle  $\alpha = 60^\circ$ ?




- (1) 10 m/s (2) 9.8 m/s (3) 5.8 m/s (4) 17.3 m/s

**Sol.** Answer (3)

$$v_A \cos 60^\circ = v_B \cos 30^\circ$$

$$10 \times \frac{1}{2} = v_B \times \frac{\sqrt{3}}{2}$$

$$v_B = \frac{10}{\sqrt{3}}$$



39. The speed of a boat is 5 km/h in still water. It crosses a river of width 1.0 km along the shortest possible path in 15 minutes. The velocity of the river water (in km/h) is

- (1) 3 (2) 1 (3) 4 (4) 5

**Sol.** Answer (1)

$$v = 5 \text{ kmh}^{-1}$$

$$d = 1.0 \text{ km}$$

$$t = 15 \text{ min}$$

40. Two racing cars of masses  $m_1$  and  $m_2$  are moving in circles of radii  $r_1$  and  $r_2$  respectively. Their speeds are such that each makes a complete circle in the same time  $t$ . The ratio of the angular speeds of the first to the second car is

- (1)  $r_1 : r_2$   
(2)  $m_1 : m_2$   
(3) 1 : 1  
(4)  $m_1 m_2 : r_1 r_2$

**Sol.** Answer (3)


If time is same then,

$$\omega_1 : \omega_2 \Rightarrow 1 : 1 \quad \left[ \because \omega = \frac{2\pi}{T} \right]$$

41. A person aiming to reach exactly opposite point on the bank of a stream is swimming with a speed of 0.5 m/s at an angle of  $120^\circ$  with the direction of flow of water. The speed of water in the stream is

- (1) 0.25 m/s (2) 0.5 m/s  
(3) 1.0 m/s (4) 0.433 m/s

**Sol.** Answer (1)



$$v \sin 30^\circ = u$$

$$0.5 \times \frac{1}{2} = u \Rightarrow \boxed{u = 0.25 \text{ ms}^{-1}}$$

42. Two projectiles of same mass and with same velocity are thrown at an angle  $60^\circ$  and  $30^\circ$  with the horizontal, then which will remain same

- (1) Time of flight (2) Range of projectile  
(3) Maximum height acquired (4) All of these

**Sol.** Answer (2)

Range is same for complimentary angles.

43. Two particles having mass  $M$  and  $m$  are moving in a circular path having radius  $R$  and  $r$ . If their time periods are same, then the ratio of their angular velocities will be

- (1)  $\frac{r}{R}$  (2)  $\frac{R}{r}$  (3) 1 (4)  $\sqrt{\frac{R}{r}}$

**Sol.** Answer (3)

44. If  $|\vec{A} + \vec{B}| = |\vec{A}| = |\vec{B}|$  then angle between  $A$  and  $B$  will be

- (1)  $90^\circ$  (2)  $120^\circ$   
(3)  $0^\circ$  (4)  $60^\circ$

**Sol.** Answer (2)

$$|\vec{A} + \vec{B}| = |\vec{A}| = |\vec{B}|$$

$$|\vec{A} + \vec{B}| = \sqrt{A^2 + B^2 + 2AB \cos \theta}$$

$$A^2 = A^2 + A^2 + 2A^2 \cos \theta$$

$$-\frac{1}{2} = \cos \theta \Rightarrow \boxed{\theta = 120^\circ}$$

45. A particle moves along a circle of radius  $\left(\frac{20}{\pi}\right)$  m with constant tangential acceleration. If the velocity of the particle is 80 m/s at the end of the second revolution after motion has begun, the tangential acceleration is
- (1) 40 m/s<sup>2</sup>
  - (2) 640π m/s<sup>2</sup>
  - (3) 160π m/s<sup>2</sup>
  - (4) 40π m/s<sup>2</sup>

**Sol.** Answer (1)

$$r = \frac{20}{\pi} \text{ m}$$

$$a_T \rightarrow \text{constant}$$

$$v = 80 \text{ ms}^{-1}, \theta = 4\pi \text{ rad}$$

$$v = r\omega$$

$$80 = \frac{20}{\pi} \cdot \omega \Rightarrow \boxed{\omega = 4\pi \text{ rad s}^{-1}}$$

$$\omega = 0,$$

$$\omega^2 = \omega_0^2 + 2\alpha\theta$$

$$4\pi \times 4\pi = 2 \times \alpha \times 4\pi$$

$$\Rightarrow \boxed{\alpha = 2\pi \text{ rad s}^{-2}}$$

$$a = r\alpha = \frac{20}{\pi} \times 2\pi$$

$$\boxed{a = 40 \text{ ms}^{-2}}$$

46. The vector sum of two forces is perpendicular to their vector differences. In that case, the forces
- (1) Are equal to each other
  - (2) Are equal to each other in magnitude
  - (3) Are not equal to each other in magnitude
  - (4) Cannot be predicted

**Sol.** Answer (2)

$$(\vec{A} + \vec{B}) \cdot (\vec{A} - \vec{B}) = 0$$

$$A^2 - B^2 - AB + BA = 0$$

$$A^2 = B^2 \Rightarrow \boxed{A = B}$$

$$\text{so, } \boxed{|\vec{A}| = |\vec{B}|}$$



47. A wheel has angular acceleration of  $3.0 \text{ rad/s}^2$  and an initial angular speed of  $2.00 \text{ rad/s}$ . In a time of  $2 \text{ s}$  it has rotated through an angle (in radian) of

- (1) 10 (2) 12  
(3) 4 (4) 6

**Sol.** Answer (1)

$$\alpha = 3 \text{ rad s}^{-2}$$

$$\omega_0 = 2 \text{ rad s}^{-1}$$

$$t = 2 \text{ s}$$

$$\omega = \omega_0 + \alpha t$$

$$\omega = 2 + 3 \times 2$$

$$\boxed{\omega = 8 \text{ rad s}^{-1}}$$

$$\omega^2 - \omega_0^2 = 2 \times \theta$$

$$64 - 4 = 2 \times 3 \times \theta$$

$$\frac{60}{6} = \theta \Rightarrow \boxed{\theta = 10 \text{ rad s}^{-1}}$$

48. A particle is moving such that its position coordinates  $(x, y)$  are

$(2 \text{ m}, 3 \text{ m})$  at time  $t = 0$ ,

$(6 \text{ m}, 7 \text{ m})$  at time  $t = 2 \text{ s}$  and

$(13 \text{ m}, 14 \text{ m})$  at time  $t = 5 \text{ s}$

Average velocity vector  $(\vec{V}_{av})$  from  $t = 0$  to  $t = 5 \text{ s}$  is

- (1)  $\frac{1}{5}(13\hat{i} + 14\hat{j})$  (2)  $\frac{7}{3}(\hat{i} + \hat{j})$  (3)  $2(\hat{i} + \hat{j})$  (4)  $\frac{11}{5}(\hat{i} + \hat{j})$

**Sol.** Answer (4)

$$\vec{V}_{av} = \frac{\vec{r}_f - \vec{r}_i}{\Delta t} = \frac{(13-2)\hat{i} + (14-3)\hat{j}}{(5-0)} = \frac{11}{5}(\hat{i} + \hat{j})$$

## SECTION - D

### Assertion - Reason Type Questions

1. A : If  $\vec{A} \perp \vec{B}$ , then  $|\vec{A} + \vec{B}| = |\vec{A} - \vec{B}|$ .

R : If  $\vec{A} \perp \vec{B}$ , then  $(\vec{A} + \vec{B})$  is perpendicular to  $\vec{A} - \vec{B}$ .

**Sol.** Answer (3)

2. A : The addition of two vectors  $\vec{P}$  and  $\vec{Q}$  is commutative.

R : By triangle law of vector addition we can prove  $\vec{P} + \vec{Q} = \vec{Q} + \vec{P}$ .

**Sol.** Answer (1)

3. A : A vector cannot be divided by other vector.  
R : A vector can be divided by a scalar.

**Sol.** Answer (2)

4. A : At the highest point the velocity of projectile is zero.  
R : At maximum height projectile comes to rest.

**Sol.** Answer (4)

5. A : Horizontal range of a projectile is always same for angle of projection  $\theta$  with horizontal or  $\theta$  with vertical.  
R : Horizontal range depends only on angle of projection.

**Sol.** Answer (4)

6. A : Horizontal motion of projectile without effect of air is uniform motion.  
R : Without air effect the horizontal acceleration of projectile is zero.

**Sol.** Answer (1)

7. A : Path of a projectile with respect to another projectile is straight line.  
R : Acceleration of a projectile with respect to another projectile is zero.

**Sol.** Answer (1)

8. A : In the case of ground to ground projection of a projectile from ground the angle of projection with horizontal is  $\theta = 30^\circ$ . There is no point on its path such that instantaneous velocity is normal to the initial velocity.  
R : Maximum deviation of the projectile is  $2\theta = 60^\circ$ .

**Sol.** Answer (1)

9. A : Three vectors having magnitudes 10, 10 and 25 cannot produce zero resultant.  
R : If three vectors are producing zero resultant, then sum of magnitude of any two is more than or equal to magnitude of third and difference is less than or equal to the magnitude of third.

**Sol.** Answer (1)

10. A : Uniform circular motion is accelerated motion still speed remains unchanged.  
R : Instantaneous velocity is always normal to instantaneous acceleration in uniform circular motion.

**Sol.** Answer (1)

11. A : When a body moves on a curved path with increasing speed, then angle between instantaneous velocity and acceleration is acute angle.  
R : When the speed is increasing, its tangential acceleration is in the direction of instantaneous velocity.

**Sol.** Answer (1)

12. A : A uniform circular motion have non uniform acceleration.  
R : The direction of acceleration of a particle in uniform circular motion changes continuously.

**Sol.** Answer (1)

13. A : Angular displacement is vector quantity only for small values.  
R : The direction of angular displacement is perpendicular to plane of rotation of object.

**Sol.** Answer (2)

