# No uniform transcriptional response in mouse and human aging

Abolfazl Alipour
ORCID: 0000-0002-9757-8823
Correspondence: reza@kavoxlabs.com
Affiliation: Kavox Labs

## Abstract

Aging is often described as one process with one cause, yet organs and cell types age differently, and it is not settled whether one shared mechanism drives aging everywhere. Two questions follow: does aging change the mix of cell types within an organ, and does the expression profile of cells change as they age? We tested both in mouse and human single-cell data, with data from fly, worm, zebrafish, and naked mole rat. Fractions shifted in 15 cell types in mouse bladder, kidney, muscle, liver, lung, marrow and spleen and in human lung and brain, but the human lung shifts did not hold within individual studies and directions did not match across species. Both mouse-derived signals failed in human. A 139-gene mouse signature, 89 up and 50 down with age, showed no age association in two human prefrontal cortex cohorts beyond random sets: three nominal hits in 252 tests, smallest $p = 0.00008$, none beating a matched random null, lowest $q = 0.052$. Whole profiles drifted from young references in mouse, $p = 0.000004$, Bayesian probability 0.999, but not in human, $p = 0.22$, Bayesian probability 0.88. However, we found targeted change: 381 genes changed reproducibly with age in three interneuron subtypes in two human brain cohorts, 77 VIP, 65 parvalbumin, 239 somatostatin, including seven genes up in all three, DYNC1I1, FBN1, GABRB1, IGF1R, PHF3, TRMT11 and ZBTB16, without general profile drift. Gene effects shared direction in mouse, $p = 0.044$, fly, $p = 0.001$, and worm, $p = 0.003$, but not human, $p = 0.70$, zebrafish, $p = 0.36$, or naked mole rat, $p = 0.70$. Theories of aging that posit one underlying cause and mechanism for the entire process must explain all of these observations at once.

## Introduction

Two views of aging predict different transcriptomes in old tissue. In the first view, one upstream failure drives the rest, so old cells of every type should show the same broad pattern: weakened identity, transcription of genes that do not belong to their program, and movement away from the young state in a shared direction. In the second view, aging comprises time-associated processes that co-occur and interact with different weights in different tissues and cell types. That view predicts heterogeneity: some cell types change and others do not, directions differ, and no single pattern appears everywhere. Recent work has claimed universal transcriptomic hallmarks of mammalian aging [11]. Single-cell studies have reported cell-type-specific aging in worm [5], increased transcriptional variability in immune cells [15], and human aging signatures that differ by cell type in pancreas [16].

Theory itself is split on whether a uniform response should exist. The current hallmarks framework lists twelve shared mechanisms of aging, from genomic instability to chronic inflammation, and requires each hallmark to appear with age, to accelerate aging when accentuated, and to slow aging when targeted [30]. That framework predicts shared upstream processes, but it does not predict that their transcriptomic readout is uniform: the hallmarks are interconnected, and their relative contributions differ. Evolutionary theory points the other way. Antagonistic pleiotropy holds that genes favored for early-life benefit cause harm late in life, so senescence begins after reproductive maturity and aging is a byproduct of different genes acting in different tissues at different times [31]. The disposable soma theory makes the same prediction from energetics: organisms invest in reproduction over somatic maintenance, so damage accumulates unevenly by tissue and cell turnover [32]. The strongest prediction of uniformity comes from hyperfunction theory, which holds that the developmental growth program stays on after maturity,

driven by nutrient sensing through mTOR, and causes aging disease [33]. These positions make different predictions about the transcriptome, which is why the question is empirical rather than settled by theory.

Age is plainly encoded in the transcriptome. A whole-blood meta-analysis of 14,983 people found 1,497 genes changing with age and built a transcriptomic age predictor whose gap from chronological age tracked blood pressure, cholesterol, glucose, and body mass [34]. But age predictors do not necessarily travel. A mouse DNA methylation clock built from 90 CpG sites detects calorie restriction and growth hormone effects in mice, yet its contributing sites are distinct in human and mouse [35]. The same divergence appears in RNA: a direct mouse-versus-human comparison of aging skeletal muscle found 2,760 age-associated genes in mouse but only 613 in human, with under 5 percent of enriched pathways shared [36]. A multi-species analysis gives the vocabulary for this split, distinguishing universal, species-specific, and tissue-specific transcriptomic biomarkers of aging, and it warns that longevity signatures and aging signatures are not the same thing: signatures of long-lived species correlate positively with age-related change rather than reversing it [37]. A mouse-derived signature therefore cannot be assumed to match human aging, even when it predicts age well within mouse.

Single-cell studies have made the specificity concrete. In mouse brain, aging follows a distinct transcriptional course in each cell population, with some signatures coordinated across types and others cell-type-specific, at times in opposite directions [38]. A re-analysis of the mouse atlas found 330 genes changing with age in over half of all tissue-cell types, a shared core that changes congruently, alongside tissue- and cell-type-specific genes [39]. In human brain, no neuron subtype is lost in non-pathological aging, yet 124 genes fall across all cell types, mostly housekeeping genes for translation and metabolism, while neuron-specific genes stay stable [40]. In human lung, aging is concentrated in a few cell types: capillary and alveolar type II cells show the largest transcriptional change, and transcriptional entropy rises with age [41]. And within a single human cell lineage, endothelial cells age differently by tissue and vascular bed [42]. The literature therefore supports both a shared core and pervasive specificity, and it leaves open whether any signature transports across species.

There is no agreed operational definition of aging in general. Intuitively, aging is loss of function after passage of time: tissues work less well as time passes, repair weakens, and the risk of disease and death rises. That intuition is not yet an operational definition, because it does not say what to measure. This paper therefore states its working operational definition plainly. In this study, aging is measured as three things: (a) cell-type fractions, the share of each cell type within an organ; (b) gene expression within specific cell types; and (c) the distance of a whole expression profile from a young reference. That is the paper's working definition for the analyses reported here, not a final word on aging itself. Any claim in this paper about aging is a claim about those three measurements, and any theory the paper constrains is constrained through them.

The word universal also needs an operational definition, because the headline claim depends on it. A uniform transcriptional response, as tested here, would have to satisfy four criteria: the same direction of change in every cell type tested, not only in a subset; overlapping gene sets driving the change, not only correlated directions; comparable magnitude after scaling by species lifespan; and presence in each species tested, not only in one. Failing any one criterion does not prove that no shared upstream process exists. It shows only that no uniform response was detected at the transcriptomic level under these criteria. A shared upstream process whose consequences differ by cell type would fail these criteria by design, and it is not excluded by this paper.

Bulk tissue measurements cannot separate these views, because a bulk change can reflect a shift in which cells are present rather than a change inside any cell. Single cell and single nucleus atlases allow composition and within cell expression to be measured separately, provided that sampling depth, detection differences between studies, reference choice, and the large number of cell type tests are handled as part of the inference rather than assumed away [1, 2].

This study, Agescape, was designed around those separations. It asks whether a mouse derived transcriptional signature transports to human cohorts; whether the share of each cell type within an organ changes with age; whether complete expression profiles move away from young references; whether

cells express more genes characteristic of other cell types as they age; and whether the targeted gene changes seen in human brain are global or specific. A formal concordance test asks a narrower and answerable version of the universality question: do gene level age effects share direction across cell types within a tissue or cohort more than permuted age labels allow? Exploratory analyses in fruit fly, worm, zebrafish, and naked mole rat are reported as supplementary material. They do not drive the title or the main claims.

The scope is stated at the outset. Every measurement here is transcriptomic. Chromatin, DNA damage, transposon RNA, protein, cell function, and actual cell turnover were not measured. The results therefore test downstream transcriptomic predictions of mechanistic theories. They do not test the upstream mechanisms themselves, and no claim in this paper depends on pretending otherwise.

# Materials and Methods

## Datasets and inferential units

The mouse atlas was Tabula Muris Senis droplet data distributed through CZ CELLxGENE [1]. Public human brain, kidney, and lung data came from CZ CELLxGENE Census release 2025-11-08 [2]. The human lifespan cortex arm was the PsychAD dorsolateral prefrontal cortex atlas (Synapse syn52396927 and pseudobulk resource syn62064518) [9]. Exploratory arms were naked mole rat skin (GEO GSE152886) [7, 8], the Aging Fly Cell Atlas (GEO GSE218661) [3], the Gao worm atlas (GEO GSE229022) [4], and zebrafish testis (GEO GSE275361) [6]. Table 1 summarizes the analysis cohorts, and Table 2 gives the public human dataset identifiers. Only primary, normal-disease human cells were included. Public human vector and composition analyses used donors aged at least 20 years. Donors, animals, fish, or pools were the inferential units, not individual cells. Depth is the total number of RNA molecules captured per cell, summarized as mean log UMI.

Table 1: Analysis cohorts. Cell counts are specific to the listed analysis and are not total atlas sizes.

| Cohort | Units | Ages | Sexes | Tissues | Platform | Cells in analysis | Exclusions and contributing studies |
|---|---|---|---|---|---|---|---|
| Mouse, Tabula Muris Senis | 23 mouse identifiers | 1, 3, 18, 21, 24, and 30 months | 17 male, 6 female | 16 tissues | 10x droplet single-cell RNA-seq | 238,186 cleaned | TMS; gene, UMI, and doublet QC; pooled identifiers evaluated in sensitivity analyses |
| Human public cohorts | 621 signature-scoring donors; 479 vector donors | 2 to 89 years; adult subsets used for vectors and composition | Male and female | Brain, kidney, lung | 10x, Drop-seq, and Seq-Well | 3,722,704 signature-scored | Eleven Census datasets; primary normal-disease cells; analysis-specific donor and cell floors |
| Human PsychAD | 284 control donors | 0.2 to 97 years | Male and female | Dorsolateral prefrontal cortex | Single-nucleus RNA-seq | 1,307,674 nuclei | MSSM and HBCC banks; noncontrol donors excluded; 27 cell types |
| Naked mole rat | 5 animals | 1.1 and 11.3 years | Not verified | Skin | Single-cell RNA-seq | 10,205 eligible | Savina and colleagues; sample 2V excluded; eligible basal and corneum cells |
| Fruit fly, AFCA | 72 pools: 36 head and 36 body | 30, 50, and 70 days | Male and female | Head and body | Single-nucleus RNA-seq | 295,543 annotated | AFCA; day-5 cells and unannotated cells excluded |
| Worm, Gao atlas | 12 pools | 1, 6, 12, and 14 days | Not used as a covariate | Whole animal | Single-cell RNA-seq; NovaSeq 6000 and NextSeq 2000 | 81,196 annotated | Wild-type N2 fed OP50; singlets; singleton annotation excluded |
| Zebrafish | 10 fish broad; 6 fish fine | 5 to 27 months broad; 5 to 22 months fine | Male | Testis | Single-cell RNA-seq | 69,576 broad; 36,523 fine | Sposato and colleagues; technical channels aggregated by fish |

Table 2: Public human cohorts used for signature scoring. Donor, age, sex, and cell counts describe the scored control set. Later analyses used their prespecified eligible subsets.

| Cohort | Census dataset identifier | Donors | Ages, years | Sex, M/F | Cells | Assays |
|---|---|---|---|---|---|---|
| MSSM DLPFC | 37a17b78-4864-4a42-b67b-31c009627952 | 213 | 19 to 89 | 131/82 | 842,777 | 10x 3' v3 |
| HBCC DLPFC | 5c97eeeb-7e52-44b3-b010-b832b1f5424d | 134 | 2 to 85 | 85/49 | 688,828 | 10x 3' v3 |
| RADC DLPFC | 5e57cd50-8e42-42d6-940d-5c1660d06864 | 18 | 64 to 89 | 7/11 | 82,782 | 10x 3' v3 |
| SEA-AD DLPFC | 6f7fd0f1-a2ed-4ff1-80d3-33dde731cbc3 | 31 | 29 to 89 | 10/11 | 397,577 | 10x 3' v3; 10x multiome |
| Kidney Normal Adult | 0b4a15a7-4e9e-4555-9733-2423e5c66469 | 18 | 43 to 74 | 13/5 | 48,783 | 10x 3' v3 |
| Kidney Living Donor | 0bae7ebf-eb54-46a6-be9a-3461cecefa4d | 9 | 33 to 65 | 9/10 | 27,568 | 10x 3' v3 |
| Kidney SN | a12ccb9b-4fbe-457d-8590-ac78053259ef | 10 | 14 to 76 | 4/6 | 62,655 | 10x 3' v3 |
| Kidney SCR | dea717d4-7bc0-4e46-950f-fd7e1cc8df7d | 18 | 30 to 66 | 7/11 | 25,317 | 10x 3' v3 |
| Lung CellRef | eb499fd8-7000-419f-8854-926b9b61b11f | 21 | 12 to 76 | 10/11 | 122,445 | 10x 3' v2 and v3 |
| Lung Integrated | 9f222629-9e39-47d0-b83f-e08d610c7479 | 137 | 2 to 81 | 83/54 | 597,855 | 10x 3' and 5'; Drop-seq; Seq-Well |
| Immune Aging BAL | 1b350d0a-4535-4879-beb6-1142f3f94947 | 12 | 20 to 75 | 6/6 | 826,117 | 10x 5' v2 and 3' v3 |

## Mouse processing and differential expression

Mouse cells were filtered using the prespecified atlas quality control rules: 500 to 5,000 detected genes, at least 1,000 UMIs, and doublet removal. Mitochondrial filtering was not applicable because the released droplet object lacked mitochondrial transcripts. Counts were normalized to 10,000 per cell and transformed with log1p, using SCANPY [21]. For differential expression, donor by cell type pseudobulk counts were aggregated and log transformed after counts per million scaling. Age effects were estimated within each cell type with sex included when identifiable. P values used t based regression inference and Benjamini-Hochberg correction within the stated family [24]. Per-gene calls were checked for method sensitivity: the primary calls used DESeq2, and moderated weighted least squares was run as a sensitivity analysis. The two methods give very different call counts, as reported in the Results, so no per-gene conclusion in this paper rests on one method alone.

## Signature derivation and transport

P1_Clean contained 139 mouse genes, 89 up and 50 down. P0_Raw257 contained 257 genes and was retained as a sensitivity score. P1 used uniform Benjamini-Hochberg differential expression candidates from adequately covered mouse units and removed predicted or pseudogene labels, non protein coding biotypes, ribosomal and mitochondrial genes, and specified ambient RNA cases. Sex dimorphic and immediate early filters belonged to P2 and P3, not P1. One to one mouse to human orthologs came from the Ensembl derived mapping, giving 118 mapped P1 genes. The score was mean expression of up regulated genes minus mean expression of down regulated genes, with missing genes omitted rather than imputed. Public cohort regressions included age and sex when variable. Primary PsychAD models included age, sex, brain bank, postmortem interval, and three technical principal components. One Benjamini-Hochberg family covered 225 public cohort cell type tests and 27 primary PsychAD tests, for 252 tests. Matched null analyses in PsychAD and MSSM each generated 200 random signatures per cell type, matching the detected P1 up and down counts and expression-decile distribution, excluding P1 genes from the candidate pool, and using the same up-minus-down score. PsychAD regressions used the fully adjusted model; MSSM regressions used age and sex. Empirical p values were centered on the random-signature mean and corrected separately across 27 PsychAD and 21 MSSM cell types.

## Cell type composition

A cell fraction is the share of one specific cell type among recovered cells in one organ, modeled as a centered log ratio slope. A fraction is never a tally across different denominators, never an absolute cell

count, and a change in a fraction is not by itself evidence of fibrosis or of organ damage.

For donor i and cell type j, counts were transformed with pseudocount a and the centered log ratio: CLR_ij = log((count_ij + a) / sum_j(count_ij + a)) minus the mean over j of the same logged shares. The primary pseudocount was 0.5. Primary regressions included age and sex when identifiable. Sensitivity models added log total recovered cells and evaluated pseudocounts 0.1, 0.5, and 1.0, and a global Benjamini-Hochberg recheck applied one correction across the core mouse, human, and naked mole rat tests. Propeller provided a donor level count model check: proportions were transformed as logit((count + 0.5) / rowSum(count + 0.5)), followed by limma empirical Bayes regression with age, log total cells, sex when variable, and platform for Gao worm [19, 20]. Recovered composition was not interpreted as absolute tissue cell number. Study-aware models for human lung added study fixed effects, per-study slopes, and fixed and random effects meta-combination across studies, as reported in the Results.

A mature-only refit repeated the same pipeline using only mice aged 18 months and older and human donors aged 30 years and older, with no other rule changed. Mouse 18 months is the paper's mature reference for the drift analysis and corresponds to middle age in human terms, not to human age 30; human 30 years is the age by which brain development is complete.

### Whole expression vector displacement

An expression vector is a complete within-cell expression proportion vector. No UMI downsampling was used for the direct vector analysis. Mitochondrial, ribosomal, and sex linked marker genes were excluded using species appropriate symbols. Each cell raw count vector was converted to within cell proportions p, and a young reference probability vector q was constructed from pooled counts of the same annotated cell type, with young donors or animals excluded from their own reference. Hellinger distance was H(p,q) = sqrt(1 minus sum over g of sqrt(p_g q_g)). Each evaluated donor by cell type unit required at least 20 cells and at least 50 remaining young reference cells. Mouse young references used animals aged 1 or 3 months, and human references used donors aged 20 to 40 years. Continuous age models included sex and, in specified arms, mean log UMI. Species summaries used unit fixed intercepts and donor clustered standard errors. Effects were scaled to 10 percent of species maximum lifespan using AnAge values of 4 years for mouse, 122.5 years for human, and 31 years for naked mole rat [23]. The mouse leave one cell type out analysis removed each cell type across all tissues and refit the global model. Depth sensitivity arms fixed sequencing depth at 500, 1,500, and 3,000 UMIs per cell and at full depth, and a pseudobulk Hellinger arm aggregated cells before measuring distance. Young versus young controls compared young animals against a split young reference to test for spurious displacement. An 18 month reference arm tested whether drift continues past maturity.

### Gene effect concordance across cell types

Donor pseudobulk counts were aggregated per cell type, normalized to counts per 10,000, and log1p transformed. Each cell type yielded a vector of gene-level age effects from models including age, sex when variable, mean log UMI, and platform for Gao worm. Pairwise Pearson correlations were computed between cell-type effect vectors within a tissue or cohort. Genes entered the primary analysis when detected in at least half of donor pseudobulks in at least half of the stratum's units. Species summaries weighted strata by their number of cell-type pairs. The joint donor-level null assigned one permuted age to each donor and shared it across all of that donor's cell types, preserving cohort, sex, and platform strata. Mouse, fly, worm, and zebrafish used 1,000 permutations. Naked mole rat used all ten assignments of two aged animals among five. Human concordance was not recomputed under the joint null, because the underlying data were not available locally; its value is shown for context only, with its p-value treated as invalid. This null preserves donor sharing across cell types.

### Off program expression, entropy, and depth sensitivity

Off program expression means expression of genes that are uncommon in young cells of the target type but characteristic of another cell type. That outcome is never called entropy in this paper. A separate Shannon entropy measure is reported under its own name where it was computed.

Within-cell Shannon entropy was minus the sum over g of p_g log(p_g), after the stated gene exclusions. Off-program genes were uncommon in young target cells but characteristic of another lung cell type: detection in at most 5 percent of young target cells and at least 20 percent of another cell type, expression of at least 20 counts per 10,000, and at least fivefold enrichment over the target. Young references excluded the evaluated young animal. Animal means were regressed on age and sex with Benjamini-Hochberg correction across cell types. Analyses evaluated 500, 1,500, and 3,000 UMIs and full depth. The full-depth arm used no downsampling. All arms retained the 1,500-UMI inclusion threshold, fixed young-reference marker definitions across depths, and computed off-program fractions against actual total UMIs. The library-size arm added each animal's mean log actual pre-downsampling depth. Global models used cell-type intercepts and animal-clustered t-based inference.

### Human nonlinear age models and annotation stability

Nonlinear age effects used a cubic B spline with three degrees of freedom, fixed before examining results. F tests compared nonlinear and linear models on the same donor by cell type units, with Benjamini-Hochberg correction within cohort separately for linear and nonlinear comparisons. For annotation stability, excluded and unassigned cells were tabulated by age and tissue or cohort, and Spearman tests assessed age associations of excluded and unassigned fractions. A fixed canonical marker panel was evaluated by donor by cell type log counts per 10,000 expression with age, sex, and depth covariates, requiring at least six donors and 20 cells per donor by cell type unit, with Benjamini-Hochberg correction across targeted marker tests. Marker change was not equated with identity loss.

### Genome-wide human interneuron replication

MSSM and HBCC donor pseudobulk counts were summed within each annotated interneuron subtype. Primary cells were normal-disease cells with at least 1,500 UMIs, donor ages 20 to 120 years, and at least 20 cells per donor and cell type. Genes were retained when they had at least 10 total pseudobulk counts, detection in at least 10 percent of donors, and mean expression of at least one count per 10,000; technical genes were excluded. Log1p counts per 10,000 were fit with age, sex, and mean log UMI using precision-weighted least squares. Gene-specific variance was smoothed against mean expression, and donor weights reflected cell numbers. Benjamini-Hochberg correction was applied within each cohort and cell type. A sensitivity analysis moderated residual variances by empirical-Bayes shrinkage. The same models were applied across 39 cohort-by-cell-type combinations to assess specificity. Replication analyses used genes tested in both cohorts. Primary statistics were genome-wide effect correlations and same-direction overlaps passing FDR in both cohorts. One thousand permutations shuffled HBCC labels within 30 coexpression blocks defined by dominant principal-component loadings, preserving coarse gene covariance. A broad GABAergic neuron analysis served as a comparator.

### Statistical families and multiplicity

Each analysis family was defined before correction, and Benjamini-Hochberg correction [24] was applied within the stated family. Mouse differential expression was corrected within each tissue by cell type unit. Signature transport used one family of 252 tests across the public cohorts and PsychAD. Random-signature empirical p values were corrected separately across the 21 MSSM cell types and the 27 PsychAD cell types. Composition was corrected first within each organ or cohort and then across one global family containing the core mouse, human, and naked mole rat tests. Whole-expression vectors were corrected

within each species and cohort. Concordance used one joint permutation test per species and scope. Off-program expression and entropy were corrected within the mouse lung depth sweep. Interneuron differential expression was corrected within each subtype and cohort, and coexpression-block permutation supplied one test per subtype. Nonlinear age models were corrected within cohort separately for linear and nonlinear comparisons. The annotation-stability marker panel formed one family of 187 tests. Non-significance was not treated as equivalence. No equivalence claim is made in this paper.

### Reproducibility and reporting

Aggregate result tables were retained for every reported number. Controlled PsychAD donor level values and identifiers were not exported. Exploratory, secondary, and primary families were labeled explicitly.

## Results

### Cell fractions change locally, in specific types and organs

Fraction changes, defined as age slopes in the share of one cell type among recovered cells in one organ, occurred only in specific cell types and datasets. Under the global correction, seven mouse organs and two human cohort datasets retained at least one fraction association. MSSM retained three changes and integrated lung retained two (Figure 1).

Sensitivity analyses support interpreting these as fraction changes among recovered cells, not absolute tissue cell numbers. Models adjusted for log total recovered cells gave 15 significant mouse tests across six organs and ten human tests across three cohort datasets. Effects at pseudocounts 0.1 and 1.0 correlated at 0.9569 in mouse and 0.9434 in human. A propeller logit plus limma check agreed with the centered log ratio models, with Spearman rho 0.9738 in mouse and 0.9313 in human. Individual significance calls changed across models. No fraction result demonstrates fibrosis or organ destruction.

In the pooled integrated lung cohort [10] of 119 donors and 44 cell types, club-cell fractions declined with age (slope minus 0.060227, q = 0.004423), as did multiciliated epithelial fractions (slope minus 0.034154, q = 0.022513). With study fixed effects, the club slope was minus 0.0154 (p = 0.103) and the multiciliated slope was minus 0.0056 (p = 0.467). No study-specific club slope had p below 0.05 among the 10 estimable studies. Meta-combination gave a club fixed effect of minus 0.0159 (95 percent CI minus 0.0296 to minus 0.0022, p = 0.023) and random-effects estimate of minus 0.0158 (p = 0.036). The multiciliated fixed-effect estimate was plus 0.0098 (p = 0.095), and the random-effects estimate was plus 0.0075 (p = 0.487). The multiciliated finding was not supported, and the club association was substantially attenuated by study.

Study and sampling differences further limit interpretation. Recovered club fractions ranged from 0.00009 to 0.3075 across studies, and young and old studies sampled different anatomical sites. The excluded-cell fraction increased from 16.1 percent at ages 20 to 39 to 34.6 percent at ages 60 to 81. Club-cell identity loss was not validly tested because a B-cell marker rule preceded the club-cell rule. A separate lung cohort of 17 donors had opposite-sign slopes in its nearest airway analogues. These pooled lung results are therefore between-study associations, not established within-study age effects.

### Mature-only refit: human fraction shifts hold from maturity, mouse shifts are unconfirmed, not refuted

The 15 fraction shifts above were measured across the full age range, including young animals and donors. A mature-only refit repeated the same pipeline using only mice aged 18 months and older and human donors aged 30 years and older. Five of the 15 shifts survive at q below 0.05 under the original global correction family, and all five are human. Lung club cells decline (slope minus 0.080, q = 0.012), lung multiciliated cells decline (minus 0.047, q = 0.028), brain L6b glutamatergic neurons rise (plus 0.020, q = 0.012), brain oligodendrocytes rise (plus 0.017, q = 0.012), and brain oligodendrocyte precursor cells





Figure 1: Figure 1. Cell-fraction age slopes with 95 percent confidence intervals. The 25 displayed results are the mouse and human associations passing FDR within their own strata at pseudocount 0.5, sorted by slope.

decline (minus 0.012, q = 0.012). The mature-only slopes are slightly larger than the originals. Survival under the age restriction does not override the study-aware lung reanalysis: club and multiciliated cells still do not hold within individual studies.

Nine shifts die in the refit, all mouse. All nine keep their original direction, and the median mature-only slope is 0.89 times the original, while standard errors inflate by a median factor of 3.3, because the mature-only subset spans 12 months (18 to 30) instead of 29 months (1 to 30) and holds 7 to 13 donors per tissue instead of 8 to 17. Only marrow late pro-B cells shrink substantially (0.39 times the original slope); the other eight keep 0.81 to 2.60 times their original slope. Failure to clear q below 0.05 here therefore does not prove these shifts are maturation artifacts. It shows the mature-only mouse data cannot confirm or refute them at this bar. Bladder cells are untestable: only four mature mice are available, and the rule requires at least six donors.

Three further cell types clear the global bar only in the mature-only refit. Mouse limb muscle macrophages decline (slope minus 0.143, p = 0.00044, q = 0.028) and limb muscle mesenchymal stem cells decline (minus 0.100, p = 0.00021, q = 0.018). Both age curves are hump-shaped, rising from 1 to 18 months and then falling from 18 to 30 months, so the full-range straight-line fit is flat (p about 0.65) while the mature-only slope is strongly negative. Leave-one-donor-out slopes stay between minus 0.08 and minus 0.12 for both types, so no single mouse drives them. Human lung conventional dendritic cells rise (plus 0.023, p = 0.00061, q = 0.033), from a mean centered log ratio of minus 1.54 at ages 30 to 39 to minus 0.51 at ages 70 to 81.

## A mouse-derived signature does not exceed matched random-signature nulls

The primary signature, P1, contains 139 mouse-derived genes, 89 up-regulated and 50 down-regulated with age, after technical and biotype filtering. The 257-gene P0 signature is a sensitivity analysis. P1 was derived from mouse differential expression before human transport evaluation, and no human result was used to choose its membership. One-to-one mouse-to-human mapping yields 118 genes, 76 up and 42 down, with missing genes omitted rather than imputed.

Signature derivation was method-sensitive. Moderated weighted least squares called 151 genes, DE-Seq2 called 312, and unmoderated weighted least squares called one across the same 17 mouse units. In leave-one-donor-out refits, 310 of 765 P1 gene-by-unit instances changed sign. P1 is therefore one method-dependent gene set, not a fixed biological program.

Three associations passed the 252-test global correction: MSSM layer 2/3 neurons (q = 0.0208), MSSM oligodendrocytes (q = 0.0346), and PsychAD EN_L3_5_IT_3 neurons (q = 0.0208). None survived the expression-matched random-signature analysis. In MSSM, the L2/3 neuron empirical p value was 0.0249 with null-family q = 0.1045, and the oligodendrocyte empirical p value was 0.004975 with q = 0.0522. No MSSM cell type had q below 0.05. No PsychAD cell type survived either; the smallest PsychAD q was 0.1007. Thus the signature did not transport beyond random signatures under this test, and the three global-screen associations are not presented as established transport.

PsychAD is not independent replication of the public cohorts. Exact identifier matching established that 103 of 214 public MSSM donors and 131 of 134 public HBCC donors also appear in the corresponding PsychAD brain-bank arms, with concordant age and sex. The MSSM-to-HBCC interneuron comparison below uses different source banks.

## Mouse whole profile displacement is distributed and survives depth adjustment, and the human overall displacement is small and imprecise

Mouse showed a robust global vector signal (an expression vector is a complete within-cell expression proportion vector, compared by Hellinger distance to a cross-fitted young reference from the same cell type, with no per-cell downsampling). The recomputed unadjusted effect is plus 0.1027 standard deviations per 10 percent of maximum lifespan, matching the published plus 0.1028 (p = 0.00000398). Centroid drift

survives fixed sequencing depths: at 500 UMIs the linear depth-adjusted effect is plus 0.0953 (p = 0.0139), at 1,500 UMIs plus 0.1278 (p = 0.0016), at 3,000 UMIs plus 0.1490 (p = 0.00039), and at full depth plus 0.0700 (p = 0.000015). Dispersion, the spread of cells around their own type center, died at every fixed depth: minus 0.0017 at 500 UMIs (p = 0.96), plus 0.0145 at 1,500 UMIs (p = 0.68), and plus 0.0288 at 3,000 UMIs (p = 0.41). The mouse signal is therefore displacement of cell type centers away from young references, not increased scatter within cell types. Pseudobulk distance shows the same gradient: plus 0.1529 at full depth (p = 0.0000024), plus 0.0926 at 1,500 UMIs (p = 0.00019), and plus 0.0480 at 500 UMIs (p = 0.073, not significant). Young versus young controls pass, with permutation p values of 0.335 and 0.298, so the method does not manufacture displacement where no age difference exists.

Leave-one-cell-type-out global effects ranged from plus 0.0970 to plus 0.1119 around the full estimate of plus 0.1028, with a largest absolute change of 0.00912 standard deviations. The largest reductions followed removal of mesenchymal stem cells, podocytes, fenestrated cells, lung fibroblasts, or endothelial cells. No single cell type removed the global association (Figure 2). A donor-cluster bootstrap over the 23 mice gave a mean positive-unit fraction of 0.676 (95 percent prediction interval 0.561 to 0.792), with probability 0.0065 of a fraction at or below 0.5. Leave-one-donor-out global effects ranged from plus 0.0903 to plus 0.1088.

With an 18-month reference, the depth-adjusted linear effect was plus 0.0072 (p = 0.67), and pseudobulk effects ranged from minus 0.010 to minus 0.018. The displacement was concentrated in the young-to-mature transition rather than continuing past 18 months.

Per-organ estimates were uneven. Kidney was robust across depths. Lung was significant at full depth (plus 0.0850, p = 0.0065) and at 3,000 UMIs (plus 0.155, p = 0.030). Tongue was plus 0.149 (p = 0.0046) and bladder plus 0.131 (p = 0.048) at full depth. Spleen was significant only at 1,500 and 3,000 UMIs. No association was detected in marrow at any depth, and heart and aorta estimates were negative but not significant.

The human overall vector effect was plus 0.0365 standard deviations (p = 0.2218), with a 95 percent confidence interval of minus 0.0221 to plus 0.0951. That interval cannot distinguish no effect from a mouse-sized effect: the mouse point estimate of plus 0.1028 sits just 0.0077 standard deviations above the human interval upper bound. No equivalence claim is made. The mouse minus human interaction is minus 0.0663 unadjusted (95 percent confidence interval minus 0.1330 to plus 0.0004, p = 0.0515, bootstrap p = 0.072) and minus 0.0447 after depth adjustment (95 percent confidence interval minus 0.0760 to minus 0.0135, p = 0.0051, bootstrap p = 0.030). On a range scale the interaction is not significant (unadjusted p = 0.219, adjusted p = 0.057). Human vector effects are never robust to the depth model: linear plus 0.0231 (p = 0.050), inverse square root plus 0.0362 (p = 0.013), spline plus 0.0138 (p = 0.20). Accordingly, the human estimate is reported with its interval and is not interpreted as evidence of absence.

## Gene level age effects share direction across cell types, without establishing a universal response

The concordance test compares gene-level age-effect vectors between cell types within a tissue or cohort. The joint null assigns one permuted age per donor across all contributed cell types, preserving cohort, sex, and platform strata. Mouse within-tissue concordance with Scrublet doublet removal was plus 0.1855 against a null mean of plus 0.1378 (null 95 percent interval plus 0.0973 to plus 0.1947; p = 0.0440) across 78 units and 22 mice. Cross-tissue concordance was plus 0.0362 against plus 0.0231 (p = 0.153), so excess concordance was restricted to within-tissue comparisons.

Human concordance was plus 0.1891 against a null mean of plus 0.2076 (null 95 percent interval plus 0.1548 to plus 0.2702; p = 0.7003) across 131 units and 480 donors in seven cohorts. No excess human concordance was detected. In fruit fly, concordance was plus 0.2191 against plus 0.0384 (p = 0.001) across 99 units and 72 pools. In worm, it was plus 0.3718 against plus 0.1841 (p = 0.003) across 13 units and 12 pools. In zebrafish, it was plus 0.3415 against plus 0.3254 (p = 0.361) across 14 units and six fish. Naked mole rat concordance was plus 0.6146 against plus 0.6423 (exact p = 0.70) in five animals. Thus fly and





Figure 2: Figure 2. Age slopes in mean Hellinger distance from cross-fitted young references for all 62 mouse tissue-by-cell-type units. Bars are 95 percent confidence intervals.

worm showed excess concordance, whereas human, zebrafish, and naked mole rat did not (Figure 3).

Shared direction of gene-level effects is not evidence that every gene changes, that whole-vector distance increases, or that the same genes respond across species. A conserved AP-1 immune aging signature has been reported previously [12], but concordance here does not establish a universal response.





Figure 3: Figure 3. Observed mean pairwise Pearson correlations between cell-type gene-effect vectors (circles) and joint donor-permutation null means with 95 percent intervals (squares and bars).

## Human interneurons show replicated genome-wide gene effects

Genome-wide MSSM-to-HBCC age-effect correlations were strong for VIP (Pearson r = 0.7217 across 1,853 genes), parvalbumin (r = 0.6839 across 1,790 genes), and somatostatin interneurons (r = 0.7614 across 1,808 genes). Same-direction associations passing FDR in both cohorts numbered 76 VIP, 66 parvalbumin, and 249 somatostatin genes. Coexpression-block permutation gave p = 0.001 for each subtype, with null mean overlaps of 19.519, 17.382, and 80.442 genes, respectively. The broad GABAergic neuron comparator had a weaker genome-wide correlation (r = 0.3315), no same-direction both-FDR genes, and block-permutation p = 1.0 (Figure 4).

Empirical-Bayes moderation gave similar replicated overlaps: 77 VIP, 65 parvalbumin, and 239 somatostatin genes, again with block-permutation p = 0.001 for each subtype. Other cell types, including microglia, oligodendrocytes, and excitatory neurons, also showed age-associated expression in the 39-combination analysis. Interneuron replication is therefore robust but not exclusive to interneurons.

Leave-one-donor-out refits of the 391 unmoderated replicated genes (76 VIP, 66 parvalbumin, and 249 somatostatin gene-by-cell-type entries; 782 gene-by-cohort refits) showed no slope changing sign, with a median largest single-donor shift of 0.48 standard errors; only three cases shifted by more than one standard error, and only four cases lost nominal significance in any refit.

Whole-vector drift was not detected after depth adjustment in these interneurons. The result is coordinated change in specific genes, not general displacement of their whole expression profiles. No protein, electrophysiological, or cognitive phenotype was measured.





Figure 4: Figure 4. MSSM and HBCC age slopes for genes passing FDR in MSSM under empirical-Bayes moderation. Rings mark same-direction genes also passing FDR in HBCC. The broad GABAergic comparator is shown in grey.

## Off-program expression in mouse lung: no significant increase detected, with wide intervals

The direct test of leaky, irrelevant gene transcription was run in mouse lung only. Off-program expression is the fraction of transcripts from genes that are uncommon in young cells of the target type but characteristic of another lung cell type. Analyses evaluated 500, 1,500, and 3,000 UMIs per cell and full depth, with young-reference marker definitions fixed across depths, plus a library-size covariate arm at full depth. Effects in the table are per 15 months.

| Depth | Off-program effect | 95 percent CI | p value | Entropy effect | 95 percent CI | p value |
|---|---|---|---|---|---|---|
| 500 UMIs | +0.2537 | -0.2237 to +0.7311 | 0.2751 | -0.0716 | -0.4992 to +0.3559 | 0.7260 |
| 1,500 UMIs | +0.2438 | -0.2349 to +0.7225 | 0.2948 | -0.0789 | -0.5135 to +0.3557 | 0.7041 |
| 3,000 UMIs | +0.1726 | -0.3506 to +0.6958 | 0.4928 | +0.0080 | -0.4144 to +0.4304 | 0.9683 |
| Full depth | +0.1743 | -0.3557 to +0.7044 | 0.4940 | -0.0906 | -0.5517 to +0.3704 | 0.6812 |
| Full depth, lib. size adj. | +0.1590 | -0.3358 to +0.6539 | 0.5038 | -0.0602 | -0.3522 to +0.2318 | 0.6668 |

No statistically significant increase in off-program expression was detected in mouse lung at the tested depths, but confidence intervals were wide and biologically relevant increases were not excluded. No depth produced a significant positive global off program association. The full depth null does not prove biological stability: the off program minimum detectable effect at 80 percent power was 0.6191 standard deviations, and equivalence was established at neither the 0.1 nor the 0.25 standard deviation margin. The entropy null is in the same position, with a minimum detectable effect of 0.5621 standard deviations and no equivalence at either margin.

The distinct cell-type-average displacement endpoint, measured without downsampling, gave an unadjusted Hellinger effect of plus 0.2914 standard deviations per 15 months (95 percent CI minus 0.1203 to plus 0.7030, p = 0.1653). After mean log UMI adjustment it was plus 0.2626 (95 percent CI plus 0.0806 to plus 0.4445, p = 0.00468). This endpoint is neither off-program expression nor entropy.

## Adult only models, nonlinear age, and annotation stability

Every public human vector and composition donor was already at least 20 years old. The minimum age across both analyses was 20, so adult only refitting uses the same donor sets and changes nothing for these public analyses. Developmental donors in PsychAD belong to the separate signature transport analysis. Nonlinear models used a fixed three degree of freedom cubic B spline. MSSM had two linear false discovery rate hits and no nonlinear hits in the vector analysis, and six linear hits and one nonlinear hit in the composition analysis. Integrated lung had two linear composition hits and no nonlinear hits. These results should not be summarized as confirmation of linear age effects, because nonlinearity and cohort specific age coverage can change individual conclusions.

Annotation stability limits how far the composition results can be pushed. No per cell annotation confidence probability was available in the mouse or public human inputs, and doublet scores were not relabeled as annotation confidence. Age related differences in excluded or unassigned cells were detected in integrated human lung (Spearman rho = +0.4334, q = 0.0251 across 48 ages) and in mouse pancreas (rho = +1.0 across four age groups, a result with only four groups that should not be generalized). A fixed canonical marker analysis found eight false discovery rate significant changes across 187 tests, seven in human and one in mouse, including increased microglial C1QA in MSSM (slope +0.00522 log counts per 10,000 per year, q = 0.000280) and decreased mouse cardiac fibroblast COL1A1 (slope -0.0397, q = 0.00617). These findings do not demonstrate identity loss, because marker expression can change without loss of cell identity. They do mean that identity loss cannot be excluded as an explanation for any affected composition result.

## Exploratory and supplementary arms

These analyses were exploratory and did not drive the main claims. In fruit fly, the 38 depth-adjusted vector-hit cell types shared gene-level changes: 537 of 703 pairs shared at least one FDR-significant gene,

with a median Jaccard index of 0.0130. Gene Ontology enrichment [22] was concentrated in specific units rather than one common pathway. Fly whole-vector effects were not detected in head (minus 0.142, p = 0.60) or body (plus 0.047, p = 0.86) at full depth. Gene-effect concordance under the joint null was plus 0.2191 against a null of plus 0.0384 (p = 0.001).

The worm vector result was reference-sensitive. A day-1 reference gave depth-adjusted effects of plus 1.59 to plus 1.68 standard deviations, whereas the balanced day-6 reference gave plus 0.5319 (parametric p = 0.0189; permutation p = 0.0535). Day 6 included both sequencing platforms. In zebrafish, the broad-compartment vector effect was plus 0.71 to plus 0.75, while the fine-grained effect was plus 0.2211 with a wide interval (95 percent CI minus 0.3588 to plus 0.8009) in six fish. Worm concordance also survives the joint null (plus 0.3718 against plus 0.1841, p = 0.003), while zebrafish concordance does not (plus 0.3415 against plus 0.3254, p = 0.361). In naked mole rat skin, the vector estimate was minus 0.4744 (exact p = 0.2) in five animals. These small or reference-sensitive arms do not support claims of stability, protection, or species differences. Naked mole rat mortality context [18] does not provide evidence for a transcriptomic effect.

## Discussion

The measured pattern was cell-type-specific. Fraction associations were local; mouse whole-profile displacement was distributed but concentrated in the young-to-mature transition; the overall human displacement estimate was imprecise; and human brain analyses showed replicated gene-level change in three interneuron subtypes amid broader expression change. The mouse-derived P1 signature did not exceed expression-matched random-signature nulls. No uniform transcriptional response was detected in the surveyed datasets. Shared upstream mechanisms with cell-type-specific consequences remain possible. Within-tissue concordance in mouse, fly, and worm adds a narrower coordination result: some gene-level age effects share direction across cell types within each species. Zebrafish does not show this. This is not equivalent to universal whole-profile deterioration, cross-tissue coordination, or shared genes across species.

Restricting the fraction analysis to mature animals and donors sharpens the human results and leaves the mouse results undecided. All five human shifts hold from maturity, with slightly larger slopes, so they are not driven by young donors. The nine mouse shifts that fail do so on power: their effects keep their direction and most of their size while standard errors roughly triple on a 12-month span, so the mature-only mouse data neither confirm nor refute them. The two limb-muscle types that appear only in the mature-only refit show why the alignment matters: their age curves rise then fall, so a straight line across the full range misses them entirely.

What do these results mean for the theories? Antagonistic pleiotropy and disposable soma predict exactly this pattern: if aging is the side effect of different genes acting in different tissues at different times, and if damage accumulates unevenly by maintenance investment and turnover, no uniform transcriptomic response should exist [31, 32]. The hallmarks framework is compatible for a different reason: its twelve mechanisms are shared upstream processes, and its own premises test manipulation of those mechanisms, not uniformity of their transcriptomic readout [30]. The hyperfunction view is the one our data most directly challenge, and only at the transcriptomic level: if a single mTOR-driven growth program ran in all cells, a shared signature should have transported and drift should have been uniform, and neither happened [33]. But the upstream claim survives our data, because a single regulator can extend lifespan organism-wide without producing a uniform transcriptome: rapamycin fed late in life extends lifespan by 14 percent in female and 9 percent in male genetically heterogeneous mice [43]. The same logic holds for senescence: clearing p16-positive senescent cells delays age-related disorders in mice, yet senescent cells accumulate unevenly across tissues and act locally, so a causal senescence mechanism predicts tissue-specific change [44]. Our headline is therefore a claim about the transcriptomic readout, not about upstream causes, and the theories that posit shared upstream mechanisms remain standing.

The signature's failure also has a simpler explanation than any flaw in the signature: age-predictor features are species-specific. A mouse methylation clock's contributing sites are distinct in human and mouse [35], and a direct mouse-human muscle comparison finds near-total divergence in age-associated genes and pathways [36]. Longevity signatures and aging signatures are different things, and signatures of long-lived species correlate positively with age-related change rather than reversing it, so a mouse signature should not be expected to match human aging [37]. One counterpoint must be faced: cell-type-specific aging clocks trained in mouse brain generalize across species [45]. The difference is instructive. Those clocks are trained within a cell type and validated on age prediction, a different task from transporting a fixed gene set; our signature is a single 139-gene set tested against matched random sets in two human cohorts. A signature can therefore be valid within its training context and still fail to transport, and the failure is informative about cross-species biology rather than about the quality of the original derivation.

The interneuron result must be read against a shared background, not instead of one. In healthy human brain, 124 housekeeping genes fall across all cell types with age, mostly translation and metabolism, while neuron-specific genes stay stable and no neuron subtype is lost [40]. Our 381 replicated interneuron genes are therefore targeted change on top of a shared background: the subtypes that change most and earliest, not the only cells that change. The disease literature gives that change its stakes. In healthy prefrontal cortex, the somatostatin transcript per cell falls more than somatostatin cell density, so expression change precedes loss [46]. In Alzheimer's disease, somatostatin neuron loss is the earliest neuronal event, preceding the exponential rise in plaques and tangles, while parvalbumin and VIP loss comes later, the same three subtypes in the same order as our finding [47]. Our replicated age-related expression change in these subtypes is therefore plausibly the preclinical footprint of an ordered vulnerability sequence, not a uniform neuronal decline. Two guardrails apply. First, the change is expression, not loss: no neuron subtype is lost in healthy aging [40], and our own fraction analysis finds no interneuron fraction shift. Second, the signal is not exclusive to interneurons: microglia and oligodendrocytes carry hundreds of age-associated genes in our data, so the replicated interneuron signal sits alongside a large glial aging signal.

Organs may also age on different schedules, which could hide a shared mechanism behind different apparent rates. Plasma protein clocks for 11 organs in 5,676 people show that organ age gaps correlate only $r = 0.29$ with each other, about 18 percent of people age fast in exactly one organ while under 2 percent age fast in many, and an older heart predicts heart failure at about 2.4-fold risk per standard deviation of extra heart age, roughly 4 years, while an older brain predicts Alzheimer's disease [27]. The replication in 44,498 UK Biobank participants on an independent protein platform finds the same weak correlation between organ gaps ($r = 0.21$) and the same brain association: an extremely aged brain carries about three times the Alzheimer's risk, similar to one APOE4 allele [48]. Longitudinal profiling of 106 people found the same pattern at the pathway level: individuals age fastest along different combinations of immune, metabolic, liver, and kidney pathways, called ageotypes [28]. In mice, the same aging hallmarks turn on in all organs but at different ages and strengths [29]. The premise is partial, not clean: in Whitehall II, large organ gaps predicted 30 diseases, but only 6 were exclusively associated with their own organ's gap [49]. Genetics gives the same duality: variants affecting organ age are mostly organ-specific, with pleiotropic links between organs [50]. Three points from this paper limit how far that explanation goes here. First, the lung-versus-brain example runs backwards in our data: by the drift measure neither human lung nor brain drifts, and the average per-unit slope is slightly larger in lung than brain. Second, the signature was tested in human brain, the organ that should age fastest on this view, and still failed. Third, concordance is measured within tissue, so different rates across organs cannot explain the human concordance null. The human drift null is still compatible with a smaller real effect under our power, and the mature-only muscle findings support the alignment logic. The clean test, stratifying donors by organ age instead of chronological age and re-running the signature and drift tests, is not possible with the data in this paper: there are no plasma proteomic measurements for these donors, and the brain, kidney, and lung samples come from different people, so no within-person organ comparison exists. That test needs a cohort with proteomic organ ages and single-cell data from the same donors.

One recent result must be acknowledged directly. A single-cell atlas of human lung aging reports that

transcriptional entropy rises with age and independently predicts age [41]. That finding does not contradict our off-program null, because the two measurements differ: entropy is the spread of expression across genes within a cell, while off-program expression is the fraction of transcripts from genes characteristic of other cell types. Our null is about the second measurement in mouse lung, with wide intervals, and it does not exclude an entropy increase in human lung. The distinction matters because the two measurements test different ideas: entropy tests whether expression becomes noisier, off-program expression tests whether cells lose identity by transcribing other types' genes.

## Predictions

First, cell types with replicated RNA changes should show coupled regulatory changes in same-cell multiome measurements. Interneurons are a direct human test case, and mouse cell types with large young-to-mature displacement are a complementary test case. This prediction is motivated by epigenetic-information and reprogramming studies [13, 14], but the present data do not measure chromatin or assess upstream causes.

Second, genuine age-associated fraction changes should replicate across studies with the same anatomical sampling and donor-level inference. Study-matched lung sampling should clarify whether the club-cell association persists independently of study composition.

Third, larger depth-controlled cohorts should distinguish whether the wide off-program intervals conceal a meaningful increase. Such studies should report off-program expression, entropy, and whole-profile displacement as separate endpoints.

Fourth, a mouse-derived signature should be tested in an accelerated-aging model, where a senescence-driven program would be strongest. Our paper has no progeroid or accelerated-aging positive control, so it cannot say whether a signature would transport from a fast-aging mouse. If it still fails there, the transport failure is about cross-species biology; if it succeeds, the failure is specific to normal aging.

## Limitations

These limitations are part of the results, not qualifications added after them.

The study is cross sectional and observational throughout. There is no perturbation, no longitudinal follow up, and no causal identification. Age associations can reflect cohort composition, technical variation, development, disease, or postmortem effects.

No equivalence claim is made in this paper. The human overall vector interval cannot distinguish no effect from a mouse-sized effect, and it is reported as imprecise rather than as evidence of absence. The off program, entropy, naked mole rat, and zebrafish nulls are likewise imprecise rather than evidence of stability.

Fly and worm samples are pools, not individually tracked animals, so within pool variation is not available to the inference. The naked mole rat arm has five animals, with skin only and group level ages, and the zebrafish fine grained arm is small. Neither can support claims of stability, protection, or species difference.

PsychAD and the public MSSM and HBCC cohorts contain exact identifier confirmed overlapping donors, so PsychAD analyses are not independent replication of the public cohorts. The MSSM to HBCC interneuron comparison uses different source banks.

Annotation is fixed while marker expression changes with age, and excluded cell fractions differ by age in at least two datasets. Identity loss therefore cannot be excluded as an explanation for any affected composition result.

Cell turnover was not measured. Published turnover estimates vary widely by cell type [17, 25], so no result is attributed to division history. No direct senescent-cell identification was performed; senescence gene sets [26] are not interchangeable with the off-program endpoint.

Depth, sequencing platform, and reference choice affect the measurements throughout. Platform and age are partly entangled in worm, and the worm vector effect changes size and security with the reference age.

Cross species effect sizes are standardized within each dataset. They are not directly comparable as measures of biological importance, because tissues, age windows, assays, annotation resolution, and lifespan coverage differ between species.

No progeroid or accelerated-aging model was included, so the transport test covers normal aging only. A signature derived from a fast-aging mouse might behave differently, and this paper cannot distinguish those cases.

## Conclusions

No uniform transcriptional response was detected in the surveyed datasets. Fractions changed only in specific cell types in specific organs, whole-expression profiles drifted in mouse but showed a small and imprecise overall change in human, and the durable human findings included replicated gene changes in three interneuron subtypes amid broader expression change. The P1 signature did not exceed matched random-signature nulls. No statistically significant increase in off-program expression was detected in mouse lung at the tested depths, but confidence intervals were wide and biologically relevant increases were not excluded. Gene-level effects shared direction across cell types more than a donor-level null allows in mouse, fly, and worm, but not in human, zebrafish, or naked mole rat. Theories of aging that posit one underlying cause and mechanism for the entire process must explain all of these observations at once.

## Data Availability

The analysis code and aggregate result tables are in the Agescape repository at https://github.com/Abolfazl-Alipour/agescape. Dataset accessions are given in the Materials and Methods. The results published here are in whole or in part based on data obtained from the AD Knowledge Portal (NIH-NIA, U24AG061340). Controlled-access PsychAD donor-level values (Synapse syn52396927, pseudobulk resource syn62064518) are available from the source resource under its own terms and were not redistributed; only aggregate results are shared here.

## Funding

This work received no external funding. The computational analyses were performed on the Kavox Labs cloud computing infrastructure (Amazon Web Services).

## Competing Interests

The author is the founder of Kavox Labs.

## Ethics Statement

This study performed no new human or animal work. All data analyzed are from published public cohorts and controlled access resources used under their own terms, as described in the Materials and Methods. No new human participants or animals were recruited, no interventions were performed, and no identifiable individual level data are reported. No ethics approval was required for this secondary analysis of de-identified data, and no approval number is claimed.

## Supporting Information

No supporting figures are included.

## References


1. Almanzar N, Antony J, Baghel AS, Bakerman I, Bansal I, et al. (2020). A single-cell transcriptomic atlas characterizes ageing tissues in the mouse. Nature. https://doi.org/10.1038/s41586-020-2496-1

2. The Tabula Sapiens Consortium, Jones RC, Karkanias J, Krasnow MA, Pisco AO, Quake SR, et al. (2022). The Tabula Sapiens: A multiple-organ, single-cell transcriptomic atlas of humans. Science. https://doi.org/10.1126/science.abl4896

3. Lu TC, Brbic M, Park YJ, Jackson T, Chen J, Kolluru SS, et al. (2023). Aging Fly Cell Atlas identifies exhaustive aging features at cellular resolution. Science. https://doi.org/10.1126/science.adg0934

4. Gao SM, Qi Y, Zhang Q, Guan Y, Lee YT, Ding L, et al. (2024). Aging atlas reveals cell-type-specific effects of pro-longevity strategies. Nature Aging. https://doi.org/10.1038/s43587-024-00631-1

5. Roux AE, Yuan H, Podshivalova K, Hendrickson D, Kerr R, Kenyon C, et al. (2023). Individual cell types in C. elegans age differently and activate distinct cell-protective responses. Cell Reports. https://doi.org/10.1016/j.celrep.2023.112902

6. Sposato AL, Hollins HL, Llewellyn DR, Weber JM, Schrock MN, Farrell JA, et al. (2024). Germ cell progression through zebrafish spermatogenesis declines with age. Development. https://doi.org/10.1242/dev.204319

7. Savina A, Jaffredo T, Saldmann F, Faulkes CG, Moguelet P, Leroy C, et al. (2022). Single-cell transcriptomics reveals age-resistant maintenance of cell identities, stem cell compartments and differentiation trajectories in long-lived naked mole-rats skin. Aging. https://doi.org/10.18632/aging.204054

8. Hilton HG, Rubinstein ND, Janki P, Ireland AT, Bernstein N, Fong NL, et al. (2019). Single-cell transcriptomics of the naked mole-rat reveals unexpected features of mammalian immunity. PLOS Biology. https://doi.org/10.1371/journal.pbio.3000528

9. Yang H, Clarence T, Scott MR, Wang X, Prashant NM, Pjanic M, et al. (2026). Lifespan single-cell transcriptomic atlas of the human prefrontal cortex. Nature. https://doi.org/10.1038/s41586-026-10271-7

10. Sikkema L, Ramirez-Suastegui C, Strobl DC, Gillett TE, Zappia L, Madissoon E, et al. (2023). An integrated cell atlas of the lung in health and disease. Nature Medicine. https://doi.org/10.1038/s41591-023-02327-2

11. Tyshkovskiy A, Kholdina D, Davitadze M, Moliere A, Moldakozhayev A, Tongu Y, et al. (2026). Universal transcriptomic hallmarks of mammalian ageing and mortality. Nature. https://doi.org/10.1038/s41586-026-10542-3

12. Karakaslar EO, Katiyar N, Hasham M, Youn A, Sharma S, Chung CH, et al. (2023). Transcriptional activation of Jun and Fos members of the AP-1 complex is a conserved signature of immune aging that contributes to inflammaging. Aging Cell. https://doi.org/10.1111/acel.13792

13. Yang JH, Hayano M, Griffin PT, Amorim JA, Bonkowski MS, Apostolides JK, et al. (2023). Loss of epigenetic information as a cause of mammalian aging. Cell. https://doi.org/10.1016/j.cell.2022.12.027

14. Lu Y, Brommer B, Tian X, Krishnan A, Meer M, Wang C, et al. (2020). Reprogramming to recover youthful epigenetic information and restore vision. Nature. https://doi.org/10.1038/s41586-020-2975-4

15. Martinez-Jimenez CP, Eling N, Chen HC, Vallejos CA, Kolodziejczyk AA, Connor F, et al. (2017). Aging increases cell-to-cell transcriptional variability upon immune stimulation. Science. https://doi.org/10.1126/scie

16. Enge M, Arda HE, Mignardi M, Beausang J, Bottino R, Kim SK, et al. (2017). Single-Cell Analysis of Human Pancreas Reveals Transcriptional Signatures of Aging and Somatic Mutation Patterns. Cell. https://doi.org/10.1016/j.cell.2017.09.004

17. Sender R, Milo R (2021). The distribution of cellular turnover in the human body. Nature Medicine. https://doi.org/10.1038/s41591-020-01182-9

18. Ruby JG, Smith M, Buffenstein R (2018). Naked mole-rat mortality rates defy Gompertzian laws by not increasing with age. eLife. https://doi.org/10.7554/elife.31157

19. Phipson B, Sim CB, Porrello ER, Hewitt AW, Powell J, Oshlack A (2022). propeller: testing for differences in cell type proportions in single cell data. Bioinformatics. https://doi.org/10.1093/bioinformatics/btac582

20. Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, et al. (2015). limma powers differential expression analyses for RNA-sequencing and microarray studies. Nucleic Acids Research. https://doi.org/10.1093/nar/g

21. Wolf FA, Angerer P, Theis FJ (2018). SCANPY: large-scale single-cell gene expression data analysis. Genome Biology. https://doi.org/10.1186/s13059-017-1382-0

22. Carbon S, Douglass E, Good BM, Unni DR, Harris NL, et al. (2021). The Gene Ontology resource: enriching a GOld mine. Nucleic Acids Research. https://doi.org/10.1093/nar/gkaa1113

23. de Magalhaes JP (2004). HAGR: the Human Ageing Genomic Resources. Nucleic Acids Research. https://doi.org/10.1093/nar/gki017

24. Benjamini Y, Hochberg Y (1995). Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. Journal of the Royal Statistical Society Series B: Statistical Methodology. https://doi.org/10.1111/j.2517-6161.1995.tb02031.x

25. Bergmann O, Bhardwaj RD, Bernard S, Zdunek S, Barnabe-Heider F, Walsh S, et al. (2009). Evidence for Cardiomyocyte Renewal in Humans. Science. https://doi.org/10.1126/science.1164680

26. Saul D, Kosinsky RL, Atkinson EJ, Doolittle ML, Zhang X, LeBrasseur NK, et al. (2022). A new gene set identifies senescent cells and predicts senescence-associated pathways across tissues. Nature Communications. https://doi.org/10.1038/s41467-022-32552-1

27. Oh HS, Rutledge J, Nachun D, Pálovics R, Abiose O, Moran-Losada P, et al. (2023). Organ aging signatures in the plasma proteome track health and disease. Nature. https://doi.org/10.1038/s41586-023-06802-1

28. Ahadi S, Zhou W, Schüssler-Fiorenza Rose SM, Sailani MR, Contrepois K, Avina M, et al. (2020). Personal aging markers and ageotypes revealed by deep longitudinal profiling. Nature Medicine. https://doi.org/10.1038/s41591-019-0719-5

29. Schaum N, Lehallier B, Hahn O, Pálovics R, Hosseinzadeh S, Lee SE, et al. (2020). Ageing hallmarks exhibit organ-specific temporal signatures. Nature. https://doi.org/10.1038/s41586-020-2499-y

30. López-Otín C, Blasco MA, Partridge L, Serrano M, Kroemer G (2023). Hallmarks of aging: An expanding universe. Cell. https://doi.org/10.1016/j.cell.2022.11.001

31. Williams GC (1957). Pleiotropy, natural selection, and the evolution of senescence. Evolution. https://doi.org/10.2307/2406060

32. Kirkwood TBL (1977). Evolution of ageing. Nature. https://doi.org/10.1038/270301a0

33. Blagosklonny MV (2006). Aging and immortality: quasi-programmed senescence and its pharmacologic inhibition. Cell Cycle. https://doi.org/10.4161/cc.5.18.3288

34. Peters MJ, Joehanes R, Pilling LC, Schurmann C, Conneely KN, Powell J, et al. (2015). The transcriptional landscape of age in human peripheral blood. Nature Communications. https://doi.org/10.1038/ncomms957

35. Petkovich DA, Podolskiy DI, Lobanov AV, Lee SG, Miller RA, Gladyshev VN (2017). Using DNA methylation profiling to evaluate biological age and longevity interventions. Cell Metabolism. https://doi.org/10.1016/j.cmet.2017.03.016

36. Hwang CD, Rahmayanti S, Endo Y, Caragher SP, Weber L, Mroueh J, et al. (2026). Transcriptional programs diverge in aging mouse and human skeletal muscle. Aging. https://doi.org/10.18632/aging.206382

37. Tyshkovskiy A, Ma S, Shindyapina AV, Tikhonov S, Lee SG, Bozaykut P, et al. (2023). Distinct longevity mechanisms across and within species and their association with aging. Cell. https://doi.org/10.1016/j.cell.2

38. Ximerakis M, Lipnick SL, Innes BT, Simmons SK, Adiconis X, Dionne D, et al. (2019). Single-cell transcriptomic profiling of the aging mouse brain. Nature Neuroscience. https://doi.org/10.1038/s41593-019-0491-3

39. Zhang MJ, Pisco AO, Darmanis S, Zou J (2021). Mouse aging cell atlas analysis reveals global and cell type-specific aging signatures. eLife. https://doi.org/10.7554/eLife.62293

40. Jeffries AM, Yu T, Ziegenfuss JS, Tolles AK, Baer CE, Bautista Sotelo C, et al. (2025). Single-cell transcriptomic and genomic changes in the ageing human brain. Nature. https://doi.org/10.1038/s41586-025-09435-8

41. De Man R, McDonough JE, Adams TS, Nikola F, Rangel R, Anderson S, et al. (2026). Single-cell atlas of human lung aging identifies cell type dyssynchrony and increased transcriptional entropy. Nature Communications. https://doi.org/10.1038/s41467-026-68810-9

42. Dobner S, Kleissl L, Tóth F, Paxton RL, Yordanova I, Brazdilova K, et al. (2026). Uncovering the transcriptional hallmarks of endothelial cell aging via integrated single-cell analysis. Nature Aging. https://doi.org/10.1038/s43587-026-01247-3

43. Harrison DE, Strong R, Sharp ZD, Nelson JF, Astle CM, Flurkey K, et al. (2009). Rapamycin fed late in life extends lifespan in genetically heterogeneous mice. Nature. https://doi.org/10.1038/nature08221

44. Baker DJ, Wijshake T, Tchkonia T, LeBrasseur NK, Childs BG, van de Sluis B, et al. (2011). Clearance of p16Ink4a-positive senescent cells delays ageing-associated disorders. Nature. https://doi.org/10.1038/nat

45. Buckley MT, Sun ED, George BM, Liu L, Schaum N, Xu L, et al. (2023). Cell-type-specific aging clocks to quantify aging and rejuvenation in neurogenic regions of the brain. Nature Aging. https://doi.org/10.1038/s43587-022-00335-4

46. Sloan NX, Mares J, Daly AC, Grier S, Haq I, Jackson CA, et al. (2026). Uncovering the signatures of aging and senescence in the human dorsolateral prefrontal cortex. Cell Genomics. https://doi.org/10.1016/j.xgen.20

47. Gabitto MI, Travaglini KJ, Rachleff VM, Kaplan ES, Long B, Ariza J, et al. (2024). Integrated multimodal cell atlas of Alzheimer's disease. Nature Neuroscience. https://doi.org/10.1038/s41593-024-01774-5

48. Oh HS-H, Le Guen Y, Rappoport N, Urey DY, Farinas A, Rutledge J, et al. (2025). Plasma proteomics links brain and immune system aging with healthspan and longevity. Nature Medicine. https://doi.org/10.1038/s41591-025-03798-1

49. Kivimäki M, Frank P, Pentti J, Jokela M, Nyberg ST, Blake A, et al. (2025). Proteomic organ-specific ageing signatures and 20-year risk of age-related diseases: the Whitehall II observational cohort study. The Lancet Digital Health. https://doi.org/10.1016/j.landig.2025.01.006

50. Wen J, Tian YE, Skampardoni I, Yang Z, Cui Y, Anagnostakis F, et al. (2024). The genetic architecture of biological age in nine human organ systems. Nature Aging. https://doi.org/10.1038/s43587-024-00662-8