
FDR-significant mouse and human cell-fraction changes
Mouse | Marrow | TMS | late pro-B cell
Mouse | Kidney | TMS | epithelial cell of proximal tubule
Mouse | Liver | TMS | B cell
Mouse | Liver | TMS | myeloid leukocyte
Mouse | Spleen | TMS | plasma cell
Mouse | Liver | TMS | NK cell
Mouse | Lung | TMS | T cell
Human | kidney | Kidney_SN_v15 | non-classical monocyte
Human | lung | Lung_Integrated | pulmonary alveolar type 2 cell
Human | brain | MSSM_DLPFC | oligodendrocyte
Human | brain | MSSM_DLPFC | L6b glutamatergic neuron of the primary motor cortex
Human | brain | MSSM_DLPFC | GABAergic neuron
Human | brain | MSSM_DLPFC | sst GABAergic cortical interneuron
Human | brain | MSSM_DLPFC | oligodendrocyte precursor cell
Human | brain | MSSM_DLPFC | smooth muscle cell
Human | lung | Lung_Integrated | multiciliated epithelial cell
Mouse | Liver | TMS | hepatic stellate cell
Mouse | Marrow | TMS | precursor B cell
Mouse | Bladder | TMS | bladder cell
Human | lung | Lung_Integrated | club cell
Mouse | Liver | TMS | endothelial cell of hepatic sinusoid
Mouse | Kidney | TMS | kidney mesangial cell
Mouse | Kidney | TMS | kidney capillary endothelial cell
Mouse | Lung | TMS | pulmonary interstitial fibroblast
Mouse | Liver | TMS | hepatocyte
−0.3
−0.2
−0.1
0.0
0.1
Total-cell-adjusted CLR age slope, with 95% CI