[DOCUMENT: ONCOLOGY_TRIAL_2025 | SOURCE PAGE: 14]
SOURCE PAGE: 14

PHASE-III CLINICAL TRIAL EFFICACY & DOSAGE TABLE (80MG COHORT)

Clinical Endpoint	80mg Cohort (n=240)	Standard Care (n=238)	p-value
Median Progression-Free Survival (months)	18.4 months	10.2 months	<0.0001
Objective Response Rate (%)	64.2%	38.7%	<0.0001
Grade >=3 Adverse Events (%)	14.1%	29.4%	0.0012

PATIENT ENROLLMENT CRITERIA AND MULTI-CENTER STUDY METHODOLOGY BACKGROUND. Adult patients diagnosed with locally advanced or metastatic non-small cell carcinoma were randomized across forty-two international academic medical centers following histological confirmation, baseline imaging assessment, and institutional review board ethics approval to evaluate long-term tolerability and safety monitoring protocols.

[DOCUMENT: ONCOLOGY_TRIAL_2025 | SOURCE PAGE: 1]
SOURCE PAGE: 1

PHASE-III ONCOLOGY TRIAL 2025 - EXECUTIVE OVERVIEW
The study evaluates a targeted treatment program for adults with locally advanced or metastatic disease. The program is designed around clinically meaningful outcomes, patient safety, consistent site operations, and transparent reporting for investigators and participants.

CLINICAL TRIAL TECHNICAL AND OPERATIONAL ARCHITECTURE

Trial operations connect electronic data capture, randomization, imaging review, laboratory workflows, pharmacovigilance, and site monitoring. A controlled data pipeline supports query resolution, protocol deviations, audit trails, and scheduled interim review.

CLINICAL TRIAL RISK GOVERNANCE

The governance plan monitors informed consent, participant privacy, adverse events, data quality, protocol adherence, and investigator oversight. Safety committees review emerging signals and document decisions, escalation thresholds, and corrective actions.