

## 22. Linear least squares problems

## Last time

- Orthogonality
- Gram–Schmidt algorithm
- QR factorization

# Goals for today

- Formulating data fitting problems
- Optimization
- Linear least squares problems
- Solution using linear algebra

## Parametric function

- Given data points  $(t_i, y_i)$
- Suppose have presumed (statistical) **model** for data

# Parametric function

- Given data points  $(t_i, y_i)$
- Suppose have presumed (statistical) **model** for data
- I.e. suspect related as  $y = f_{\mathbf{p}}(x)$ , with noise:

$$y_i = f_{\mathbf{p}}(t_i) + \epsilon_i$$

where  $\epsilon_i$  represents the noise on observation  $i$

# Parametric function

- Given data points  $(t_i, y_i)$
- Suppose have presumed (statistical) **model** for data
- I.e. suspect related as  $y = f_{\mathbf{p}}(x)$ , with noise:

$$y_i = f_{\mathbf{p}}(t_i) + \epsilon_i$$

where  $\epsilon_i$  represents the noise on observation  $i$

- $f_{\mathbf{p}}$  is **parametric function** with parameter vector  $\mathbf{p}$

# Parametric function

- Given data points  $(t_i, y_i)$
- Suppose have presumed (statistical) **model** for data
- I.e. suspect related as  $y = f_{\mathbf{p}}(x)$ , with noise:

$$y_i = f_{\mathbf{p}}(t_i) + \epsilon_i$$

where  $\epsilon_i$  represents the noise on observation  $i$

- $f_{\mathbf{p}}$  is **parametric function** with parameter vector  $\mathbf{p}$
- Most common example: straight line  $f_{\alpha,\beta}(x) = \alpha + \beta x$

# Best fit

- Want to find **best fit** to the data
- How can we formalize this?



# Best fit

- Want to find **best fit** to the data
- How can we formalize this?
- Find values of  $\alpha, \beta$  that **minimize** distance of data from function
- **Parameter estimation** in statistics

## Loss function

- Distance: **loss function** or **cost function** (optimization)
- E.g.

# Loss function

- Distance: **loss function** or **cost function** (optimization)
- E.g.

$$\mathcal{L}(\alpha, \beta) := \sum_i \ell(y_i, f_{\alpha, \beta}(x_i))$$

where  $\ell$  is a measure of distance

# Loss function

- Distance: **loss function** or **cost function** (optimization)
- E.g.

$$\mathcal{L}(\alpha, \beta) := \sum_i \ell(y_i, f_{\alpha, \beta}(x_i))$$

where  $\ell$  is a measure of distance

- Most common choice

$$\ell(x, y) = \|x - y\|_2^2$$

- **Least squares**

# Least squares

- Want to minimize

$$\mathcal{L}(\mathbf{p}) = \sum_i [y_i - f_{\mathbf{p}}(t_i)]^2$$

# Least squares

- Want to minimize

$$\mathcal{L}(\mathbf{p}) = \sum_i [y_i - f_{\mathbf{p}}(t_i)]^2$$

- Of form  $\sum r_i^2$ , where  $r_i := y_i - f_{\mathbf{p}}(t_i)$  is  $i$ th **residual**
- Sum of squares, so minimum is when all  $r_i = 0$

# Least squares

- Want to minimize

$$\mathcal{L}(\mathbf{p}) = \sum_i [y_i - f_{\mathbf{p}}(t_i)]^2$$

- Of form  $\sum r_i^2$ , where  $r_i := y_i - f_{\mathbf{p}}(t_i)$  is  $i$ th **residual**
- Sum of squares, so minimum is when all  $r_i = 0$
- But data do not lie exactly on any  $y = f_{\alpha,\beta}(x)$
- Choose curve that **minimizes** square sum of residuals

# Least squares

- Want to minimize

$$\mathcal{L}(\mathbf{p}) = \sum_i [y_i - f_{\mathbf{p}}(t_i)]^2$$

- Of form  $\sum r_i^2$ , where  $r_i := y_i - f_{\mathbf{p}}(t_i)$  is  $i$ th **residual**
- Sum of squares, so minimum is when all  $r_i = 0$
- But data do not lie exactly on any  $y = f_{\alpha,\beta}(x)$
- Choose curve that **minimizes** square sum of residuals
- **least squares** minimization
- Statistics: **linear regression**



# Optimization

- Example of an **optimization** problem
- Optimization problems are usually **very hard**; huge field

# Optimization

- Example of an **optimization** problem
- Optimization problems are usually **very hard**; huge field
- Special methods for problems with certain **structure**

# Optimization

- Example of an **optimization** problem
- Optimization problems are usually **very hard**; huge field
- Special methods for problems with certain **structure**
  
- There are many numerical methods for optimization
- Sometimes analytical solutions are possible – e.g. linear least squares

## Matrix formulation of linear least squares

- Look for a matrix formulation:  $\mathcal{L} = \sum_i r_i^2 = r^T r$

## Matrix formulation of linear least squares

- Look for a matrix formulation:  $\mathcal{L} = \sum_i r_i^2 = \mathbf{r}^T \mathbf{r}$

$$\mathbf{r} = \begin{bmatrix} r_1 \\ r_2 \\ \vdots \\ r_m \end{bmatrix} = \begin{bmatrix} 1 & t_1 \\ 1 & t_2 \\ \vdots & \vdots \\ 1 & t_m \end{bmatrix} \begin{bmatrix} \alpha \\ \beta \end{bmatrix} - \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_m \end{bmatrix} =: \mathbf{A}\mathbf{x} - \mathbf{b}$$

where  $\mathbf{x} = \begin{bmatrix} \alpha \\ \beta \end{bmatrix}$  are the unknowns

## Matrix formulation II

- Ideally would like to solve  $Ax = b$

## Matrix formulation II

- Ideally would like to solve  $Ax = b$
- But have **overdetermined system**:
- Too many equations ( $n$ ) for number of unknowns (2)

## Matrix formulation II

- Ideally would like to solve  $Ax = b$
- But have **overdetermined system**:
- Too many equations ( $n$ ) for number of unknowns (2)
- Best we can do is minimize  $r^T r$



## General linear least squares

- General case: want to “solve”  $A\mathbf{x} = \mathbf{b}$   
for **tall and narrow** matrix  $A$

## General linear least squares

- General case: want to “solve”  $A\mathbf{x} = \mathbf{b}$   
for **tall and narrow** matrix  $A$
- i.e.  $(m \times n)$  matrix with  $m > n$
- Again overdetermined system

## General linear least squares

- General case: want to “solve”  $A\mathbf{x} = \mathbf{b}$   
for **tall and narrow** matrix  $A$
- i.e.  $(m \times n)$  matrix with  $m > n$
- Again overdetermined system
  
- Best we can do is minimize

$$\|A\mathbf{x} - \mathbf{b}\|^2$$

# General linear least squares

- General case: want to “solve”  $A\mathbf{x} = \mathbf{b}$   
for **tall and narrow** matrix  $A$
- i.e.  $(m \times n)$  matrix with  $m > n$
- Again overdetermined system
  
- Best we can do is minimize

$$\|A\mathbf{x} - \mathbf{b}\|^2$$

- E.g. fit a polynomial of degree  $< n$  to  $(n + 1)$  points

## Geometrical interpretation

- Suppose  $\mathbf{x}$  ranges over all of  $\mathbb{R}^n$
- Then  $A\mathbf{x}$  ranges over column space / range of  $A$

## Geometrical interpretation

- Suppose  $\mathbf{x}$  ranges over all of  $\mathbb{R}^n$
- Then  $A\mathbf{x}$  ranges over column space / range of  $A$
- **Column space** of  $A$ : vec. space spanned by columns of  $A$

## Geometrical interpretation

- Suppose  $\mathbf{x}$  ranges over all of  $\mathbb{R}^n$
- Then  $A\mathbf{x}$  ranges over column space / range of  $A$
- **Column space** of  $A$ : vec. space spanned by columns of  $A$
- Column space is a hyperplane

## Geometrical interpretation

- Suppose  $\mathbf{x}$  ranges over all of  $\mathbb{R}^n$
- Then  $A\mathbf{x}$  ranges over column space / range of  $A$
- **Column space** of  $A$ : vec. space spanned by columns of  $A$
- Column space is a hyperplane
- Looking for point  $\mathbf{x}$  whose image  $A\mathbf{x}$  is **closest** to  $b$



## Geometrical interpretation

- Suppose  $\mathbf{x}$  ranges over all of  $\mathbb{R}^n$
- Then  $A\mathbf{x}$  ranges over column space / range of  $A$
- **Column space** of  $A$ : vec. space spanned by columns of  $A$
  
- Column space is a hyperplane
- Looking for point  $\mathbf{x}$  whose image  $A\mathbf{x}$  is **closest** to  $b$
- Intuitively: when  $r := A\mathbf{x} - \mathbf{b}$  **perpendicular** to column space

## Solving linear least squares

- Want to find  $\mathbf{x}$  minimizing  $\|\mathbf{Ax} - \mathbf{b}\|$

## Solving linear least squares

- Want to find  $\mathbf{x}$  minimizing  $\|\mathbf{Ax} - \mathbf{b}\|$
- Recall:  $\|\mathbf{x}\|^2 = \mathbf{x}^\top \mathbf{x}$

## Solving linear least squares

- Want to find  $\mathbf{x}$  minimizing  $\|\mathbf{A}\mathbf{x} - \mathbf{b}\|$
- Recall:  $\|\mathbf{x}\|^2 = \mathbf{x}^\top \mathbf{x}$
- Take any vector  $\mathbf{y}$  and look at  $\|\mathbf{A}(\mathbf{x} + \mathbf{y}) - \mathbf{b}\|$ :

## Solving linear least squares

- Want to find  $\mathbf{x}$  minimizing  $\|\mathbf{Ax} - \mathbf{b}\|$
- Recall:  $\|\mathbf{x}\|^2 = \mathbf{x}^\top \mathbf{x}$
- Take any vector  $\mathbf{y}$  and look at  $\|\mathbf{A}(\mathbf{x} + \mathbf{y}) - \mathbf{b}\|$ :

$$\|(\mathbf{Ax} - \mathbf{b}) + \mathbf{Ay}\|^2 = [(\mathbf{Ax} - \mathbf{b}) + \mathbf{Ay}]^\top [(\mathbf{Ax} - \mathbf{b}) + \mathbf{Ay}]$$

## Solving linear least squares

- Want to find  $\mathbf{x}$  minimizing  $\|\mathbf{Ax} - \mathbf{b}\|$
- Recall:  $\|\mathbf{x}\|^2 = \mathbf{x}^\top \mathbf{x}$
- Take any vector  $\mathbf{y}$  and look at  $\|\mathbf{A}(\mathbf{x} + \mathbf{y}) - \mathbf{b}\|$ :

$$\begin{aligned} \|\mathbf{A}(\mathbf{x} + \mathbf{y}) - \mathbf{b}\|^2 &= [\mathbf{A}(\mathbf{x} + \mathbf{y}) - \mathbf{b}]^\top [\mathbf{A}(\mathbf{x} + \mathbf{y}) - \mathbf{b}] \\ &= (\mathbf{Ax} - \mathbf{b})^\top (\mathbf{Ax} - \mathbf{b}) + 2\mathbf{y}^\top \mathbf{A}^\top (\mathbf{Ax} - \mathbf{b}) + (\mathbf{Ay})^\top (\mathbf{Ay}) \end{aligned}$$

## Solving linear least squares

- Want to find  $\mathbf{x}$  minimizing  $\|\mathbf{Ax} - \mathbf{b}\|$
- Recall:  $\|\mathbf{x}\|^2 = \mathbf{x}^\top \mathbf{x}$
- Take any vector  $\mathbf{y}$  and look at  $\|\mathbf{A}(\mathbf{x} + \mathbf{y}) - \mathbf{b}\|$ :

$$\|(\mathbf{Ax} - \mathbf{b}) + \mathbf{Ay}\|^2 = [(\mathbf{Ax} - \mathbf{b}) + \mathbf{Ay}]^\top [(\mathbf{Ax} - \mathbf{b}) + \mathbf{Ay}]$$

$$= (\mathbf{Ax} - \mathbf{b})^\top (\mathbf{Ax} - \mathbf{b}) + 2\mathbf{y}^\top \mathbf{A}^\top (\mathbf{Ax} - \mathbf{b}) + (\mathbf{Ay})^\top (\mathbf{Ay})$$

- Here used  $\mathbf{y}^\top \mathbf{z} = \mathbf{z}^\top \mathbf{y}$ , so  $\mathbf{y}^\top \mathbf{z} + \mathbf{z}^\top \mathbf{y} = 2\mathbf{y}^\top \mathbf{z}$

$$= \|\mathbf{Ax} - \mathbf{b}\|^2 + \|\mathbf{Ay}\|^2 + 2\mathbf{y}^\top \mathbf{A}^\top (\mathbf{Ax} - \mathbf{b})$$

## Solving linear least squares II

■ Arrived at  $\|A\mathbf{x} - \mathbf{b}\|^2 + \|A\mathbf{y}\|^2 + 2\mathbf{y}^\top A^\top (A\mathbf{x} - \mathbf{b})$



## Solving linear least squares II

- Arrived at  $\|A\mathbf{x} - \mathbf{b}\|^2 + \|A\mathbf{y}\|^2 + 2\mathbf{y}^\top A^\top (A\mathbf{x} - \mathbf{b})$
- Take  $\mathbf{x}$  such that  $A^\top (A\mathbf{x} - \mathbf{b}) = 0$

## Solving linear least squares II

- Arrived at  $\|A\mathbf{x} - \mathbf{b}\|^2 + \|A\mathbf{y}\|^2 + 2\mathbf{y}^\top A^\top (A\mathbf{x} - \mathbf{b})$
- Take  $\mathbf{x}$  such that  $A^\top (A\mathbf{x} - \mathbf{b}) = 0$
- Then for any  $\mathbf{y}$  we have

$$\|A(\mathbf{x} + \mathbf{y}) - \mathbf{b}\|^2 \geq \|A\mathbf{x} - \mathbf{b}\|^2$$

## Solving linear least squares II

- Arrived at  $\|A\mathbf{x} - \mathbf{b}\|^2 + \|A\mathbf{y}\|^2 + 2\mathbf{y}^\top A^\top (A\mathbf{x} - \mathbf{b})$
- Take  $\mathbf{x}$  such that  $A^\top (A\mathbf{x} - \mathbf{b}) = 0$
- Then for any  $\mathbf{y}$  we have

$$\|A(\mathbf{x} + \mathbf{y}) - \mathbf{b}\|^2 \geq \|A\mathbf{x} - \mathbf{b}\|^2$$

- So  $\mathbf{x}$  is minimizer

## Solving linear least squares II

- Arrived at  $\|A\mathbf{x} - \mathbf{b}\|^2 + \|A\mathbf{y}\|^2 + 2\mathbf{y}^\top A^\top (A\mathbf{x} - \mathbf{b})$
- Take  $\mathbf{x}$  such that  $A^\top (A\mathbf{x} - \mathbf{b}) = 0$
- Then for any  $\mathbf{y}$  we have

$$\|A(\mathbf{x} + \mathbf{y}) - \mathbf{b}\|^2 \geq \|A\mathbf{x} - \mathbf{b}\|^2$$

- So  $\mathbf{x}$  is minimizer
- Unique solution of least squares problem given by

$$A^\top A\mathbf{x} = A^\top \mathbf{b}$$

## Solving linear least squares III

- Reduced linear least squares problem to *linear* equation:

$$A^T A x = A^T \mathbf{b}$$

– called **normal equations**

- Square matrix  $A^T A$

## Solving linear least squares III

- Reduced linear least squares problem to *linear* equation:

$$A^T A x = A^T \mathbf{b}$$

– called **normal equations**

- Square matrix  $A^T A$
- Solvable if  $A$  is **full rank**, i.e. columns span the space

# Pseudoinverse

- Mathematically get  $\mathbf{x} = (\mathbf{A}^\top \mathbf{A})^{-1} \mathbf{A}^\top \mathbf{b}$

# Pseudoinverse

- Mathematically get  $\mathbf{x} = (\mathbf{A}^\top \mathbf{A})^{-1} \mathbf{A}^\top \mathbf{b}$
- Notation:  $\mathbf{x} = \mathbf{A}^+ \mathbf{b}$
- Where  $\mathbf{A}^+ := (\mathbf{A}^\top \mathbf{A})^{-1} \mathbf{A}^\top$  is **pseudo-inverse**



# Pseudoinverse

- Mathematically get  $\mathbf{x} = (\mathbf{A}^\top \mathbf{A})^{-1} \mathbf{A}^\top \mathbf{b}$
- Notation:  $\mathbf{x} = \mathbf{A}^+ \mathbf{b}$
- Where  $\mathbf{A}^+ := (\mathbf{A}^\top \mathbf{A})^{-1} \mathbf{A}^\top$  is **pseudo-inverse**
- $\mathbf{A}^\top \mathbf{A}$  is invertible iff  $\mathbf{A}$  has full rank

## Solving normal equations

- Possible to solve normal equations using elimination, but to do so need to construct  $A^T A$

## Solving normal equations

- Possible to solve normal equations using elimination, but to do so need to construct  $A^T A$
- Slightly more efficient version of elimination for **symmetric** matrices: **Cholesky factorization**

## Solving normal equations

- Possible to solve normal equations using elimination, but to do so need to construct  $A^T A$
- Slightly more efficient version of elimination for **symmetric** matrices: **Cholesky factorization**
- Alternative: factorize  $A = QR$

## Solving normal equations

- Possible to solve normal equations using elimination, but to do so need to construct  $A^T A$
- Slightly more efficient version of elimination for **symmetric** matrices: **Cholesky factorization**
- Alternative: factorize  $A = QR$
- Need generalization of QR when  $A$  tall and narrow,  $m \times n$

## Solving normal equations

- Possible to solve normal equations using elimination, but to do so need to construct  $A^T A$
- Slightly more efficient version of elimination for **symmetric** matrices: **Cholesky factorization**
- Alternative: factorize  $A = QR$
- Need generalization of QR when  $A$  tall and narrow,  $m \times n$
- **Thin**  $QR$  factorization:  $A = QR$  where
  - $Q$  is  $(m \times n)$  with orthonormal columns
  - $R$  is  $(n \times n)$  upper triangular

## Solving normal equations

- Possible to solve normal equations using elimination, but to do so need to construct  $A^T A$
- Slightly more efficient version of elimination for **symmetric** matrices: **Cholesky factorization**
- Alternative: factorize  $A = QR$
- Need generalization of QR when  $A$  tall and narrow,  $m \times n$
- **Thin**  $QR$  factorization:  $A = QR$  where
  - $Q$  is  $(m \times n)$  with orthonormal columns
  - $R$  is  $(n \times n)$  upper triangular
- If  $A$  is of full rank then  $R$  is non-singular

## Solving normal equations II

- Want to solve  $A^T A = A^T \mathbf{b}$
- Substitute  $A = QR$



## Solving normal equations II

- Want to solve  $A^T A = A^T \mathbf{b}$
- Substitute  $A = QR$
- So  $R^T Q^T QR \mathbf{x} = R^T Q^T \mathbf{b}$

## Solving normal equations II

- Want to solve  $A^T A = A^T \mathbf{b}$
- Substitute  $A = QR$
- So  $R^T Q^T QR \mathbf{x} = R^T Q^T \mathbf{b}$
- $R^T R \mathbf{x} = R^T Q^T \mathbf{b}$

## Solving normal equations II

- Want to solve  $A^T A = A^T \mathbf{b}$
- Substitute  $A = QR$
- So  $R^T Q^T QR \mathbf{x} = R^T Q^T \mathbf{b}$
- $R^T R \mathbf{x} = R^T Q^T \mathbf{b}$
- $R \mathbf{x} = Q^T \mathbf{b}$

## Solving normal equations II

- Want to solve  $A^T A = A^T \mathbf{b}$
- Substitute  $A = QR$
- So  $R^T Q^T QR \mathbf{x} = R^T Q^T \mathbf{b}$
- $R^T R \mathbf{x} = R^T Q^T \mathbf{b}$
- $R \mathbf{x} = Q^T \mathbf{b}$
- So  $\mathbf{x} = R^{-1} Q^T \mathbf{b}$

## Solving normal equations II

- Want to solve  $A^T A = A^T \mathbf{b}$
- Substitute  $A = QR$
- So  $R^T Q^T QR \mathbf{x} = R^T Q^T \mathbf{b}$
- $R^T R \mathbf{x} = R^T Q^T \mathbf{b}$
- $R \mathbf{x} = Q^T \mathbf{b}$
- So  $\mathbf{x} = R^{-1} Q^T \mathbf{b}$
- Solve  $R \mathbf{x} = Q^T \mathbf{b}$  by backsubstitution

# Backslash

- Backslash in Julia is overloaded to give this least-squares solution when  $A$  is an  $(m \times n)$  matrix
- It is not necessarily clear that this is a good idea!

# Backslash

- Backslash in Julia is overloaded to give this least-squares solution when  $A$  is an  $(m \times n)$  matrix
- It is not necessarily clear that this is a good idea!
- E.g. simple linear fit: use `\` with above matrix

# Summary

- Linear least squares for overdetermined systems
- Solvable using linear algebra
- Solution given by normal equations – linear system
- Solve using QR decomposition