

19. Linear algebra

Last time

- Mid-term review
- Designing auxiliary problems
- Convergence

Goals for today

- Conceptual review of linear algebra
- Emphasising **geometry**
- Vectors
- Linear transformations
- Matrices
- Mostly think about \mathbb{R}^2

What is linear algebra?

- **Linear:** Something related to “lines”

What is linear algebra?

- **Linear:** Something related to “lines”
 - **flatness**

What is linear algebra?

- **Linear:** Something related to “lines”
 - flatness
- **Algebra:** Manipulating, solving, understanding “structure”

What is linear algebra?

- **Linear:** Something related to “lines”
 - **flatness**
- **Algebra:** Manipulating, solving, understanding “structure”
- Study of **vectors** and **linear transformations** between them

Why linear algebra?

- Have repeatedly seen systems of linear equations crop up:
 - First-order Taylor expansion
 - Newton's method (linear equations to solve nonlinear equations)
 - Coefficients of interpolating polynomial (Vandermonde matrix)

Why linear algebra?

- Have repeatedly seen systems of linear equations crop up:
 - First-order Taylor expansion
 - Newton's method (linear equations to solve nonlinear equations)
 - Coefficients of interpolating polynomial (Vandermonde matrix)
- Discretising many PDEs gives linear systems
- Numerical linear algebra: fundamental importance in science and engineering

Why linear algebra II

- **Algebraic:** simplest relation between ≥ 2 variables:

$$ax + by = c$$

- linear equation relating variables x and y

Why linear algebra II

- **Algebraic:** simplest relation between ≥ 2 variables:

$$ax + by = c$$

– linear equation relating variables x and y

- (Relative) simplicity allows us to obtain much information

Why linear algebra II

- **Algebraic:** simplest relation between ≥ 2 variables:

$$ax + by = c$$

– linear equation relating variables x and y

- (Relative) simplicity allows us to obtain much information
- **Geometry:** “flat” objects – lines, planes, hyperplanes
- Flatness makes much easier to understand than general curved surfaces
 - “rigidity”

Vectors

- **Vectors** are the cornerstone of linear algebra
- What is a vector?

Vectors

- **Vectors** are the cornerstone of linear algebra
- What is a vector?
- Directed / oriented piece of line
- Displacement through space
- Column of numbers

Operations on vectors

- Must be able to operate on vectors in 2 ways:

- 1 **Scalar multiplication:** αv

Operations on vectors

- Must be able to operate on vectors in 2 ways:

- 1 Scalar multiplication:** αv

- Geometry: Stretch vector by factor α in same direction

Operations on vectors

- Must be able to operate on vectors in 2 ways:

- 1 Scalar multiplication:** αv

- Geometry: Stretch vector by factor α in same direction
- “Scaling”

Operations on vectors

- Must be able to operate on vectors in 2 ways:

1 **Scalar multiplication:** αv

- Geometry: Stretch vector by factor α in same direction
- “Scaling”

2 **Addition:** $v_1 + v_2$

Operations on vectors

- Must be able to operate on vectors in 2 ways:

1 **Scalar multiplication:** αv

- Geometry: Stretch vector by factor α in same direction
- “Scaling”

2 **Addition:** $v_1 + v_2$

- Geometry: Move along one vector and then the other

Linear combinations

- In general, + and scalar multiplication *only* operations
- So we can only form (finite) **linear combinations**

$$\alpha_1 v_1 + \cdots + \alpha_n v_n$$

Linear combinations

- In general, + and scalar multiplication *only* operations
- So we can only form (finite) **linear combinations**

$$\alpha_1 v_1 + \cdots + \alpha_n v_n$$

- Define **span** $\langle v_1, \dots, v_n \rangle$ as set of *all* linear combinations

Geometry of spans

- What are spans geometrically?

Geometry of spans

- What are spans geometrically?
- Points in space that are **reachable** using given vectors

Geometry of spans

- What are spans geometrically?
- Points in space that are **reachable** using given vectors
- Simplest case: 1 vector

Geometry of spans

- What are spans geometrically?
- Points in space that are **reachable** using given vectors
- Simplest case: 1 vector
 $\langle v \rangle$ is line through origin in direction of v

Geometry of spans

- What are spans geometrically?
- Points in space that are **reachable** using given vectors

- Simplest case: 1 vector
 $\langle v \rangle$ is line through origin in direction of v

- 2 vectors:
 $\langle v_1, v_2 \rangle$: unique plane containing v_1 and v_2

Geometry of spans

- What are spans geometrically?
- Points in space that are **reachable** using given vectors

- Simplest case: 1 vector
 $\langle v \rangle$ is line through origin in direction of v

- 2 vectors:
 $\langle v_1, v_2 \rangle$: unique plane containing v_1 and v_2
except if v_2 is in $\langle v_1 \rangle$
 i.e. if $v_2 = \beta v_1$ – **linearly dependent**

Geometry of spans III

- 3 vectors:

$\langle v_1, v_2, v_3 \rangle$: 3D hyperplane containing the vectors

Geometry of spans III

- 3 vectors:

$\langle v_1, v_2, v_3 \rangle$: 3D hyperplane containing the vectors

- all of space if $v_i \in \mathbb{R}^3$

Geometry of spans III

- 3 vectors:

$\langle v_1, v_2, v_3 \rangle$: 3D hyperplane containing the vectors

- all of space if $v_i \in \mathbb{R}^3$
- **unless** e.g. $v_3 \in \langle v_1, v_2 \rangle$

Geometry of spans III

- 3 vectors:

$\langle v_1, v_2, v_3 \rangle$: 3D hyperplane containing the vectors

- all of space if $v_i \in \mathbb{R}^3$

- **unless** e.g. $v_3 \in \langle v_1, v_2 \rangle$

- All spans contain 0

- A span is a **vector subspace**

Lines

- Vector subspaces must pass through the origin
- More general: **affine subspace**

Lines

- Vector subspaces must pass through the origin
- More general: **affine subspace**
- Simplest is straight line: need one point on line and direction

Lines

- Vector subspaces must pass through the origin
- More general: **affine subspace**
- Simplest is straight line: need one point on line and direction
- $\ell(t) = p + tv$ – parametric equation

Lines

- Vector subspaces must pass through the origin
- More general: **affine subspace**
- Simplest is straight line: need one point on line and direction
- $\ell(t) = p + tv$ – parametric equation
- E.g. Line joining points a and b :

$$(1 - \alpha)a + \alpha b$$

- $\alpha = 0$ gives a and $\alpha = 1$ gives b

Vector equations

- Suppose $\mathbf{u} := \begin{bmatrix} u_1 \\ u_2 \end{bmatrix}$ and $\mathbf{v} := \begin{bmatrix} v_1 \\ v_2 \end{bmatrix}$ span \mathbb{R}^2
- Suppose $\mathbf{b} := \begin{bmatrix} b_1 \\ b_2 \end{bmatrix} \in \mathbb{R}^2$

Vector equations

- Suppose $\mathbf{u} := \begin{bmatrix} u_1 \\ u_2 \end{bmatrix}$ and $\mathbf{v} := \begin{bmatrix} v_1 \\ v_2 \end{bmatrix}$ span \mathbb{R}^2
- Suppose $\mathbf{b} := \begin{bmatrix} b_1 \\ b_2 \end{bmatrix} \in \mathbb{R}^2$
- Then there must exist scalars x and y such that

$$x \mathbf{u} + y \mathbf{v} = \mathbf{b}$$

Vector equations

- Suppose $\mathbf{u} := \begin{bmatrix} u_1 \\ u_2 \end{bmatrix}$ and $\mathbf{v} := \begin{bmatrix} v_1 \\ v_2 \end{bmatrix}$ span \mathbb{R}^2
- Suppose $\mathbf{b} := \begin{bmatrix} b_1 \\ b_2 \end{bmatrix} \in \mathbb{R}^2$
- Then there must exist scalars x and y such that

$$x \mathbf{u} + y \mathbf{v} = \mathbf{b}$$

- x, y : “how much \mathbf{u} and \mathbf{v} need to reach \mathbf{b} ”

Vector equations II

- Suppose $x\mathbf{u} + y\mathbf{v} = \mathbf{b}$
- Rewrite in terms of components:

$$x \begin{bmatrix} u_1 \\ u_2 \end{bmatrix} + y \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \end{bmatrix}$$

Vector equations II

- Suppose $x\mathbf{u} + y\mathbf{v} = \mathbf{b}$
- Rewrite in terms of components:

$$x \begin{bmatrix} u_1 \\ u_2 \end{bmatrix} + y \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \end{bmatrix}$$

- Take components:

$$u_1x + v_1y = b_1$$

$$u_2x + v_2y = b_2$$

Vector equations II

- Suppose $x\mathbf{u} + y\mathbf{v} = \mathbf{b}$
- Rewrite in terms of components:

$$x \begin{bmatrix} u_1 \\ u_2 \end{bmatrix} + y \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \end{bmatrix}$$

- Take components:

$$u_1x + v_1y = b_1$$

$$u_2x + v_2y = b_2$$

- So linear system can be interpreted as finding correct linear combination of vectors

Linear maps

- Consider a function / map $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$
- In general: complicated and hard to understand

Linear maps

- Consider a function / map $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$
- In general: complicated and hard to understand
- To understand and calculate, study maps with **structure**
- i.e. restrict to some *subset* of all maps

Linear maps

- Consider a function / map $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$
- In general: complicated and hard to understand
- To understand and calculate, study maps with **structure**
- i.e. restrict to some *subset* of all maps
- One very important subclass are **linear** maps

Linear maps II

- A map $f : X \rightarrow Y$ is **linear** if

$$f(\alpha x + \beta y) = \alpha f(x) + \beta f(y) \quad \forall x, y \in X$$

Linear maps II

- A map $f : X \rightarrow Y$ is **linear** if

$$f(\alpha x + \beta y) = \alpha f(x) + \beta f(y) \quad \forall x, y \in X$$

- A linear map **preserves** operations $+$ and \star
- Doesn't matter whether we do them before or after applying the map

Linear maps II

- Geometrical interpretation of this linearity condition?

Linear maps II

- Geometrical interpretation of this linearity condition?
- What is action of linear map on a line?

Linear maps II

- Geometrical interpretation of this linearity condition?
- What is action of linear map on a line?

$$f((1 - \alpha)a + \alpha b) = (1 - \alpha)a' + \alpha b'$$

where $a' := f(a)$ and $b' := f(b)$

Linear maps II

- Geometrical interpretation of this linearity condition?
- What is action of linear map on a line?

$$f((1 - \alpha)a + \alpha b) = (1 - \alpha)a' + \alpha b'$$

where $a' := f(a)$ and $b' := f(b)$

- So linear map maps lines to lines
and moves along them at the same speed

Linear maps III

- Linearity *greatly* restricts what f can do

Linear maps III

- Linearity *greatly* restricts what f can do
- Suppose x and y span \mathbb{R}^2
- Suppose f is linear map from \mathbb{R}^2 to \mathbb{R}^2
- Then we know *exactly* what f does for *every* point on the plane!

Geometry of linear maps

- What does a linear map (or **linear transformation**) do geometrically?

Geometry of linear maps

- What does a linear map (or **linear transformation**) do geometrically?
- Squishes unit square into parallelogram
- Or rotates it
- Or reflects it

How do we specify a linear map?

- What does a general linear map look like?
- Specifying action on any two (non-collinear) vectors

$$f(a) =: a' \text{ and } f(b) =: b'$$

specifies everything

How do we specify a linear map?

- What does a general linear map look like?
- Specifying action on any two (non-collinear) vectors

$$f(a) =: a' \text{ and } f(b) =: b'$$

specifies everything

- In \mathbb{R}^2 encode via action on **standard basis vectors**

$$e_1 := \begin{bmatrix} 1 \\ 0 \end{bmatrix} \text{ and } e_2 := \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

Matrices

- e_1 mapped to $a_1 := f(e_1)$
- e_2 mapped to $a_2 := f(e_2)$
- Need 4 parameters; encode as **matrix**

$$\begin{bmatrix} | & | \\ a_1 & a_2 \\ | & | \end{bmatrix}$$

- First **column** is image of $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$
- Second **column** is image of $\begin{bmatrix} 1 \\ 0 \end{bmatrix}$
- $Ae_1 = a_{1,1}e_1 + a_{2,1}e_2$

Matrix–vector multiplication

- Let matrix A represent linear map f
- Define **matrix–vector product** by

$$A x := f(x)$$

Matrix–vector multiplication

- Let matrix A represent linear map f
- Define **matrix–vector product** by

$$A x := f(x)$$

- Let A have columns a_1 and a_2
- Let x be $\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$
- Then $f(x) = f(x_1 e_1 + x_2 e_2) = x_1 f(e_1) + x_2 f(e_2) = x_1 a_1 + x_2 a_2$
- So $A x$ is **linear combination of columns of A**

Matrix–vector multiplication

- For matrix A and vector x define $A \cdot x$ as

$$A \cdot x := \begin{bmatrix} | & | \\ a_1 & a_2 \\ | & | \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} := xa_1 + ya_2$$

- Matrix–vector multiplication is **linear combination** of columns of A

Matrix–vector multiplication

- For matrix A and vector x define $A \cdot x$ as

$$A \cdot x := \left[\begin{array}{c|c} | & | \\ a_1 & a_2 \\ | & | \end{array} \right] \begin{bmatrix} x \\ y \end{bmatrix} := xa_1 + ya_2$$

- Matrix–vector multiplication is **linear combination** of columns of A
- Linear system becomes

$$Ax = b$$

Matrix–matrix multiplication

- Let A, B represent linear maps f, g resp.

Matrix–matrix multiplication

- Let A, B represent linear maps f, g resp.
- Define **matrix–matrix product** $A B$ by action on vectors:

$$(A B)(x) := A(B x) = f(g(x)) = (f \circ g)(x)$$

Matrix–matrix multiplication

- Let A, B represent linear maps f, g resp.
- Define **matrix–matrix product** AB by action on vectors:

$$(AB)(x) := A(Bx) = f(g(x)) = (f \circ g)(x)$$

- Calculate as

$$AB = \left[\begin{array}{c|c} (A|b_1) & (A|b_2) \end{array} \right]$$

Matrix–matrix multiplication

- Let A, B represent linear maps f, g resp.
- Define **matrix–matrix product** AB by action on vectors:

$$(AB)(x) := A(Bx) = f(g(x)) = (f \circ g)(x)$$

- Calculate as

$$AB = \left[\begin{array}{c|c} (A|b_1) & (A|b_2) \end{array} \right]$$

- Since $(AB)(e_1) = A(Be_1) = Ab_1$

Summary

- Vectors are basic objects
- Linearly independent spans form a basis
- Define matrices as recording action of a linear map