

University of the People

Discrete Mathematics Assignment

Instructor: Bhumika Modi

Abraham Chileshe

May 21, 2025

1 Partial Ordering Relation


A partial ordering relation is a binary relation that is reflexive, antisymmetric, and transitive. I'll explain this using the “divides” relation ($|$) on the set $\{1, 2, 3, 6, 12\}$.

In the divides relation, we say that $a|b$ if a divides b evenly (without remainder). Formally, $a|b$ if there exists an integer k such that $b = a \cdot k$.

Let's verify that the divides relation is a partial ordering:

1. **Reflexive:** For any element a , $a|a$ (since $a = a \cdot 1$). ✓
2. **Antisymmetric:** If $a|b$ and $b|a$, then $a = b$. ✓
(If a divides b and b divides a , they must be equal)
3. **Transitive:** If $a|b$ and $b|c$, then $a|c$. ✓
(If a divides b and b divides c , then a divides c as well)

Here's the Hasse diagram for this relation:



Explanation of the Hasse Diagram:

- In a Hasse diagram, elements are represented as nodes, and the relation between elements is shown by lines connecting them.
- Elements are arranged vertically according to their “rank” - if $a|b$, then b appears above a .
- We omit drawing edges implied by transitivity and reflexivity to keep the diagram clean.
- In this diagram, 1 is at the bottom since it divides all other numbers.
- 12 is at the top since it's divisible by all other numbers in the set.
- If we can follow a path upward from element a to element b , then $a|b$.
- Note that not all pairs of elements are comparable. For example, 2 and 3 are not related by the divides relation (neither divides the other), which is why there's no path between them.

2 Transitive Closure of Relations

2.1 Definition and Uses

The transitive closure of a relation R on a set A is the smallest transitive relation that contains R . It's denoted as R^+ or R^* and includes all pairs that are connected directly or indirectly through the relation.

Formally, (a, c) is in the transitive closure of R if either:

- (a, c) is in R , or
- There exists some b such that (a, b) and (b, c) are both in R , or
- There's a longer chain connecting a to c through the relation.

Uses of Transitive Closure:

1. **Graph Theory:** Finding all reachable nodes from a given node in a graph
2. **Database Systems:** Computing all dependencies between attributes
3. **Network Analysis:** Determining all nodes that can be reached from a starting point
4. **Programming Language Semantics:** Defining the inheritance relation between classes

Example: Consider a relation $R = \{(a, b), (b, c), (d, e)\}$ on the set $\{a, b, c, d, e\}$. R itself is not transitive because (a, b) and (b, c) are in R , but (a, c) is not.

The transitive closure $R^+ = \{(a, b), (b, c), (d, e), (a, c)\}$ R^+ adds (a, c) to make the relation transitive.

2.2 Equivalence Relation

Consider the relation R defined on the set of integers as $R = \{(x, y) : x, y \in \mathbb{Z}, (x - y) \text{ is a multiple of } 11\}$.

To show R is an equivalence relation, I need to prove it's reflexive, symmetric, and transitive.

1. **Reflexive:** For any $x \in \mathbb{Z}$, $(x - x) = 0$, which is a multiple of 11 ($0 = 0 \times 11$). So $(x, x) \in R$ for all x . ✓
2. **Symmetric:** If $(x, y) \in R$, then $(x - y)$ is a multiple of 11. This means $(x - y) = 11k$ for some integer k . Then $(y - x) = -(x - y) = -11k = 11(-k)$, which is also a multiple of 11. So $(y, x) \in R$. ✓

3. **Transitive:** If $(x, y) \in R$ and $(y, z) \in R$, then:

$$(x - y) = 11k_1 \text{ for some integer } k_1$$

$$(y - z) = 11k_2 \text{ for some integer } k_2$$

$$\text{Adding: } (x - y) + (y - z) = (x - z) = 11k_1 + 11k_2 = 11(k_1 + k_2)$$

So $(x - z)$ is a multiple of 11, which means $(x, z) \in R$. ✓

Since R is reflexive, symmetric, and transitive, it is an equivalence relation.

2.3 Equivalence Class of 0

The equivalence class of 0, denoted $[0]$, is the set of all integers y such that $(0, y) \in R$. This means $(0 - y) = -y$ must be a multiple of 11.

So $y = 11k$ for some integer k . Therefore, $[0] = \{\dots, -33, -22, -11, 0, 11, 22, 33, \dots\} = \{11k : k \in \mathbb{Z}\}$

2.4 Number of Equivalence Classes

For the relation R , two integers x and y are in the same equivalence class if and only if $(x - y)$ is a multiple of 11.

We can classify any integer by its remainder when divided by 11. There are exactly 11 possible remainders: 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10.

So there are exactly 11 distinct equivalence classes:

$$[0] = \{x \in \mathbb{Z} : x \equiv 0 \pmod{11}\}$$

$$[1] = \{x \in \mathbb{Z} : x \equiv 1 \pmod{11}\}$$

$$[2] = \{x \in \mathbb{Z} : x \equiv 2 \pmod{11}\}$$

$$\vdots$$

$$[10] = \{x \in \mathbb{Z} : x \equiv 10 \pmod{11}\}$$

3 Mathematical Induction Proof

Prove that for all positive integers $n \geq 1$, $5n > 4n$ using mathematical induction.

Base Case ($n = 1$):

$$5(1) = 5$$

$$4(1) = 4$$

$$5 > 4 \quad \checkmark$$

Inductive Hypothesis: Assume that for some $k \geq 1$, $5k > 4k$.

Inductive Step: Need to prove that $5(k + 1) > 4(k + 1)$

$$5(k + 1) = 5k + 5$$

$$4(k + 1) = 4k + 4$$


From the inductive hypothesis, we know $5k > 4k$.
Adding 5 to both sides: $5k + 5 > 4k + 5$
Since $5 > 4$, we know $4k + 5 > 4k + 4$
Therefore, by transitivity: $5k + 5 > 4k + 4$
So $5(k + 1) > 4(k + 1)$ ✓
By the principle of mathematical induction, $5n > 4n$ holds for all positive integers $n \geq 1$.

4 Relation $R = ' \geq '$ on a Set of Five Integers

Let's choose the set $A = \{5, 8, 13, 21, 34\}$ (five integers from the Fibonacci sequence).

The relation $R = ' \geq '$ (greater than or equal to) is defined on A .

4.1 Diagrammatical Representation of R on A



4.2 Adjacency Matrix for R on A

For the relation $R = ' \geq '$ on the set $A = \{5, 8, 13, 21, 34\}$, the adjacency matrix is:

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 1 & 1 \end{pmatrix}$$


In this matrix:

- Row i and column j contain a 1 if element $i \geq$ element j
- Row i and column j contain a 0 if element $i <$ element j

For example:

- The entry at row 8, column 5 is 1 because $8 \geq 5$ (true)
- The entry at row 5, column 8 is 0 because $5 \geq 8$ (false)
- All diagonal entries are 1 because every number is equal to itself ($x \geq x$)

4.3 Hasse Diagram for $R = ' \geq '$ on A



Explanation of the Hasse Diagram:

- In a Hasse diagram for the \geq relation, larger elements appear higher in the diagram.
- The diagram shows the transitive reduction of the relation, omitting edges that can be inferred through transitivity.
- For this relation $R = ' \geq '$ on $A = \{5, 8, 13, 21, 34\}$, the elements form a total ordering.
- The diagram shows a single vertical chain with 34 at the top (greatest) and 5 at the bottom (smallest).
- Each element is connected only to its immediate predecessor/successor, as the other relationships can be inferred through transitivity.
- $34 \geq 21 \geq 13 \geq 8 \geq 5$, and from this we can infer all other relations like $34 \geq 5$.

This Hasse diagram illustrates that ' \geq ' is a total order on this set, meaning every pair of elements is comparable (either $a \geq b$ or $b \geq a$ for any a, b in the set).

References

- [1] Rosen, K. H. (2019). *Discrete Mathematics and Its Applications* (8th ed.). McGraw-Hill Education.
- [2] Grimaldi, R. P. (2017). *Discrete and Combinatorial Mathematics: An Applied Introduction* (5th ed.). Pearson.
- [3] Epp, S. S. (2019). *Discrete Mathematics with Applications* (4th ed.). Cengage Learning.