CROSS-VALIDATION ERROR BARS WHEN ROWS ARE CLUSTERED: WHAT GROUPED CV ESTIMATES AND WHICH ERROR BARS COVER IT

SOPHIA QIN, JENNA YANG, AND ABRAHAM YEUNG

Abstract. In the Columbia speed-dating data (21 events, 4,184 woman–man pairs) every row shares people and an evening with other rows, so cross-validation (CV) error bars that treat rows as independent are too narrow, and random-row folds estimate the wrong quantity. We collect what the team has established about CV error bars when rows are clustered. Holding out whole clusters makes CV estimate the error on new clusters (Lemma 1). On grouped folds, the Nadeau–Bengio variance equals the random-group, or cluster-robust, variance times $(J + K - 1)/(J - 1)$ (Proposition 6). A coverage study that keeps the clusters fixed overstates the coverage of the cluster-robust interval (Proposition 7). A first-order expansion (Proposition 8) shows that the cluster-robust variance is first-order right for the full model's new-cluster error, and that the Nadeau–Bengio variance differs from it by the factor $\mathrm{Var}(a - \beta/(K-1))/\mathrm{Var}(a)$, near 1.14 to 1.22 for the 0–1 error rate and near 1 for the AUC in our designs. Of 27 pre-registered simulation hypotheses, 21 were supported and 6 were not; we report all of them. Cluster-robust intervals with t quantiles, and the wild cluster bootstrap-t, covered the new-cluster error 0.853 to 0.915 of the time at nominal 0.90 in every design with a precisely scored target. The Nadeau–Bengio limit was predicted out of sample at $K = 3$ and 10 to within 0.053. The nested CV of Bates, Hastie and Tibshirani, as published with random-row folds, covered 0.460 to 0.735. For the AUC, pooling held-out scores across fold models biases the CV estimate downward by order $1/G$ when clusters differ in outcome rate (Proposition 9). A second-order expansion predicted that bias to within 5.1% at $K = 5$; at $K = 2$ under strong clustering the registered test missed by 18.6%, mostly sampling error according to a later review. An estimate that uses only pairs from different clusters within a fold removed the bias, and its interval covered 0.848 to 0.938 for $G \geq 40$. With unequal cluster sizes the first-order results carry over with cluster totals and a row-weighted target (Proposition 11), but with the 21 real event sizes the Nadeau–Bengio ratio followed a finite-G heuristic at the actual sizes and partitions rather than the first-order limit; the four registered tests passed, three of them near their band edges. Most ingredients are known, and we say which parts may be new.

1. Introduction

The data come from the Columbia speed-dating experiment (Fisman, Iyengar, Kamenica and Simonson, 2002 to 2004), in its public release. At each of 21 events (waves), every woman met every man for four minutes, and each said yes or no; a match is a mutual yes. The team's main analysis predicts a match from information available before the date, with one row per woman–man pair (a *dyad*): 4,184 dyads, with a match rate of 0.165. Events range from 25 to 484 dyads.[1]

The rows are not independent: a woman at an event with ten men appears in ten dyads, and her pickiness and desirability enter all ten. Every person attends one event, so all dependence through shared people sits inside events, together with whatever is common to an evening. We treat rows from different events as independent. This has two consequences for CV. First, if rows are dealt into folds one at a time (*random-row* folds), a held-out dyad's two people have other dyads in the training set, so CV measures the error for people already seen rather than for new people. Second, the usual standard error of a CV estimate, $s/\sqrt{n}$ with s the standard deviation of the per-row losses, ignores the covariance between rows of the same event and is too small even when the folds are grouped.

Date: 15 October 2026 (draft); updated 3 November 2026.

Math 195 capstone, Stanford University. **Draft for the team's review**, 15 October 2026, updated 3 November 2026 (Part 11); the theory has not yet been checked by the team's mathematical lead. Numbering of results follows the theory note `proposals/06`.

[1]Computed from `data/pairs.csv`; two events have 441 and 484 dyads. `PREREG.md` Part 7 says 25 to 440, corrected in its change log.

The question of this note is therefore: when rows are clustered, what does CV estimate, and which error bar gives an interval that covers the fitted model's error on rows from new clusters? The project's headline numbers use event-held-out 5-fold CV with a cluster-robust standard error. Sections 2 and 3 give the setup and theory, Section 4 the simulation evidence, organized around hypotheses registered in `PREREG.md` before each test ran, Section 5 the real-data error bars, Section 6 the cross-validated AUC, and Section 7 unequal cluster sizes. Sections 8 to 10 give recommendations, what is known and what may be new, and limitations.

2. Setup and the two targets

Rows $z_i = (x_i, y_i)$, $i = 1, \dots, n$, each lie in one cluster $g(i) \in \{1, \dots, G\}$; in the theory the clusters are balanced, $n = Gm$. We use a *two-level model*: cluster effects $U_1, \dots, U_G$ are i.i.d., and given U_g the rows of cluster g are i.i.d. from a distribution P_{U_g}. Rows in different clusters are independent; rows in the same cluster are exchangeable but dependent. A learner maps a training set D to a fitted function $\hat{f}_D$ and is symmetric (it does not depend on the order of the training rows). The loss $\ell(z; f)$ has finite second moments. Folds $F_1, \dots, F_K$ are assigned at random, independently of the data. Under *grouped* folds each cluster lies entirely in one fold, with G/K clusters per fold; under *random-row* folds rows are assigned individually, so clusters straddle folds. With $\hat{f}_{-k}$ trained on all folds but F_k,

$$e_i = \ell(z_i; \hat{f}_{-k(i)}), \qquad \widehat{\mathrm{Err}}_{\mathrm{CV}} = \frac{1}{n}\sum_{i=1}^{n} e_i, \qquad \bar{e}_g = \text{mean of } e_i \text{ over cluster } g.$$

The two targets. For a model trained on D,

$$\mathrm{Err}^{\mathrm{new}}(D) = \mathbb{E}\big[\ell(Z'; \hat{f}_D) \mid D\big], \qquad Z' \text{ a row from a fresh cluster},$$

$$\mathrm{Err}^{\mathrm{same}}(D) = \frac{1}{G}\sum_{g} \mathbb{E}\big[\ell(Z'_g; \hat{f}_D) \mid D, U_g\big], \qquad Z'_g \text{ a fresh row from existing cluster } g.$$

In the dating data a new cluster is a new event of people we have not seen, so the target is $\mathrm{Err}_{\mathrm{new}} := \mathrm{Err}^{\mathrm{new}}(D)$ for the model trained on all the data. The simulations also report coverage of the *average* error $\mathbb{E}[\mathrm{Err}^{\mathrm{new}}(D)]$ over training sets. One more distinction matters below: an interval for $\mathrm{Err}_{\mathrm{new}}$ must be built from $\mathrm{Var}(\widehat{\mathrm{Err}}_{\mathrm{CV}} - \mathrm{Err}_{\mathrm{new}})$, not from $\mathrm{Var}(\widehat{\mathrm{Err}}_{\mathrm{CV}})$, and the two differ when the CV estimate and the target are correlated.

Variance estimates and intervals. The *naive* variance is s^2/n, with s^2 the sample variance of the e_i. The *cluster-robust* variance, in the unbalanced form every script uses, is $\hat{V}_{\mathrm{CR}} = \frac{G}{G-1}\, n^{-2} \sum_g \big(\sum_{i \in g} (e_i - \widehat{\mathrm{Err}}_{\mathrm{CV}})\big)^2$, which for balanced clusters equals $\frac{1}{G(G-1)} \sum_g (\bar{e}_g - \widehat{\mathrm{Err}}_{\mathrm{CV}})^2$; the Wald interval of Bayle et al. [6] with clusters as units omits the factor $G/(G-1)$. *CR2* divides each cluster's residual sum by $\sqrt{1 - n_g/n}$ and uses t_{G-1} quantiles; alternatively $\hat{V}_{\mathrm{CR}}$ is used with t quantiles on $G^* - 1$, where $G^* = (\sum_g n_g)^2 / \sum_g n_g^2$ is the effective number of clusters (13.8 for the real events). The *wild cluster bootstrap-t* puts Rademacher signs on each cluster's residual sum (999 draws), recomputes the cluster-robust t, and takes a symmetric interval from the 0.90 quantile of $|t^*|$. The *Nadeau–Bengio* variance [22, 9] uses the $J = RK$ fold mean losses $e_1, \dots, e_J$ of R partitions into K folds: $\hat{\sigma}^2_{\mathrm{NB}} = (\frac{1}{J} + \frac{1}{K-1}) S^2$, with S^2 their sample variance and t_{J-1} quantiles. *Nested CV* [5] runs, for each outer fold k, an inner CV on D_{-k} giving $\hat{e}^{(k)}_{\mathrm{in}}$, and the outer fold gives $e^{(k)}_{\mathrm{out}}$; with $a_k = (\hat{e}^{(k)}_{\mathrm{in}} - e^{(k)}_{\mathrm{out}})^2$ and b_k an estimate of $\mathrm{Var}(e^{(k)}_{\mathrm{out}} \mid D_{-k})$, the MSE estimate is $\mathrm{mean}_k\, a_k - \mathrm{mean}_k\, b_k$, averaged over repetitions, optionally with the bias adjustment of [5]. The published version uses random-row folds and a within-fold row variance for b_k. The team's *cluster-nested* version groups both levels and uses a cluster-robust b_k, and the *floor* uses $\max(\hat{V}_{\mathrm{CR}}, \widehat{\mathrm{MSE}}_{\mathrm{CN}})$.

3. Theory

3.1. What each CV scheme estimates.

Lemma 1. *Under the two-level model with grouped folds and a symmetric learner, for every row i in fold k, $\mathbb{E}[e_i \mid D_{-k}] = \mathrm{Err}^{\mathrm{new}}(D_{-k})$, so $\mathbb{E}[\widehat{\mathrm{Err}}_{\mathrm{CV}}] = \mathbb{E}[\mathrm{Err}^{\mathrm{new}}(D')]$ for a training set D' of $G(K-1)/K$ clusters. Under random-row folds, a held-out row's cluster typically has other rows in D_{-k}, and the corresponding identity holds with a same-cluster estimand at the same training size.*

Proof. With grouped folds the cluster of row i is disjoint from D_{-k}, so z_i is independent of D_{-k} and has the marginal law of a row from a fresh cluster. Conditioning on D_{-k} gives the first identity; averaging over i and k gives the second. Under random-row folds, condition on D_{-k} and on $U_{g(i)}$: then z_i is a fresh draw from $P_{U_{g(i)}}$, which is the same-cluster estimand. □

Remark. The gap $\mathrm{Err}^{\mathrm{new}} - \mathrm{Err}^{\mathrm{same}}$ is the *cluster leakage*. It is not signed in general, but it is nonnegative whenever the learner can use cluster-level structure in x to absorb the cluster effect in y. No variance estimate can repair a random-row interval for the new-cluster error, because the interval is centred on a different quantity.

3.2. Variance of the grouped CV estimate.

By exchangeability, the symmetry of the learner and the random fold assignment, the joint law of a pair (e_i, e_j), $i \neq j$, depends only on the pair's type. Let $\sigma^2 = \mathrm{Var}(e_i)$; let σ_{cc}, σ_{sf} and σ_{df} be $\mathrm{Cov}(e_i, e_j)$ for i, j in the same cluster, in different clusters of the same fold, and in different folds; and let $\delta_c = \sigma_{cc} - \sigma_{sf}$.

Proposition 1. *Under the two-level model with grouped balanced folds,*

$$\mathrm{Var}(\widehat{\mathrm{Err}}_{\mathrm{CV}}) = \frac{\sigma^2}{n} + \Big(\frac{1}{K} - \frac{1}{n}\Big)\sigma_{sf} + \Big(1 - \frac{1}{K}\Big)\sigma_{df} + \frac{m-1}{n}\,\delta_c.$$

Proof sketch. There are $n(m-1)$ ordered pairs in the same cluster, $n(n/K - m)$ in the same fold but different clusters, and $n(n - n/K)$ in different folds. Sum the covariances, divide by n^2, and substitute $\sigma_{cc} = \sigma_{sf} + \delta_c$. □

The first three terms are the independent-rows formula ($m = 1$); the last is the cluster term. Since $\mathbb{E}[s^2] = \frac{n}{n-1}(\sigma^2 - \mathrm{Var}(\widehat{\mathrm{Err}}_{\mathrm{CV}}))$ for any e_i with common mean and variance, $\mathbb{E}[\hat{V}_{\mathrm{naive}}] \approx \sigma^2/n$, which, ignoring the between-cluster terms, falls short by the Kish design effect $1 + (m-1)\sigma_{cc}/\sigma^2$ [16].

Let $v = \mathrm{Var}(\bar{e}_g)$ and let $\bar{c}$ be the average of $\mathrm{Cov}(\bar{e}_g, \bar{e}_{g'})$ over $g' \neq g$.

Proposition 2. *Under the conditions of Proposition 1, with $V = \mathrm{Var}(\widehat{\mathrm{Err}}_{\mathrm{CV}})$,*

$$V = \frac{v}{G} + \frac{G-1}{G}\,\bar{c}, \qquad \mathbb{E}[\hat{V}_{\mathrm{CR}}] = \frac{v}{G} - \frac{\bar{c}}{G}, \qquad V - \mathbb{E}[\hat{V}_{\mathrm{CR}}] = \bar{c}.$$

Proof sketch. The first identity is Proposition 1 at the cluster level. For the second, apply the sample-variance identity above to the G cluster means, whose mean is $\widehat{\mathrm{Err}}_{\mathrm{CV}}$: $\mathbb{E}[\hat{V}_{\mathrm{CR}}] = (v - V)/(G-1)$. Substitute the first identity. □

Remark. Two corrections from the adversarial review of the theory note apply. First, $\bar{c}$ *can have either sign*: by Proposition 8 below, $\bar{c} = [2\,\mathrm{Cov}(a, \beta) + \mathrm{Var}(\beta)]/G + o(1/G)$, which is first order and of either sign. (In the review's own Monte Carlo with 0–1 loss, as reported in the theory note, $\hat{V}_{\mathrm{CR}}$ overstated $\mathrm{Var}(\widehat{\mathrm{Err}}_{\mathrm{CV}})$ by about 25% at every G.) Second, $\mathrm{Var}(\widehat{\mathrm{Err}}_{\mathrm{CV}})$ is not the interval's target; $\mathrm{Var}(\widehat{\mathrm{Err}}_{\mathrm{CV}} - \mathrm{Err}_{\mathrm{new}})$ is (Proposition 8(i)). Propositions 1 and 2 are exact only for balanced clusters; the unbalanced estimator the scripts use is not covered.

Remark (Proposition 3 of the theory note). With grouped folds at both levels and a symmetric learner, the unclipped cluster-nested MSE estimate has expectation $\mathbb{E}[(\hat{e}_{\mathrm{in}}^{(k)} - \mathrm{Err}^{\mathrm{new}}(D_{-k}))^2]$, the MSE of grouped $(K-1)$-fold CV on $G(K-1)/K$ clusters for the new-cluster error of the model trained on them: given D_{-k}, the held-out clusters' mean errors are i.i.d. with mean $\mathrm{Err}^{\mathrm{new}}(D_{-k})$, so the cluster-robust b_k is unbiased. With the independent-rows b_k the estimate is biased upward by about the within-fold cluster covariance and over-covers. The estimate is unbiased in mean but noisy at $K = 5$; the floor is a heuristic, conservative by design.

3.3. **The Nadeau–Bengio correction on grouped folds.** This part concerns the test-sampling part of the variance, with the losses held fixed (no refitting). Let $\ell_1, \dots, \ell_n$ be fixed losses in $G = Kq$ balanced clusters of m rows, with mean $\bar{y}$, cluster means $\bar{\ell}_g$, $s_c^2 = \frac{1}{G-1}\sum_g(\bar{\ell}_g - \bar{y})^2$ and $s^2 = \frac{1}{n-1}\sum_i(\ell_i - \bar{y})^2$. A grouped partition deals the clusters at random into K folds of q clusters; a row partition deals the rows at random into K folds of n/K rows. Take R partitions and let $e_1, \dots, e_J$, $J = RK$, be the fold mean losses.

Proposition 6. *Taking expectation over the random partitions given the losses (the R partitions need not be independent),*

$$\mathbb{E}[\hat{\sigma}^2_{\mathrm{NB}}] = \frac{J+K-1}{J-1}\cdot\frac{s_c^2}{G} \quad \textit{(grouped)}, \qquad \mathbb{E}[\hat{\sigma}^2_{\mathrm{NB}}] = \frac{J+K-1}{J-1}\cdot\frac{s^2}{n} \quad \textit{(rows)}.$$

Proof. (1) Every partition's folds average to $\bar{y}$, so the mean of all J fold means is $\bar{y}$ exactly, and $\mathbb{E}[S^2] = \frac{1}{J-1}\sum_j \mathbb{E}[(e_j - \bar{y})^2] = \frac{J}{J-1}\operatorname{Var}_{\mathrm{sel}}(e_j)$, the selection variance of one fold mean. (2) A grouped fold mean is the mean of q cluster means drawn without replacement from G, so $\operatorname{Var}_{\mathrm{sel}}(e_j) = (1-q/G)\, s_c^2/q = (K-1)s_c^2/G$; for rows it is $(K-1)s^2/n$. (3) Multiply by $\frac{1}{J} + \frac{1}{K-1} = \frac{J+K-1}{J(K-1)}$. □

With balanced clusters, s_c^2/G is exactly $\hat{V}_{\mathrm{CR}}$ (Section 2) and s^2/n is the naive variance. So on grouped folds the test-sampling part of the Nadeau–Bengio variance is the cluster-robust variance times $1 + K/(J-1)$, and on row folds it is the naive variance times the same factor.

Remark (where this sits). The case $J = K$ is the random group method of survey sampling [31, Chapter 2]: averaged over random divisions of a fixed sample, the random-group variance equals the with-replacement variance, and in multistage samples the groups must be formed from whole clusters. Nadeau and Bengio [22] derived their $\rho_0 = n_2/n$ from the same fixed-loss approximation for random train/test splits; Bouckaert and Frank [9] used it in repeated K-fold CV without a derivation; Markatou et al. [20] found empirically that the correction on a single K-fold run is about twice their variance approximation. Proposition 6 is the random-group estimator rewritten in CV notation, with a small exact extension and a reading for grouped against row folds.

Consequences. (a) Under the two-level model $\mathbb{E}[s_c^2/G] \approx \operatorname{Var}(\bar{y})$, while s^2/n falls short by the design effect: the correction handles clustering *only because the folds are grouped*, which N7 tested. (b) With one CV run ($R = 1$, $J = K$) the factor is $(2K-1)/(K-1)$, or 2.25 at $K = 5$. (c) With refitting, fold errors also vary because the model changes; the conjecture that the excess over $\frac{J+K-1}{J-1}\hat{V}_{\mathrm{CR}}$ tracks $\bar{c}$ was refuted by the review (Proposition 8 gives the excess).

3.4. **Fixed against redrawn clusters in a coverage study.**

Proposition 7. *Hold the clusters fixed, in balanced clusters of m rows; cluster g has expected loss μ_g, set by its fixed composition. Let $\ell_i = \mu_{g(i)} + u_{g(i)} + v_i$ with independent mean-zero u_g (variance σ_u^2) and v_i (variance σ_v^2), and $S_\mu^2 = \frac{1}{G-1}\sum_g(\mu_g - \bar{\mu})^2$. Then*

$$\operatorname{Var}(\bar{y} \mid \textit{design}) = \frac{\sigma_u^2}{G} + \frac{\sigma_v^2}{n}, \qquad \mathbb{E}[\hat{V}_{\mathrm{CR}} \mid \textit{design}] = \operatorname{Var}(\bar{y} \mid \textit{design}) + \frac{S_\mu^2}{G}.$$

If instead the clusters are redrawn, with μ_g i.i.d. of variance σ_μ^2, then $\mathbb{E}[\hat{V}_{\mathrm{CR}}] = \operatorname{Var}(\bar{y}) = (\sigma_\mu^2 + \sigma_u^2)/G + \sigma_v^2/n$ exactly.

Proof. The cluster means are $\bar{\ell}_g = \mu_g + u_g + \bar{v}_g$. With the design fixed, $\mathbb{E}\sum_g(\bar{\ell}_g - \bar{y})^2 = \sum_g(\mu_g - \bar{\mu})^2 + (G-1)(\sigma_u^2 + \sigma_v^2/m)$; divide by $G(G-1)$. With the design random, add $\mathbb{E}[S_\mu^2] = \sigma_\mu^2$. □

A coverage study that reuses the same clusters in every replicate measures the conditional variance, while $\hat{V}_{\mathrm{CR}}$ also counts the spread of the clusters' expected losses, so the cluster-robust interval over-covers there by an amount set by how different the clusters are. The proposition has no refitting and assumes balance, so it explains the A2 over-coverage of Section 4.1 qualitatively, not quantitatively.

3.5. **First-order theory: what grouped CV estimates and which error bar fits it.** Clusters are i.i.d. and balanced. The model is fitted by an M-estimator on a set T of clusters, with first-order expansion $\hat{\theta}_T - \theta^* \approx |T|^{-1} \sum_{g \in T} \psi_g$, the influence functions ψ_g being i.i.d. across clusters with mean 0. The evaluation loss may differ from the training criterion (in the project, 0–1 loss with the threshold at the training match rate, for a model fitted by log-likelihood). Let $A_h(\theta)$ be cluster h's mean evaluation loss, $A(\theta) = \mathbb{E}[A_h(\theta)]$, $a_h = A_h(\theta^*)$, $\mu = A(\theta^*)$, $d = \nabla A(\theta^*)$ and $\beta_g = d^\top \psi_g$. Write $\bar{a}$, $\bar{\beta}$ for averages over all G clusters and $\bar{a}_F$, $\bar{\beta}_F$ for averages over a set F of clusters.

Assumptions for Proposition 8. (1) K divides G, and the partition is a uniform random balanced grouped partition. (2) $\hat{\theta}$ is $\sqrt{G}$-consistent with the stated linearization, and θ^* is interior and unique. (3) Smooth losses: A_h and A are twice differentiable near θ^*, with finite second moments. (4) 0–1 loss: the threshold is part of θ, with influence $\psi_g^t = \bar{y}_g - t^*$, and $\partial A/\partial t$ is included in d; the linear score has a bounded continuous density near the threshold.

Proposition 8 (first order, balanced grouped K-fold CV)**.** *To first order,*

$$\widehat{\mathrm{Err}}_{\mathrm{CV}} = \bar{a} + \bar{\beta} + r_1, \qquad \mathrm{Err}_{\mathrm{new}}(\hat{\theta}_{\mathrm{all}}) = \mu + \bar{\beta} + r_2, \qquad \widehat{\mathrm{Err}}_{\mathrm{CV}} - \mathrm{Err}_{\mathrm{new}} = \bar{a} - \mu + r_1 - r_2.$$

For smooth losses $r_1 - r_2 = O_p(1/G)$, which includes an $O(1/G)$ bias from the smaller training sets. For 0–1 loss $r_1 - r_2$ is only $O_p(G^{-3/4})$: a cluster's loss is a step function of θ, so its change is bounded by a flip probability of order $|\hat{\theta} - \theta^*|$. In both cases $r_1 - r_2 = o_p(G^{-1/2})$, which suffices for (i) to (iii).

- *(i)* $G \operatorname{Var}(\widehat{\mathrm{Err}}_{\mathrm{CV}} - \mathrm{Err}_{\mathrm{new}}) \to \operatorname{Var}(a)$ *(with uniform integrability), and* $G \hat{V}_{\mathrm{CR}} \to_p \operatorname{Var}(a)$.
- *(ii) The fold errors of R grouped partitions are* $\bar{\tilde{a}}_{F_j} + K\bar{\beta}/(K-1)$ *with* $\tilde{a}_h = a_h - \beta_h/(K-1)$, *the same in every partition. So by Proposition 6,* $G\,\mathbb{E}[\hat{\sigma}^2_{\mathrm{NB}}] \to \frac{J+K-1}{J-1} \operatorname{Var}(a - \beta/(K-1))$. *In expectation, the Nadeau–Bengio variance therefore exceeds the relevant variance whenever* $\operatorname{Var}(\beta) > 2(K-1)\operatorname{Cov}(a, \beta)$ *(sufficient, not necessary, at finite J).*
- *(iii) If* $\operatorname{Var}(\beta) > 0$, *then* $\operatorname{Corr}(\widehat{\mathrm{Err}}_{\mathrm{CV}}, \mathrm{Err}_{\mathrm{new}}) \to \dfrac{\operatorname{Cov}(a, \beta) + \operatorname{Var}(\beta)}{\sqrt{\operatorname{Var}(a + \beta)\operatorname{Var}(\beta)}}$.
- *(iv) Suppose three things hold: the evaluation loss is the training criterion, with any penalty $o(1)$; training and evaluation weight rows the same way (balanced clusters, or a correctly specified model); and θ^* is interior and unique. Then $d = 0$ and $\beta \equiv 0$, and the Nadeau–Bengio variance tends to $\frac{J+K-1}{J-1}$ times the cluster-robust variance.* $\mathrm{Err}_{\mathrm{new}}$ *then varies only at second order, and the correlation in (iii) goes to* 0 *at rate* $G^{-1/2}$.

Derivation. (1) Condition on the training set T of fold F; the q test clusters are independent of $\hat{\theta}_T$, so $\frac{1}{q}\sum_{h \in F}[A_h(\hat{\theta}_T) - A_h(\theta^*)] - [A(\hat{\theta}_T) - A(\theta^*)]$ has mean 0 and variance at most $\mathbb{E}[(A_h(\hat{\theta}_T) - A_h(\theta^*))^2]/q$, which is $O(G^{-2})$ for smooth losses and $O(G^{-3/2})$ for 0–1 loss. (2) $A(\hat{\theta}_T) - A(\theta^*) = d^\top(\hat{\theta}_T - \theta^*) + O_p(1/G)$, by Taylor's theorem and the linearization remainder. (3) So fold j's error is $\bar{a}_{F_j} + \bar{\beta}_{T_j}$ plus remainders. Each cluster lies in $K-1$ training sets of equal size, so averaging over folds gives $\bar{a} + \bar{\beta}$. (4) The full model's error is $A(\hat{\theta}_{\mathrm{all}}) = \mu + \bar{\beta} + O_p(1/G)$. (5) Within a partition, $\bar{\beta}_{T_j} - \bar{\beta} = -\frac{1}{K-1}(\bar{\beta}_{F_j} - \bar{\beta})$, so fold j's error is $\bar{a}_{F_j} - \bar{\beta}_{F_j}/(K-1) + K\bar{\beta}/(K-1)$; the last term is common to all folds of all partitions and does not affect S^2, and Proposition 6 applied to the losses $\tilde{a}_h$ gives (ii). Parts (i) and (iii) follow from the displayed decomposition: $\operatorname{Var}(\widehat{\mathrm{Err}}_{\mathrm{CV}}) \approx \operatorname{Var}(a + \beta)/G$, $\operatorname{Var}(\mathrm{Err}_{\mathrm{new}}) \approx \operatorname{Var}(\beta)/G$ and $\operatorname{Cov}(\widehat{\mathrm{Err}}_{\mathrm{CV}}, \mathrm{Err}_{\mathrm{new}}) \approx [\operatorname{Cov}(a, \beta) + \operatorname{Var}(\beta)]/G$. □

Remark (relation to the literature). Bayle et al. [6] target the average of the K fold models' conditional test errors. They assume i.i.d. points and a loss-stability condition but no smoothness, and with clusters as the units their central limit theorem already gives (i) for *that* target, including 0–1 loss. Proposition 8 adds the step to the full-data model: $\frac{1}{K}\sum_j A(\hat{\theta}_{T_j}) - A(\hat{\theta}_{\mathrm{all}}) = O_p(1/G)$ because $\frac{1}{K}\sum_j \bar{\beta}_{T_j} = \bar{\beta}$, a step that needs smooth A and carries an $O(1/G)$ training-size bias. Wager [30] overlaps (i) and (iv).

Remark. With unbalanced events and a misspecified marginal model (for example an omitted random intercept), $d \neq 0$ even for log-loss when the target weights every cluster equally; for the row-weighted

target that the scripts score, $d = 0$ (Proposition 11(e)). Proposition 8 also explains how the cluster-robust interval can be right although $\bar{c} \neq 0$: to first order it estimates $\mathrm{Var}(\widehat{\mathrm{Err}}_{\mathrm{CV}} - \mathrm{Err}_{\mathrm{new}})$, not $\mathrm{Var}(\widehat{\mathrm{Err}}_{\mathrm{CV}})$.

4. Simulation evidence

Every hypothesis below was committed to `PREREG.md` before its test ran, and none was edited afterwards. Table 1 lists all of them; Part 9 (N16 to N21) is discussed in Section 6 and Part 11 (N22 to N25) in Section 7. Unless stated otherwise, intervals are nominal 90%, the procedure is a logistic regression with 0–1 loss at the training match rate, and CV uses 5 folds. The realistic designs (Parts 2 to 5 and 7) use the *A2 truth*: the real dyads and events, a logistic model for the match on four pre-date predictors, and woman, man and event random effects with SDs and coefficients from a crossed random-effects fit to the real matches; the target is the full-data model's error on new events.

4.1. **The published nested CV fails on clustered data (A2, N2).** In A2 (500 replicates, seed 20261906; Table 2) the 21 real events were kept in every replicate and the target was scored on 300 new events. The conventional interval (random-row CV, naive SE) covered 0.790, just under A2's threshold of 0.80, and event CV with the cluster-robust SE covered 0.992; A2 was supported. The published nested CV of [5] on random-row folds covered 0.736 unadjusted and 0.678 bias-adjusted. The cluster-robust 0.992 is over-coverage, as Proposition 7 explains: with the events fixed, the variation from drawing new events is absent from the truth but present in $\hat{V}_{\mathrm{CR}}$ (tested by N4).

The design grid of Part 3 (Table 2, right) varied $G \in \{10, 21, 40\}$, the person-effect SD and the event-effect SD. N2 was supported: in the 12 designs with person SD at least 0.5, the published, bias-adjusted interval covered 0.460 to 0.735. N1 was not supported: the cluster-nested, bias-adjusted interval covered 0.735 to 0.920, with 6 of 15 designs in the band 0.85 to 0.95. The theory note traces this to the variance of the MSE estimate (Proposition 3 concerns only its mean): in 4% to 15% of data sets it was zero or negative, giving a zero-width interval, and among the rest the interval covered 84% to 98%.

4.2. **The repair, and Nadeau–Bengio on grouped and row folds (N3 to N7).** Part 4 redrew the events in every replicate (Table 3). N3 was supported: the floor covered 0.890 to 0.965 in all 15 designs with person or event effects. N4 was supported: with the A2 events redrawn, the cluster-robust interval covered 0.874 rather than 0.992, confirming the fixed-design explanation. N5 was supported: the cluster-nested, bias-adjusted interval covered 0.836 rather than 0.926; in the grid its MSE estimate was not positive in 0.065 to 0.150 of data sets per design. Exploratory: the Nadeau–Bengio t on grouped folds covered 0.900 to 0.975 with mean half-width 0.0377, against 0.0466 for the floor; grouped nested CV with a row-level variance over-covered (0.980 to 0.995), as the remark after Proposition 3 predicts; internal–external CV under-covered at $G = 40$ (0.720 to 0.790).

Part 5 tested the Nadeau–Bengio interval on a fresh grid (Table 4). N6 was *not* supported: on event folds it covered 0.880 to 0.975, with 8 of 12 designs in the band; the four designs with 10 events over-covered (0.960 to 0.975). Its half-width was 0.73 to 0.90 times the floor's, so the width condition held; only the coverage band failed. N7 was supported, as Proposition 6 predicted before the run: on random-row folds the same correction covered 0.515 to 0.760. The median ratio of $\hat{\sigma}^2_{\mathrm{NB}}$ to $\frac{J+K-1}{J-1}\hat{V}_{\mathrm{CR}}$ was 1.29 to 1.44, so the Nadeau–Bengio variance exceeded the scaled cluster-robust variance; Proposition 8(ii) gives the form of such a factor, but not its value for unbalanced events.

4.3. **First-order predictions (N9 to N11, N14, N15).** Part 6 tested Proposition 8 where it applies cleanly: $G \in \{20, 40, 80, 160, 320\}$ balanced clusters of 40 rows, $x = u_g + v_i$ with $u_g, v_i \sim N(0, I_3)$, $y \sim \mathrm{Bernoulli}(\mathrm{logistic}(-1.5 + x^\top(1, 0.5, 0) + a_g))$, $a_g \sim N(0, 1)$, a logistic regression on x, one grouped 5-fold partition for $\widehat{\mathrm{Err}}_{\mathrm{CV}}$ and $\hat{V}_{\mathrm{CR}}$, and $R = 10$ more ($J = 50$) for $\hat{\sigma}^2_{\mathrm{NB}}$. The losses are 0–1 at the training positive rate and log-loss, the training criterion, for which (iv) applies. $\mathrm{Err}_{\mathrm{new}}$ was scored on 2,000 new clusters (Part 6), then on 20,000 with its Monte Carlo variance subtracted (Part 6b); see Table 5.

TABLE 1. Every pre-registered hypothesis of the error-bar track, with its outcome. Verdicts are read from the `*_hypotheses.csv` files and were re-derived from the coverage tables by the table script (Part 11's from `sim_cv_unequal_summary.csv`). Part 11's ratios are Nadeau–Bengio ratios $\hat{\sigma}^2_{\mathrm{NB}}/(\frac{J+K-1}{J-1}\hat{V}_{\mathrm{CR}})$.

Hyp.	Part	Registered claim (abbreviated)	Result	Verdict
A2	2	Random-row < 0.80; event CV, cluster-robust SE ≥ 0.85 (events fixed)	0.790; 0.992	supported
N1	3	Cluster-nested, bias-adj. covers 0.85–0.95 in all 15 designs	0.735–0.920; 6/15 in band	**not supported**
N2	3	Published BHT (random rows) < 0.80 if person SD ≥ 0.5	0.460–0.735; 12/12 below	supported
N3	4	Floor covers ≥ 0.85 in all 15 designs (events redrawn)	0.890–0.965	supported
N4	4	A2, events redrawn: cluster-robust covers 0.85–0.95	0.874	supported
N5	4	A2, events redrawn: cluster-nested, bias-adj. below 0.926	0.836	supported
N6	5	NB t, event folds: 0.85–0.95 in all 12; width $\leq 0.9\times$ floor's	0.880–0.975; 8/12; 0.73–0.90	**not supported**
N7	5	NB t, random-row folds: below 0.80 in all 12 designs	0.515–0.760	supported
N8	6	$\mathrm{Var}(\mathrm{CV}-\mathrm{Err}_{\mathrm{new}})/\overline{V}_{\mathrm{CR}} \in [0.85, 1.15]$, $G \geq 80$, both losses	1.060–1.155	**not supported**
N9	6	NB ratio in $[0.9, 1.1]$ (log, $G = 160, 320$); > 1.1 at all G (0–1)	1.005, 0.994; 1.145–1.223	supported
N10	6	Log-loss $\mathrm{Corr}(\mathrm{CV}, \mathrm{Err}_{\mathrm{new}}) < 0.2$, $G \geq 80$	-0.106 to -0.043	supported
N11	6	Cluster-robust t_{G-1} covers 0.85–0.95, every G, both losses	0.875–0.902	supported
N8b	6b	As N8 with a precise target; corrected ratio in $[0.9, 1.1]$	0.911–1.165	**not supported**
N12	7	Wild cluster bootstrap-t covers 0.85–0.95 in all 8 designs	0.853–0.910	supported
N13	7	Cluster-robust (normal) < 0.85 in some $G = 40$ design	0.867–0.883	**not supported**
N14	8	0–1 loss: NB ratio within 0.07 of prediction, $K = 3, 10$	max. |diff| 0.053	supported
N15	8	Log-loss: NB ratio within 0.05 of 1, $K = 3, 10$	max. |diff| 0.032	supported
N16	9	Pooled CV AUC biased down (90% interval < 0), outcome clustering, $G \leq 80$	-0.004 to -0.040	supported
N17	9	Mixing bias at $K = 5$ within 15% of $C_{\mathrm{mix}} + c_2/4$, four designs	off by 1.3%–5.1%	supported
N18	9	Mixing bias at $K = 2, 10$ within 15% of $C_{\mathrm{mix}} + c_2/(K-1)$	3 of 4; one off by 18.6%	**not supported**
N19	9	BCWF-pooled |bias| ≤ 0.25 SE for $G \geq 40$; pooled > 0.25 SE when $s_a = 2$	max. 0.11; 0.48–0.87	supported
N20	9	BCWF-pooled mean coverage 0.87–0.93; pooled < 0.85 when $s_a = 2$	0.899; 0.841	supported
N21	9	NB ratio for fold AUCs within 0.07 of prediction (about 1)	0.978–0.999	supported
N22	11	21 real sizes, 0–1: own-row ratio within 0.04 of P2 ($K = 10$); within 0.03 of P2 and below the P1/P2 midpoint ($K = 5$)	1.007; 1.230	supported
N23	11	84 resampled sizes, 0–1: own-row ratio within 0.03 of P1, $K = 5, 10$	1.266; 1.078	supported
N24	11	A2, 21 redrawn events: own-row ratio within 0.08 of its plug-in; own $-$ residual-total ≥ 0.10	-0.009, $+0.068$; 0.192, 0.203	supported
N25	11	21 real sizes, log: residual-total within 0.03 of 1, own-row within 0.03 of P2, $K = 5, 10$	1.007, 0.994; 1.092, 0.993	supported

N9 was supported: under log-loss the ratio of $\hat{\sigma}^2_{\mathrm{NB}}$ to $\frac{J+K-1}{J-1}\hat{V}_{\mathrm{CR}}$ was 1.005 and 0.994 at $G = 160$ and 320, and under 0–1 loss it stayed near 1.2 (1.145 to 1.223; 1.183 to 1.209 in Part 6b). N10 was supported: under log-loss $\mathrm{Corr}(\widehat{\mathrm{Err}}_{\mathrm{CV}}, \mathrm{Err}_{\mathrm{new}})$ was -0.106 to -0.043 for $G \geq 80$ (-0.106 to -0.021 over both runs); under 0–1 loss it was positive (0.080 to 0.208), as (iii) allows. N11 was supported: the cluster-robust t_{G-1} interval covered $\mathrm{Err}_{\mathrm{new}}$ 0.875 to 0.902 at every G for both losses (0.875 to 0.915 in Part 6b); with normal quantiles, 0.863 to 0.915.

Predicted against observed Nadeau–Bengio ratio. Under 0–1 loss the variances and covariance of $\widehat{\mathrm{Err}}_{\mathrm{CV}}$ and $\mathrm{Err}_{\mathrm{new}}$ determine $\mathrm{Var}(a)$, $\mathrm{Var}(\beta)$ and $\mathrm{Cov}(a, \beta)$ to first order (see the derivation of Proposition 8), and hence a predicted ratio $\mathrm{Var}(a - \beta/4)/\mathrm{Var}(a)$ (Table 6). In Part 6b the observed ratio fell inside

TABLE 2. Coverage of nominal 90% intervals. Left: A2, the 21 real events in every replicate (500 replicates, seed 20261906), for $\mathrm{Err}_{\mathrm{new}}$ and for the average error, with mean half-width. Right: the Part 3 design grid, coverage of $\mathrm{Err}_{\mathrm{new}}$, range over the five designs with person or event effects at each G (200 replicates per design, seeds 20262806 + design index, $R = 10$); $G = 21$ keeps the real events, $G = 10$ and 40 redraw them. N2 uses the 12 designs with person SD at least 0.5.

	A2 (events fixed)			Design grid		
Procedure	$\mathrm{Err}_{\mathrm{new}}$	Average	Half-width	$G = 10$	$G = 21$	$G = 40$
Random-row CV, naive SE (conventional)	0.790	0.630	0.0125	0.550–0.650	0.725–0.890	0.490–0.580
Random-row nested CV (BHT)	0.736	0.622	0.0139	0.515–0.680	0.690–0.765	0.495–0.610
Random-row nested CV (BHT), bias-adj.	0.678	0.566	0.0124	0.465–0.600	0.645–0.735	0.460–0.540
Event CV, naive SE	0.746	0.604	0.0125	0.535–0.635	0.680–0.840	0.480–0.585
Event CV, cluster-robust SE	0.992	0.938	0.0288	0.855–0.945	0.965–1.000	0.840–0.915
Event CV, cluster-nested	0.938	0.908	0.0401	0.850–0.870	0.870–0.935	0.765–0.860
Event CV, cluster-nested, bias-adj.	0.926	0.886	0.0359	0.830–0.860	0.865–0.920	0.735–0.820
Event CV, floor	1.000	0.982	0.0431	–	–	–

TABLE 3. Part 4 repair grid, events redrawn in every data set: coverage range and mean half-width over the 15 designs with person or event effects (200 data sets each; seeds 20263006 + design), and A2 with its 21 events redrawn (500 data sets; seeds 20263106 + chunk). Event folds unless named.

	Grid, 15 designs		A2, events redrawn	
Procedure	Coverage	Mean half-width	Coverage	Half-width
Random-row CV, naive SE	0.465–0.660	0.0133	0.540	0.0127
Event CV, cluster-robust SE	0.815–0.920	0.0298	0.874	0.0280
Cluster-nested, bias-adjusted	0.720–0.855	0.0381	0.836	0.0366
Floor (N3)	0.890–0.965	0.0466	0.956	0.0440
Nadeau–Bengio t, grouped folds	0.900–0.975	0.0377	0.956	0.0358
Cluster bootstrap of grouped CV	0.820–0.965	0.0335	0.894	0.0322
Internal–external CV, random effects	0.720–0.915	0.0319	0.812	0.0298
Grouped nested CV, row-level variance	0.980–0.995	0.0624	0.992	0.0594

the prediction's 90% bootstrap interval at 5 of 5 values of G, with $\mathrm{Corr}(a, \beta)$ from -0.568 to -0.493. In Part 6 it fell inside at only 2 of 5; there the estimate of $\mathrm{Var}(\beta)$ absorbs the target's Monte Carlo variance ($G\,\mathrm{Var}(\beta) = 0.0134$ at $G = 320$, against 0.0083 in Part 6b).

Out-of-sample test at $K = 3$ and 10. Before Part 8 ran, the components were estimated from Part 6b's $K = 5$ runs pooled over $G = 80, 160, 320$: $G\,\mathrm{Var}(a) = 0.02266$, $G\,\mathrm{Var}(\beta) = 0.00946$, $G\,\mathrm{Cov}(a, \beta) = -0.00787$. Proposition 8(ii), $\mathrm{Var}(a - \beta/(K-1))/\mathrm{Var}(a)$, then predicts 1.452 at $K = 3$, 1.082 at $K = 10$, 2.112 at $K = 2$ and 1.200 at $K = 5$, and 1 at every K for log-loss (Table 7). N14 was supported: under 0–1 loss the observed ratio was within 0.053 of the prediction at $K = 3$ and 10 at every G, and the $K = 3$ discrepancy shrank as G grew. N15 was supported: under log-loss the ratio was within 0.032 of 1. In the exploratory $K = 2$ runs the 0–1 ratio was 2.184 to 2.251, 3.4% to 6.6% above the prediction, where half the data is held out and the expansion is least accurate; under log-loss it reached 1.056. Monte Carlo SEs of the observed ratios were 0.009 to 0.032.

4.4. **The failed variance-ratio tests (N8, N8b) and why.** Proposition 8(i) predicts $\mathrm{Var}(\widehat{\mathrm{Err}}_{\mathrm{CV}} - \mathrm{Err}_{\mathrm{new}})/\widehat{V}_{\mathrm{CR}} \to 1$. N8 registered the band $[0.85, 1.15]$ at $G = 80, 160, 320$ for both losses; the ratios were 1.060 to 1.155, 1 of 6 outside, so N8 was not supported. The diagnosis was that $\mathrm{Err}_{\mathrm{new}}$, scored on 2,000 new clusters, carries Monte Carlo noise that adds to $\mathrm{Var}(\widehat{\mathrm{Err}}_{\mathrm{CV}} - \mathrm{Err}_{\mathrm{new}})$, most at large G. Under log-loss the SD of $\mathrm{Err}_{\mathrm{new}}$ at $G = 80, 160, 320$ was 0.0046, 0.0044, 0.0043 in Part 6, where it stops

TABLE 4. Part 5 confirmatory grid, events redrawn (200 data sets per design; seeds 20263406 + design index; $J = RK = 50$). Coverage of nominal 90% intervals for $\mathrm{Err}_{\mathrm{new}}$. NB: Nadeau–Bengio t; CR: cluster-robust, normal quantiles; CR2 with t_{G-1}. Width ratio: NB (event folds) half-width over the floor's. Last column: median of $\hat{\sigma}^2_{\mathrm{NB}}/(\frac{J+K-1}{J-1}\hat{V}_{\mathrm{CR}})$.

G	person SD	event SD	NB, events (N6)	NB, rows (N7)	Floor	CR	CR2-t	Width ratio	NB ratio
10	0.5	0.0	0.970	0.760	0.985	0.940	0.955	0.75	1.44
10	0.5	0.4	0.975	0.720	0.965	0.890	0.930	0.80	1.41
10	1.0	0.0	0.975	0.665	0.990	0.915	0.950	0.73	1.39
10	1.0	0.4	0.960	0.665	0.980	0.895	0.925	0.74	1.35
21	0.5	0.0	0.945	0.600	0.955	0.885	0.905	0.85	1.42
21	0.5	0.4	0.940	0.595	0.955	0.880	0.900	0.81	1.44
21	1.0	0.0	0.905	0.630	0.920	0.845	0.865	0.82	1.41
21	1.0	0.4	0.930	0.630	0.945	0.880	0.905	0.84	1.36
40	0.5	0.0	0.935	0.610	0.945	0.895	0.905	0.90	1.31
40	0.5	0.4	0.880	0.515	0.880	0.770	0.795	0.89	1.29
40	1.0	0.0	0.950	0.540	0.945	0.840	0.850	0.88	1.33
40	1.0	0.4	0.935	0.555	0.940	0.845	0.870	0.89	1.34

TABLE 5. Parts 6 and 6b, balanced synthetic clusters (400 data sets per G; seeds 20264106 + G and 20264506 + G). Var ratio: $\mathrm{Var}(\widehat{\mathrm{Err}}_{\mathrm{CV}} - \mathrm{Err}_{\mathrm{new}})/\overline{\hat{V}_{\mathrm{CR}}}$ (Part 6; N8) and the same with the target's Monte Carlo variance subtracted (Part 6b; N8b). NB ratio: mean of $\hat{\sigma}^2_{\mathrm{NB}}/(\frac{J+K-1}{J-1}\hat{V}_{\mathrm{CR}})$ (N9). Corr: $\mathrm{Corr}(\widehat{\mathrm{Err}}_{\mathrm{CV}}, \mathrm{Err}_{\mathrm{new}})$ (N10). Coverage: cluster-robust t_{G-1} interval for $\mathrm{Err}_{\mathrm{new}}$ (N11). Grouped 5-fold CV.

		Var ratio		NB ratio		Corr		Coverage	
G	loss	6	6b	6	6b	6	6b	6	6b
20	0–1	1.059	0.988	1.145	1.183	0.118	0.126	0.902	0.905
20	log	1.131	1.066	1.020	1.012	−0.088	−0.167	0.875	0.880
40	0–1	0.989	0.951	1.200	1.187	0.158	0.208	0.895	0.897
40	log	1.113	1.027	1.016	0.998	−0.060	−0.165	0.882	0.895
80	0–1	1.146	1.114	1.223	1.209	0.091	0.080	0.885	0.875
80	log	1.060	1.165	1.027	1.018	−0.043	−0.063	0.882	0.882
160	0–1	1.061	1.039	1.200	1.206	0.140	0.147	0.880	0.895
160	log	1.072	1.096	1.005	0.999	−0.053	−0.063	0.887	0.882
320	0–1	1.070	0.911	1.208	1.193	0.159	0.156	0.895	0.907
320	log	1.155	0.969	0.994	0.994	−0.106	−0.021	0.890	0.915

TABLE 6. Proposition 8(ii) under 0–1 loss: the Nadeau–Bengio ratio predicted from the estimated variance components against the observed mean ratio (`firstorder_check`; bootstrap over data sets, 500 resamples, seed 20264606). †Observed value outside the prediction's 90% interval.

	Part 6 (target on 2,000 clusters)			Part 6b (20,000 clusters, MC variance removed)			
G	Predicted	90% interval	Observed	Predicted	90% interval	Observed	$\mathrm{Corr}(a, \beta)$
20	1.227	[1.198, 1.253]	1.145†	1.193	[1.173, 1.215]	1.183	−0.53
40	1.199	[1.172, 1.225]	1.200	1.196	[1.169, 1.220]	1.187	−0.49
80	1.225	[1.199, 1.251]	1.223	1.205	[1.184, 1.224]	1.209	−0.57
160	1.255	[1.228, 1.280]	1.200†	1.203	[1.180, 1.225]	1.206	−0.53
320	1.271	[1.241, 1.301]	1.208†	1.189	[1.163, 1.212]	1.193	−0.51

TABLE 7. Part 8: Nadeau–Bengio ratio $\hat{\sigma}^2_{\mathrm{NB}}/(\frac{J+K-1}{J-1}\hat{V}_{\mathrm{CR}})$ at other fold counts (400 data sets per G; seeds 20264906 + G; $J = 60$). Predictions from Part 6b's $K = 5$ components, fixed before the run. *Exploratory ($K = 2$; $K = 5$ replication).

		0–1 loss, observed			log-loss, observed (predicted 1)		
K	Predicted (0–1)	$G = 80$	160	320	$G = 80$	160	320
2*	2.112	2.184	2.251	2.223	1.056	1.051	1.029
3	1.452	1.505	1.474	1.463	1.032	1.027	0.984
5*	1.200	1.162	1.193	1.213	0.995	1.001	1.013
10	1.082	1.071	1.082	1.084	0.994	0.988	0.989

TABLE 8. Part 7, realistic designs with events redrawn (300 data sets per design; seeds 20264706 + design index; target on 2,000 new events; event 5-fold CV). Coverage of nominal 90% intervals for $\mathrm{Err}_{\mathrm{new}}$. Var ratio: $\mathrm{Var}(\widehat{\mathrm{Err}}_{\mathrm{CV}} - \mathrm{Err}_{\mathrm{new}})$, less the target's Monte Carlo variance, over the mean of $\hat{V}_{\mathrm{CR}}$. MC share: the target's Monte Carlo variance as a share of $\mathrm{Var}(\widehat{\mathrm{Err}}_{\mathrm{CV}} - \mathrm{Err}_{\mathrm{new}})$.

G	person SD	event SD	mean G^*	CR, normal (N13)	CR2, t_{G-1}	CR, t_{G^*-1}	Wild (N12)	Var ratio	MC share
21	0.5	0.0	14.0	0.897	0.907	0.913	0.910	1.053	0.009
21	0.5	0.4	13.9	0.883	0.903	0.913	0.907	0.929	0.010
21	1.0	0.0	14.0	0.823	0.857	0.860	0.853	1.115	0.008
21	1.0	0.4	14.0	0.873	0.903	0.907	0.910	0.985	0.009
40	0.5	0.0	26.6	0.883	0.893	0.893	0.893	1.034	0.018
40	0.5	0.4	26.4	0.867	0.877	0.877	0.880	1.109	0.018
40	1.0	0.0	26.6	0.867	0.877	0.883	0.887	1.018	0.019
40	1.0	0.4	26.5	0.867	0.870	0.877	0.880	1.153	0.016

shrinking (the registered diagnosis put the Monte Carlo SD at about 0.0043), against 0.0024, 0.0018, 0.0014 in Part 6b. N8b used 20,000 new clusters, subtracted the Monte Carlo variance and narrowed the band to $[0.9, 1.1]$; the corrected ratios were 0.911 to 1.165, 2 of 6 outside, so N8b was not supported either. Pooled over $G \geq 80$ and both losses the corrected ratio is 1.049 (90% bootstrap interval 0.982 to 1.105), consistent with 1. The bands were too tight for 400 data sets, where each ratio has a sampling error of about 7% ($\sqrt{2/399}$ for a sample variance). So (i) is not confirmed at the registered precision: the evidence is consistent with it but does not exclude a ratio about 10% above 1 at these G.

4.5. **Unequal cluster sizes and the wild cluster bootstrap (N12, N13).** In Part 5 the cluster-robust interval had under-covered at $G = 40$; very unequal cluster sizes are a known cause [19], and the real events are very unequal. Part 7 (Table 8) reran the realistic designs with $G \in \{21, 40\}$ events redrawn with replacement, the target on 2,000 new events, and fresh seeds. N12 was supported: the wild cluster bootstrap-t covered 0.853 to 0.910 in all 8 designs. N13, which predicted that the plain cluster-robust interval with normal quantiles would still cover below 0.85 in some $G = 40$ design, was not supported: 0.867 to 0.883 at $G = 40$ (0.823 to 0.897 over all designs). The t-based intervals covered 0.857 to 0.913, with mean G^* about 14.0 at $G = 21$ and 26.5 at $G = 40$, and the corrected variance ratio was 0.929 to 1.153. A miss of the normal interval correlated -0.114 to 0.055 with a data set's size heterogeneity. So the earlier shortfall at $G = 40$ was mostly the imprecise target plus sampling variation, not unequal event sizes.

4.6. **The Monte Carlo caveat for the target.** In a coverage study the target $\mathrm{Err}_{\mathrm{new}}$ is itself a Monte Carlo average over new clusters. Its Monte Carlo variance adds to the empirical $\mathrm{Var}(\widehat{\mathrm{Err}}_{\mathrm{CV}} - \mathrm{Err}_{\mathrm{new}})$, lowering measured coverage and raising measured variance ratios; since $\mathrm{Var}(\widehat{\mathrm{Err}}_{\mathrm{CV}} - \mathrm{Err}_{\mathrm{new}})$ shrinks like $1/G$ and the Monte Carlo variance does not, the distortion grows with G. Its share of $\mathrm{Var}(\widehat{\mathrm{Err}}_{\mathrm{CV}} - \mathrm{Err}_{\mathrm{new}})$

TABLE 9. Real data, event 5-fold CV (21 events, 4,184 dyads; headline folds, seed 20261006; NB partitions and wild bootstrap, seed 20264806). Half-widths of 90% intervals for the error rate: cluster-robust with normal and t_{20} quantiles ($t_{0.95,20} = 1.725$), wild cluster bootstrap-t, and Nadeau–Bengio t on event folds. NB ratio: $\hat{\sigma}^2_{\mathrm{NB}}/(\frac{J+K-1}{J-1}\hat{V}_{\mathrm{CR}})$. AUC with a clustered 90% interval.

			Half-width of 90% interval						
Set	Model	Error	CR, z	CR, t_{20}	Wild	NB	NB ratio	AUC	90% interval
`pre`	Logistic	0.418	0.0286	0.0300	0.0339	0.0322	1.109	0.542	[0.507, 0.578]
`pre`	LDA	0.417	0.0333	0.0350	0.0388	0.0378	1.123	0.553	[0.519, 0.587]
`pre+date`	Logistic	0.252	0.0243	0.0254	0.0260	0.0266	1.054	0.780	[0.757, 0.804]
`pre+date`	LDA	0.268	0.0221	0.0232	0.0265	0.0294	1.539	0.790	[0.769, 0.810]

was 0.001 to 0.017 with 20,000 new clusters (Part 6b) and 0.008 to 0.019 with 2,000 new events (Part 7). The realistic grids of Parts 2 to 5 scored the target on only 300 new events. The theory note puts that target's Monte Carlo SD at about 0.0045, a variance of 3% to 10% of $\mathrm{Var}(\widehat{\mathrm{Err}}_{\mathrm{CV}} - \mathrm{Err}_{\mathrm{new}})$, so the coverages of those grids are slightly understated, most at $G = 40$. With it removed, the theory note reports corrected cluster-robust variance ratios over the four confirmatory designs at each G of 0.55 to 0.84 at $G = 10$ (mean 0.72; a finite-population effect, since 10 events are drawn without replacement from 21), 0.96 to 1.20 at $G = 21$ (mean 1.04) and 0.89 to 1.46 at $G = 40$ (mean 1.16); Part 7's precise rerun resolved most of that $G = 40$ excess.

5. Real-data error bars

Table 9 applies the recommended error bars to the real headline models: the full logistic regression and LDA, on `pre` (information available before the date) and `pre+date` (adding the in-date ratings), all 4,184 dyads, event 5-fold CV on the headline folds, and the loss a wrong prediction at the training match rate. The Nadeau–Bengio variance uses $R = 10$ random event partitions. The AUC uses the clustered placement-value variance [23] with t_{20} quantiles. Because the threshold is the training match rate, as pre-registered, the error rate is not comparable with that of always predicting no match; the AUC is the more interpretable summary.

For the pre-date logistic model the error rate is 0.418 with a cluster-robust t_{20} half-width of 0.0300, and the AUC is 0.542 (0.507 to 0.578), against 0.780 on `pre+date`. In all four models the wild bootstrap-t interval is the widest of the three cluster-robust ones, and the Nadeau–Bengio ratio is 1.054 to 1.539. With balanced clusters Proposition 8(ii) would make the ratio an estimate of $\mathrm{Var}(a - \beta/4)/\mathrm{Var}(a)$, the factor by which the Nadeau–Bengio interval is conservative. The real events are unequal and the fold values are means over each fold's own rows, so the ratio also carries the fold-size normalization effect of Section 7, and the plug-in prediction of it is not reliable for these models. Each ratio is one draw from $R = 10$ partitions, and the partition noise is about as large as the differences between models: the noise variance of one $R = 10$ ratio is about 0.054, against a variance of 0.050 across the four models (Section 7). So we do not interpret their differences. The AUC intervals in Table 9 use the pooled estimate. Part 9 (Section 6) shows that this estimate runs low by order $1/G$ when clusters differ in outcome rate. On the real data, averaged over 20 random event partitions, the pooled AUC minus BCWF-pooled (the estimate Part 9 found close to unbiased) is -0.004 to $+0.002$ (Table 13), so the headline AUCs are not materially biased on average.

6. The cross-validated AUC under clustering (Part 9)

The project reports a pooled event-CV AUC: one AUC over all held-out scores, which come from K different fold models, with the clustered (Obuchowski) SE. That interval had been checked by simulation only for a fixed score. The target here is the full model's AUC on new clusters, pooled over clusters: $A(\theta) = P\{s_\theta(X_1) > s_\theta(X_0)\}$ for a match and a non-match from different clusters, with $s_\theta(x) = \tilde{x}^\top\theta$. We compare four CV estimates: the *pooled* AUC; the *fold-averaged* AUC, the mean of

the K within-fold AUCs; the *between-cluster within-fold* (BCWF) AUC, which in each fold uses only match–non-match pairs from different clusters and averages over folds; and *BCWF-pooled*, the same pairs' concordances summed over folds and divided by their count. The *pooled between-cluster* AUC (the pooled AUC without same-cluster pairs) separates two of the biases below.

6.1. **Theory.** The setup is Proposition 8's, with H the per-row Fisher information of the logistic fit and π the match rate. F_0 and S_1 are the population distribution function of non-match scores and survival function of match scores at θ^*, and $A^* = A(\theta^*)$. Cluster h's Hájek projection is $a_h = \frac{1}{m\pi}\sum_{i\in h,\, y_i=1}\{F_0(s_i^*) - A^*\} + \frac{1}{m(1-\pi)}\sum_{i\in h,\, y_i=0}\{S_1(s_i^*) - A^*\}$; $d = \nabla A(\theta^*)$ and $\beta_h = d^\top \psi_h$. Write $s_h = \sum_{i\in h}\tilde{x}_i(y_i - p_i^*)$ for the cluster's score sum and $D = s_i^* - s_j^*$ for a match i and a non-match j from different clusters; δ is the Dirac delta, so $\mathbb{E}[\delta(D)\cdot]$ weights the event $D = 0$ by the density of D. A_w and A_b are the AUCs over match–non-match pairs from the same cluster and from different clusters, and $\kappa = \mathbb{E}[m_{1h}m_{0h}]/(m^2\pi(1-\pi))$, with m_{1h} and m_{0h} the numbers of matches and non-matches in cluster h.

Proposition 9. (i) *First order.* For the pooled, fold-averaged and BCWF estimates $\hat{A}$, $\hat{A} - A(\hat{\theta}_{\text{all}}) = \bar{a} + o_p(G^{-1/2})$, and the Obuchowski variance $\hat{V}$ of the pooled scores satisfies $G\hat{V} \to_p \operatorname{Var}(a)$.

(ii) *Nadeau–Bengio.* As in Proposition 8(ii), $G\,\mathbb{E}[\hat{\sigma}^2_{\text{NB}}] \to \frac{J+K-1}{J-1}\operatorname{Var}(a - \beta/(K-1))$ for the fold AUCs. Since A does not depend on location or scale, d has no intercept part and $d \perp \theta^*$. When the model's index direction maximizes the pooled AUC among linear scores, $d = 0$; this holds under correct specification (Neyman–Pearson) and for monotone single-index truths with elliptical x [18]. Otherwise d is first order in the misspecification angle.

(iii) *Second order: three biases of order* $1/G$.

(a) *Same-cluster pairs.* A within-fold AUC puts weight about $(K/G)\kappa$ on same-cluster pairs, so the fold-averaged estimate carries $K\kappa(A_w - A_b)/G$ and the pooled estimate $\kappa(A_w - A_b)/G$, with the sign of $A_w - A_b$.

(b) *Mixing.* With $C_{\text{mix}} = \frac{1}{m}\mathbb{E}[\delta(D)\{\tilde{x}_i^\top H^{-1}s_{h(j)} - \tilde{x}_j^\top H^{-1}s_{h(i)}\}]$, $c_2 = -g'(0)$ and $g(t) = \mathbb{E}[\tilde{x}_i^\top \operatorname{Var}(\psi)\tilde{x}_j\,\delta(D-t)]$,

$$\mathbb{E}[\text{pooled between-cluster} - \text{BCWF-pooled}] = \frac{1}{G-1}\Big(C_{\text{mix}} + \frac{c_2}{K-1}\Big) + o(1/G),$$

and $C_{\text{mix}} < 0$ under outcome clustering. The next term, half the expectation of $\delta''(D)$ times the mixing shift times the cross-fold noise variance, scales with $K/(K-1)^2$ and is of order $1/G^2$.

(c) *Training size.* All CV estimates carry the bias $\operatorname{tr}(H_A \operatorname{Var}\psi)/(2(K-1)G)$, where H_A is the AUC Hessian in the slopes.

Sketch. (i) A cross-fold pair compares two models, but the AUC $A(\theta_k, \theta_{k'})$ expanded to first order sums, over the folds, to $\nabla A^\top(\hat{\theta}_{\text{all}} - \theta^*)$; the intercept parts cancel because the AUC does not depend on location. (ii) follows from Proposition 6 as in Proposition 8(ii). (iii)(b) In a cross-fold pair (i in fold k, j in fold k'), model k was fitted on j's cluster and model k' on i's, so the score difference is shifted by $\{\tilde{x}_i^\top H^{-1}s_{h(j)} - \tilde{x}_j^\top H^{-1}s_{h(i)}\}/n'$ with $n' = n(K-1)/K$; a within-fold pair from different clusters has no shift. Given a match in fold k, a non-match from another cluster lies in another fold with probability $G(K-1)/(K(G-1))$; times $1/n'$ this is $1/(m(G-1))$, so the leading term is $C_{\text{mix}}/(G-1)$, free of K. The two fold models also differ by noise $\xi = -\tilde{x}_j^\top(\theta_{k'} - \theta_k)$ correlated with the common error. A second-order Taylor step in the Heaviside function gives, per cross-fold pair,

$$\mathbb{E}[\delta'(D)\{\operatorname{Cov}(\varepsilon_0, \xi) + \tfrac{1}{2}\operatorname{Var}\xi\}] = \frac{K}{K-1}\cdot\frac{\mathbb{E}[\delta'(D)\,\tilde{x}_i^\top \operatorname{Var}(\psi)\tilde{x}_j]}{(K-1)G},$$

and times the cross-fold share this is $c_2/((K-1)(G-1))$. □

Parts (i) and (ii) are derived as in Proposition 8. Part (iii) is a *heuristic second-order expansion*, not a proved result: a formal expansion with no remainder bound, which assumes a smooth density for D,

enough moments and balanced clusters. Its constants are computed numerically from the population. An adversarial review of Propositions 9 and 10 by an automated reviewer (`proposals/09`) confirmed the first- and second-order terms order by order and made two corrections that are folded in above: the factor is $1/(G-1)$, not the $1/G$ of the first version, which used a cross-fold share of $(K-1)/K$ (a difference of $o(1/G)$, but 1% to 2.6% at the G used); and the per-pair identity needs the factor $K/(K-1)$ before the cross-fold share is applied. The mixing bias is the pooling bias of [25, 15] seen through clustering: with independent rows it is of order $1/n$, with clusters of order $1/G$. Its leading term does not shrink with more folds; only the c_2 part does. By exact quadrature (review), c_2/C_{mix} is 0.50000, 0.50000, 0.50183, 0.50027, 0.50104 and 0.50316 in the six linear designs and 0.52 to 0.53 with random slopes. The binned estimates in `sim_cv_auc_components.csv` cannot resolve deviations below about 1%.

Proposition 10 (independent rows). Suppose the rows are independent, the logistic model is correctly specified, and the score densities are smooth. Then $c_2 = C_{\text{mix}}/2$ exactly, and to leading order the pooling bias of K-fold CV is

$$-\frac{2K-1}{2(K-1)} \cdot \frac{1}{n}\,\mathbb{E}[\delta(D)\,\tilde{x}_i^\top H^{-1}\tilde{x}_j]$$

plus the training-size term.

Proof. (1) With independent rows, $\mathbb{E}[s_{h(j)} \mid j] = -p_j\tilde{x}_j$ for a non-match and $(1-p_i)\tilde{x}_i$ for a match, and $\operatorname{Var}\psi = H^{-1}/m$. On $D=0$ the two scores are equal, so $p_i = p_j$ and $1 - p_i + p_j = 1$. Hence $C_{\text{mix}} = -\frac{1}{m}\mathbb{E}[\delta(D)\,q]$ with $q = \tilde{x}_i^\top H^{-1}\tilde{x}_j$, and $c_2 = -\frac{1}{m}I'(0)$ with $I(t) = \mathbb{E}[q\,\delta(D-t)]$. (2) Under correct specification $P(y=1 \mid x)$ depends on x only through the score s, so x is independent of y given s. With $\tilde{\mu}(s) = \mathbb{E}[\tilde{x} \mid s]$ and the symmetric $Q(a,b) = \tilde{\mu}(a)^\top H^{-1}\tilde{\mu}(b)$,

$$I(t) = \int f_1(s+t)f_0(s)Q(s+t,s)\,ds = \int f_1(u)f_0(u-t)Q(u,u-t)\,du.$$

(3) Average the derivatives of the two forms at $t=0$. Because Q is symmetric the ∂Q terms cancel, leaving $I'(0) = \frac{1}{2}\int (f_1'f_0 - f_1f_0')\,Q(s,s)\,ds = \frac{1}{2}\int f_1f_0\,(\operatorname{logit} P(y=1 \mid s))'\,Q(s,s)\,ds$. (4) Under a correct logistic model $(\operatorname{logit} P)' = 1$, so $I'(0) = \frac{1}{2}\mathbb{E}[\delta(D)q]$ and $c_2 = C_{\text{mix}}/2$. With $m(G-1) \approx n$, the bias $(C_{\text{mix}} + c_2/(K-1))/(G-1)$ is the display. □

Remark. This shorter proof, by conditioning on the score, is the review's. The first version assumed Gaussian covariates and kept the factor $(1-p_i+p_j)$, which is identically 1 on $D=0$; neither is needed. In design 0 (no clustering, correct model) exact quadrature gives 0.500000, against 0.4985 from the binned estimator, and with logistic, t_5, Laplace, bimodal and chi-square covariates the review found 0.49993 to 0.49999. Under misspecification the identity is off by a term proportional to $\nabla A(\theta^*)$, which is small (theory note).

Lemma (the clustered case) (review, `proposals/09`). Assume a single index (y independent of x given s) and a correct marginal logistic model, and let $B(x) = \mathbb{E}[s_h \mid x, y=1] - \mathbb{E}[s_h \mid x, y=0]$. Then $c_2 = C_{\text{mix}}/2$ whenever B is affine in $\tilde{x}$.

Independent rows give $B = \tilde{x}$, which is Proposition 10. With clusters, B also carries the cluster mates' conditional score difference, which is not affine in general, and the identity is *false in general*. The review's counterexamples, by exact quadrature: a two-point random intercept of ± 3 gives 0.454 (and 0.452 to 0.456 with the project's binned code on fresh seeds, checked independently), ± 4 gives 0.422, a Gaussian intercept with strongly shared covariates ($s_a = 1$, $s_u = 3$, $m = 400$) gives 0.475 to 0.563, and a bridge-distributed intercept, whose marginal model is exactly logistic, gives 0.507. With a Gaussian random intercept the mates' term is nearly affine, so the ratio minus 1/2 is $O(s_a^2)$ with a tiny coefficient (1.4×10^{-3} at $s_a = 0.8$); that is why our designs come out near 1/2. Under clustering the closed form $C_{\text{mix}}(2K-1)/(2(K-1))$ for $G-1$ times the mixing bias is therefore an approximation, with error under 0.4% in our designs (review). The quadrature values are reproduced by `scripts/auc_ratio_quadrature.py` (`results/auc_ratio_quadrature.csv`), adopted from the

TABLE 10. Part 9, mixing bias: $G\times$(pooled between-cluster AUC − BCWF-pooled), observed mean (SE), against the registered prediction $C_{\text{mix}} + c_2/(K-1)$ and the corrected one, $G/(G-1)$ times that (review, `proposals/09`, E1). $K = 5$: mean over $G = 40, 80, 160$ (N17); $K = 2$ and 10: $G = 80$ (N18). Binned constants from `sim_cv_auc_components.csv`; 400 data sets per cell. †Outside the registered 15% band.

				Predicted			Rel. error	
K	(s_a, s_u)	C_{mix}	c_2	registered	$\times G/(G-1)$	Observed (SE)	registered	corrected
5	1, 0	−0.267	−0.133	−0.300	−0.305	−0.304 (0.008)	+1.3%	−0.2%
5	1, 1	−0.210	−0.103	−0.236	−0.239	−0.242 (0.011)	+2.5%	+1.0%
5	2, 0	−0.753	−0.374	−0.846	−0.859	−0.865 (0.021)	+2.2%	+0.7%
5	2, 1	−0.629	−0.315	−0.708	−0.719	−0.745 (0.022)	+5.1%	+3.6%
2	1, 0	−0.267	−0.133	−0.400	−0.405	−0.397 (0.024)	−0.8%	−2.1%
2	2, 0	−0.753	−0.374	−1.127	−1.142	−0.917 (0.060)	−18.6%†	−19.7%
10	1, 0	−0.267	−0.133	−0.282	−0.285	−0.281 (0.014)	−0.3%	−1.5%
10	2, 0	−0.753	−0.374	−0.794	−0.805	−0.783 (0.030)	−1.4%	−2.7%

review and tested against the binned constants and against Proposition 10; the Lemma's derivation is in `proposals/09`.

Which target matters depends on the use: for ranking pairs across events (an app) it is the pooled between-cluster AUC, which BCWF estimates without the mixing and same-cluster biases; for ranking within an event (a matchmaker) it is A_w, estimated by a within-cluster AUC.

6.2. **Simulation evidence (N16 to N21).** Part 9 used clusters of 40 rows with $x = s_u u_g + v_i$ and $y \sim \text{Bernoulli}(\text{logistic}(-1.5 + x^\top(1, 0.5, 0) + s_a a_g))$; designs 0 to 5 cross $s_a \in \{0, 1, 2\}$ with $s_u \in \{0, 1\}$, and the Nadeau–Bengio designs ($s_a = s_u = 1$) are linear, hinge ($+1.5\max(x_3, 0) - 1$) and random slope (x_1 slope $1 + b_g$). The model is a logistic regression on x, and AUC_{new} is scored on 20,000 new clusters (Monte Carlo share of the variance at most 0.008). The constants and predictions were computed before any CV run, and every test used fresh seeds. `PREREG.md` records two labelled pilots run before registration (scratch outputs, not in `results/`).

N16 was supported: in the 12 cells with outcome clustering ($s_a \in \{1, 2\}$, $G \le 80$) the pooled bias was −0.004 to −0.040, with the largest 90% upper limit −0.0022. G times the bias averaged −0.81 with $s_a = 2$ and −0.29 with $s_a = 1$, so the bias is of order $1/G$; at $G = 20$ with $s_a = 2$, $s_u = 0$ it was 1.07 SE. With no clustering at all (design 0) |bias| was at most 0.0028, but covariate clustering alone ($s_a = 0$, $s_u = 1$) gave −0.0064 at $G = 20$. N17 was supported: at $K = 5$ the observed $G\times$ mixing bias was within 1.3% to 5.1% of the registered prediction $C_{\text{mix}} + c_2/4$ in all four designs (Table 10). With the corrected factor $G/(G-1)$ and the binned constants the gaps are −0.2%, +1.0%, +0.7%, +3.6%; with the review's exact constants they become −1.0%, −0.8%, +0.5% and +3.7% (`proposals/09`). N18 was *not* supported, and that verdict stands: at $G = 80$ three of the four cells at $K = 2$ and 10 were within 1.4% of $C_{\text{mix}} + c_2/(K-1)$, but at $K = 2$ with $s_a = 2$ the registered prediction was −1.127 against an observed −0.917 (SE 0.060), off by 18.6% against a 15% band (19.7% against −1.142 with the factor $G/(G-1)$). The review (`proposals/09`, E2) traced most of the miss to sampling error. At $K = 2$ the statistic is roughly minus a squared difference of two fold means, so it is heavily skewed (skewness −2.4), and 400 data sets understate its SE. An exploratory rerun with 10,000 data sets on fresh seeds (seeds 20265906 + chunk, reported in `proposals/09` and the theory note; it has no CSV in `results/`) gave −1.075 (SE 0.014) against −1.142, a 5.8% shortfall; the reviewer's own rerun gave 7%. The rest is the third-order term, which scales with $K/(K-1)^2$ and decays with G; $K = 2$ was also the weakest fold count in Part 8.

N19 was supported: over the 18 cells with $G \ge 40$, BCWF-pooled had |bias| at most 0.11 SE, while the pooled estimate's bias was 0.48 to 0.87 SE in the $s_a = 2$ cells at $G = 40$ and 80. N20 was supported (Table 11): the BCWF-pooled interval (t_{G-1}, Obuchowski SE) covered 0.899 on average over the 18

TABLE 11. Part 9, coverage of nominal 90% intervals for $\mathrm{AUC}_{\mathrm{new}}$ (t_{G-1}, Obuchowski SE of the pooled scores; 400 data sets per cell; event 5-fold CV; seeds 20265106 + 10 × design + G index). Second column: $G\times$ the pooled estimate's bias, averaged over G.

		Pooled				BCWF-pooled			
(s_a, s_u)	$G\times$bias	$G = 20$	40	80	160	$G = 20$	40	80	160
0, 0	−0.08	0.890	0.910	0.912	0.868	0.890	0.910	0.912	0.885
0, 1	−0.10	0.890	0.877	0.885	0.907	0.853	0.882	0.875	0.910
1, 0	−0.25	0.825	0.870	0.858	0.907	0.877	0.920	0.887	0.915
1, 1	−0.32	0.868	0.855	0.895	0.892	0.863	0.848	0.882	0.890
2, 0	−0.80	0.705	0.782	0.830	0.860	0.910	0.915	0.938	0.917
2, 1	−0.81	0.797	0.828	0.873	0.873	0.823	0.887	0.927	0.880

TABLE 12. Part 9, Nadeau–Bengio ratio $\hat{\sigma}^2_{\mathrm{NB}}/(\frac{J+K-1}{J-1}\,\hat{V}_{\mathrm{Obu}})$ for within-fold AUCs ($R = 12$, $J = 60$; 400 data sets per cell; $s_a = s_u = 1$), mean (SE), against Proposition 9(ii)'s prediction (N21). Last two columns: the same ratio for per-fold BCWF AUCs (exploratory).

Design	Predicted	$G = 80$	$G = 160$	BCWF, $G = 80$	BCWF, $G = 160$
Linear	1.000	0.988 (0.010)	0.991 (0.010)	1.059	1.025
Hinge	1.003	0.999 (0.010)	0.994 (0.010)	1.056	1.021
Random slope	0.991	0.979 (0.010)	0.978 (0.009)	1.021	0.999

cells with $G \geq 40$ (0.848 to 0.938), and the pooled interval 0.841 on average in the $s_a = 2$ cells with $G \geq 40$, down to 0.705 at $G = 20$. The wild cluster bootstrap does not rescue the pooled interval (0.695 in that cell), because the problem is bias, not variance. N21 was supported: the Nadeau–Bengio ratio for fold AUCs was 0.978 to 0.999, within 0.013 of its predictions of 0.991 to 1.003 (Table 12). In every design computed $|d| \leq 0.014$, so unlike the error rate's ratio of about 1.2 the AUC's Nadeau–Bengio correction is not conservative.

Exploratory results. Where the same-cluster term dominates ($s_a = 0$, $s_u = 1$) the fold-averaged bias followed $(c_{\mathrm{train}} + K\kappa(A_w - A_b))/G$: −0.0150 against −0.0137 at $G = 20$ and −0.0015 against −0.0017 at $G = 160$. With $s_a = 2$, $s_u = 0$ at $G = 20$, per-fold BCWF was biased by 0.0079 against 0.0011 for BCWF-pooled, the per-fold ratio bias that pooling the counts removes. In the realistic A2 design with 21 redrawn events (300 data sets per design), the pooled interval covered 0.770 to 0.873 and BCWF-pooled 0.847 to 0.870, with biases of −0.004 to −0.015 against at most 0.002 in absolute value.

6.3. **Real data.** Table 13 gives the four estimates for the real headline models, on the headline folds and averaged over 20 random event partitions (seed 20265406), with the within-event AUC, the matchmaker's target, for reference. On the headline folds the pooled AUC minus BCWF-pooled is −0.003 to −0.012; averaged over the random partitions it is −0.004 to +0.002, with an SD across partitions of 0.005 to 0.009. So the reported AUCs are not materially biased on average, but the gap on any one partition varies by more than its mean.

7. Unequal cluster sizes (Part 11)

Propositions 6 and 8 assume balanced clusters and $K \mid G$, but the 21 real events hold 25 to 484 dyads. The scripts use three things that balance makes trivial: the pooled estimate $\widehat{\mathrm{Err}}_{\mathrm{CV}}$, which weights every row equally and is therefore a ratio estimator; the unbalanced form of $\hat{V}_{\mathrm{CR}}$; and Nadeau–Bengio fold values that are each fold's mean loss over its own rows (*own-row* fold means). With unequal clusters the folds of a grouped partition hold different numbers of rows, so the training sets differ in size, and a partition's fold means no longer average to the pooled estimate. Section 6g of the theory

TABLE 13. Real data, event 5-fold CV: the four CV AUC estimates on the headline folds and their mean over 20 random event partitions (seed 20265406; exploratory). Gap: pooled − BCWF-pooled, with its SD over partitions. Within: AUC over pairs from the same event.

Set	Model	Folds	Pooled	Fold-avg.	BCWF	BCWF-pooled	Gap (SD)	Within
`pre`	Logistic	headline	0.542	0.544	0.542	0.546	−0.003	0.539
		mean of 20	0.544	0.553	0.555	0.545	−0.001 (0.009)	0.532
`pre`	LDA	headline	0.553	0.558	0.559	0.559	−0.006	0.544
		mean of 20	0.555	0.562	0.568	0.558	−0.004 (0.007)	0.540
`pre+date`	Logistic	headline	0.780	0.781	0.775	0.793	−0.012	0.800
		mean of 20	0.790	0.794	0.788	0.788	0.002 (0.005)	0.801
`pre+date`	LDA	headline	0.790	0.799	0.798	0.797	−0.007	0.794
		mean of 20	0.794	0.799	0.799	0.795	−0.001 (0.005)	0.795

note extends Proposition 8 to this case with cluster totals in place of cluster means; this section states the result and its registered test.

7.1. **Theory.** Cluster h has m_h rows, $n = \sum_h m_h$, $\bar{m} = \mathbb{E}[m]$, and $\tilde{m} = \mathbb{E}[m^2]/\mathbb{E}[m]$ is the size-biased mean size ($\bar{m} = 199.2$ and $\tilde{m} = 302.5$ for the real events). Let $L_h(\theta) = \sum_{i \in h} \ell(z_i; \theta)$ be cluster h's loss total. The target is row-weighted: $A(\theta) = \mathbb{E}[L_h(\theta)]/\bar{m}$, the expected loss of a row drawn at random from the rows of new clusters, and $\mathrm{Err}_{\mathrm{new}} = A(\hat{\theta}_{\mathrm{all}})$. This is what the simulations score when they pool the loss over new events. The cluster-weighted target $\mathbb{E}[L_h(\theta)/m_h]$, estimated by the unweighted mean of the events' CV error rates, coincides with it only when size carries no information about the rows; in the A2 designs it does, because each event template has its own covariates. The M-estimator weights rows equally: $\hat{\theta}_T$ solves $\sum_{g \in T} S_g(\theta) = 0$ with S_g the cluster's score total, H is the per-row information, and $\psi_g = H^{-1} S_g(\theta^*)$ is a total. With $\mu = A(\theta^*)$, $a_h = L_h(\theta^*) - m_h \mu$, $d = \nabla A(\theta^*)$ and $\beta_h = d^\top \psi_h$, every variance below is a variance of totals. Fold j has n_j rows, and $N_j = n - n_j$.

Assumptions for Proposition 11. (U1) The pairs $(m_h, \text{rows of } h)$ are i.i.d., with $\mathbb{E}[m^4] < \infty$ and $\mathbb{E}\|\nabla L_h(\theta^*)\|^2 < \infty$. (U2) The partition deals the clusters at random into K folds of $\lfloor G/K \rfloor$ or $\lceil G/K \rceil$ clusters, independently of the data, so the n_j are random; K is fixed as $G \to \infty$. Assumptions (2) to (4) for Proposition 8 hold with the linearization $\hat{\theta}_T - \theta^* = \sum_{g \in T} \psi_g / n_T + O_p(1/G)$, $n_T = \sum_{g \in T} m_g$; for 0–1 loss the threshold's influence total is $Y_g - m_g t^*$.

Proposition 11 (pooled grouped K-fold CV, row-weighted target)**.** *(a) To first order, with $\omega_h = \sum_{j \neq j(h)} n_j/N_j$, cluster h's total weight over the training sets that contain it,*

$$\widehat{\mathrm{Err}}_{\mathrm{CV}} = \mu + \frac{1}{n} \sum_h (a_h + \omega_h \beta_h) + r_1, \qquad \mathrm{Err}_{\mathrm{new}} = \mu + \frac{1}{n} \sum_h \beta_h + r_2.$$

With $n_j/n = 1/K + \varepsilon_j$, $\omega_h = 1 - \frac{K^2}{(K-1)^2} \varepsilon_{j(h)} + O_p(1/G)$, and $\omega_h \equiv 1$ when the folds have equal row counts. The term $\frac{1}{n} \sum_h (\omega_h - 1)\beta_h$ is $O_p(1/G)$, so $\widehat{\mathrm{Err}}_{\mathrm{CV}} - \mathrm{Err}_{\mathrm{new}} = \frac{1}{n} \sum_h a_h + o_p(G^{-1/2})$.

(b) $G \operatorname{Var}(\widehat{\mathrm{Err}}_{\mathrm{CV}} - \mathrm{Err}_{\mathrm{new}}) \to \operatorname{Var}(a)/\bar{m}^2$ (with uniform integrability), the linearized variance of a ratio estimator, and $G\hat{V}_{\mathrm{CR}} \to_p \operatorname{Var}(a)/\bar{m}^2$. If sizes carry no information about the rows, and the losses at θ^* have per-row variance σ_a^2 and within-cluster correlation ρ_a, then $\operatorname{Var}(\widehat{\mathrm{Err}}_{\mathrm{CV}} - \mathrm{Err}_{\mathrm{new}}) \approx \sigma_a^2\{1 + (\tilde{m} - 1)\rho_a\}/n$: Kish's design effect with m replaced by $\tilde{m}$.

(c) In expectation, $G\,\mathbb{E}[\hat{\sigma}^2_{\mathrm{NB}}] \to \frac{J+K-1}{J-1} \operatorname{Var}(a - \beta/(K-1))/\bar{m}^2$, *so*

$$\frac{\mathbb{E}[\hat{\sigma}^2_{\mathrm{NB}}]}{\frac{J+K-1}{J-1}\,\mathbb{E}[\hat{V}_{\mathrm{CR}}]} \to \frac{\operatorname{Var}(a - \beta/(K-1))}{\operatorname{Var}(a)},$$

whether the fold values are own-row means or residual totals divided by n/K. The per-data-set ratio does not converge in probability at fixed J.

(d) Proposition 8(iii) holds with totals.

(e) Proposition 8(iv) holds without balance for the row-weighted target: if the evaluation loss is the training criterion, training and target weight rows the same way, so θ^* minimizes A and $d = 0$, even when the model is misspecified (for example by an omitted random intercept).

Derivation sketch. The steps are those of Proposition 8 with totals; the theory note gives them in full. Step (3) changes: fold j contributes $\frac{1}{n}[\sum_{h\in F_j}(a_h + m_h\mu) + n_j B_{T_j}/N_j]$, with $B_T = \sum_{g\in T}\beta_g$, and $\sum_j (n_j/N_j)B_{T_j} = \sum_h \omega_h\beta_h$. Expanding $p/(1-p)$ about $p = 1/K$ gives ω_h, and $\varepsilon_j = O_p(G^{-1/2})$ makes the weight term $O_p(1/G)$. For (c), the step of Proposition 6 that fails is that a partition's fold values average to $\widehat{\mathrm{Err}}_{\mathrm{CV}}$: with own-row means $\mathbb{E}[S^2] = (Jv - u)/(J-1)$, where v is the variance of one fold value and u is K times the variance of a partition's mean fold value. The pooled estimate varies across partitions only at order $G^{-3/4}$ for 0–1 loss and $1/G$ for smooth losses, so $u = o(v)$; with residual totals $u = 0$ exactly. Proposition 6 applied to the totals $a_h - \beta_h/(K-1)$ then gives (c). □

Parts (a), (b), (d) and (e) are first-order results under (U1), (U2) and the assumptions for Proposition 8, and (c) holds in expectation only; the reduction to $\tilde{m}$ in (b) needs sizes that carry no information about the rows. The order-$1/G$ terms are not small at $G = 21$: with the real sizes, $\max_h |\omega_h - 1|$ has median 0.17 over partitions at $K = 5$ (theory note).

Two predictions. P1 is the limit in (c), computed from the population constants of the synthetic design below: for 0–1 loss 1.260 at $K = 5$, 1.107 at $K = 10$ and 1.586 at $K = 3$, and for log-loss 1 by (e). *P2* is a first-order heuristic at finite G. Condition on the sizes and the partitions and keep only the linear terms of (a): cluster h's CV loss total is $m_h\mu + a_h + m_h B_{T(h)}/N_{(h)}$, and fold j's own-row mean is $\mu + A_{F_j}/n_j + B_{T_j}/N_j$, with $A_F = \sum_{h\in F} a_h$. Then $\widehat{\mathrm{Err}}_{\mathrm{CV}}$, $\mathrm{Err}_{\mathrm{new}}$, $\hat{V}_{\mathrm{CR}}$ and $\hat{\sigma}^2_{\mathrm{NB}}$ are quadratic forms in the independent totals (a_h, β_h), and P2 is the mean of the per-data-set ratio over 40,000 Gaussian draws of the totals at the actual sizes and partitions (`scripts/prop11_linear.py`). P2 keeps some order-$1/G$ terms: the weights ω_h, the leverage term of $\hat{V}_{\mathrm{CR}}$, the covariance of a_h with its own fold's training shift, and the division by each fold's own row count. It omits others of the same or larger order: the pooled estimate's variation across partitions, through flip noise for 0–1 loss and curvature for smooth losses, which adds about $\mathrm{Var}_P(\widehat{\mathrm{Err}}_{\mathrm{CV}})/((K-1)\hat{V}_{\mathrm{CR}})$ to the own-row ratio (by the theory note's estimate +0.006 to +0.008 at $K = 5$, about +0.03 at $K = 3$, and nothing with residual totals); the test cluster's own gradient deviation times its fold's training deviation; and curvature in the fold values and in $\mathrm{Err}_{\mathrm{new}}$. As a mean of ratios it also depends on the Gaussian law assumed for the totals. So P2 is a heuristic, not a second-order expansion. For log-loss ($\beta \equiv 0$) it has a closed form: with the real sizes and $\mathrm{Var}(a_h \mid m_h) \propto m_h^2$, the leverage bias of the CR1 form puts $\mathbb{E}[\hat{V}_{\mathrm{CR}}]$ at 0.945 times $\mathrm{Var}(\widehat{\mathrm{Err}}_{\mathrm{CV}} - \mathrm{Err}_{\mathrm{new}})$, and own-row normalization puts the Nadeau–Bengio numerator at 1.009, 0.982 and 0.869 times its target at $K = 3$, 5 and 10. The algebra was checked against known answers (`tests/test_sim_cv_unequal.py`): the linear representation matches a separate implementation to 10^{-14} and reproduces Proposition 6 for fixed totals.

7.2. **Simulation evidence (N22 to N25).** Part 11 used Part 6's generator, with a logistic regression on x, and cluster sizes from the real events: *S21f*, the 21 real sizes as they are (2,000 data sets), and *S84r*, 84 sizes drawn with replacement from them (1,000 data sets). Here $\mathrm{Err}_{\mathrm{new}}$ is the full model's exact row-weighted population loss (400,000 rows from the row marginal, with $P(y = 1 \mid x)$ by quadrature), so it carries no Monte Carlo noise. Two A2 designs (person SD 0.5 with event SD 0; person SD 1.0 with event SD 0.4) redrew 21 events from the real templates (500 data sets each), with $\mathrm{Err}_{\mathrm{new}}$ on 2,000 new events. In every data set one grouped 5-fold run gave $\widehat{\mathrm{Err}}_{\mathrm{CV}}$ and $\hat{V}_{\mathrm{CR}}$, and $J = 60$ fold values ($R = 20$, 12 and 6 partitions at $K = 3$, 5 and 10) gave $\hat{\sigma}^2_{\mathrm{NB}}$ in three versions: own-row means, as in the scripts; residual totals about the partition's pooled estimate, divided by n/K; and that residual value plus the partition's pooled estimate. The own-row minus residual-total difference then splits into a *normalization part* (own-row minus the third version) and a *partition part* (the third version minus

TABLE 14. Part 11, registered tests: mean Nadeau–Bengio ratio $\hat{\sigma}^2_{\mathrm{NB}}/(\frac{J+K-1}{J-1}\hat{V}_{\mathrm{CR}})$ over data sets, with its SE (S21f: 2,000 data sets; S84r: 1,000; A2: 500 per design; $J = 60$; $\hat{V}_{\mathrm{CR}}$ from one grouped 5-fold run; seeds 20267106 + 100 × design index + chunk for S21f and S84r, 20267306 + 100 × design index + chunk for A2). Fold values: own-row means (own) or residual totals divided by n/K (res.). P1 and P2 fixed before the run (seeds 20267006 and 20267016; 40,000 draws). Distance: observed minus the value named in the band, before rounding. A2 designs are labelled by person SD and event SD. [a]Mean per-data-set plug-in P2 (in-sample residual totals, jackknife $\hat{\beta}$). [b]P1 estimated from the moments of $\widehat{\mathrm{Err}}_{\mathrm{CV}}$ and $\mathrm{Err}_{\mathrm{new}}$ across data sets, as in Table 6.

Hyp.	Design	Loss	K	Folds	Observed (SE)	P2	P1	Distance	Band	Verdict
N22	S21f	0–1	10	own	1.007 (0.007)	0.998	1.107	+0.009	P2 ±0.04	supported
	S21f	0–1	5	own	1.230 (0.006)	1.217	1.260	+0.013	P2 ±0.03; < 1.2385	
N23	S84r	0–1	5	own	1.266 (0.008)	1.251	1.260	+0.006	P1 ±0.03	supported
	S84r	0–1	10	own	1.078 (0.006)	1.092	1.107	−0.029	P1 ±0.03	
N24	A2, 0.5, 0	0–1	5	own	1.406 (0.016)	1.414[a]	1.270[b]	−0.009	plug-in ±0.08	supported
	A2, 0.5, 0	0–1	5	own − res.	0.192 (0.011)	–	–	–	≥ 0.10	
	A2, 1.0, 0.4	0–1	5	own	1.477 (0.019)	1.410[a]	1.275[b]	+0.068	plug-in ±0.08	
	A2, 1.0, 0.4	0–1	5	own − res.	0.203 (0.011)	–	–	–	≥ 0.10	
N25	S21f	log	5	res.	1.007 (0.005)	0.998	1	+0.007	1 ±0.03	supported
	S21f	log	5	own	1.092 (0.006)	1.065	1	+0.027	P2 ±0.03	
	S21f	log	10	res.	0.994 (0.004)	0.997	1	−0.006	1 ±0.03	
	S21f	log	10	own	0.993 (0.007)	0.984	1	+0.009	P2 ±0.03	

residual totals), the latter being the pooled estimate's variation across partitions that P2 omits. P1 and P2 were computed before any CV run. In the A2 designs sizes carry information, so N24 compares the observed ratio with a per-data-set plug-in P2 (in-sample residual totals, jackknife $\hat{\beta}$), and P1 is estimated from the moments of $\widehat{\mathrm{Err}}_{\mathrm{CV}}$ and $\mathrm{Err}_{\mathrm{new}}$ across data sets, as in Table 6. `PREREG.md` records a labelled exploratory pilot (scratch seeds 997000 to 997610) and an adversarial review with its own checks (scratch seeds 998100 to 998300), both before registration; every test used fresh seeds.

All four hypotheses were supported (Table 14), but three of the four came close to their band edges: N23 at $K = 10$ was 0.029 from P1 against a band of 0.03; N24 in the second A2 design missed its plug-in by +0.068 against a band of 0.08, 3.8 SE from zero; and N25's own-row ratio at $K = 5$ was 0.027 above P2 against 0.03. N22, the test that separates P2 from P1, was 1.007 (SE 0.007) at $K = 10$, against 0.998 for P2 and 1.107 for P1. At $K = 5$ it was 1.230 (SE 0.006) against P2's 1.217, and 0.008 (1.3 SE) below the P1/P2 midpoint of 1.2385; its excess over P2, +0.013, is about the omitted partition part, measured at +0.009. At $G = 84$ (N23) the ratio was 1.266 (SE 0.008) against P1's 1.260 at $K = 5$, but 1.078 (SE 0.006) at $K = 10$, below both P1 (1.107) and P2 (1.092). In the A2 designs (N24) the own-row ratios were 1.406 and 1.477, and exceeded the residual-total ratios by 0.192 and 0.203 (SE 0.011), against the registered minimum of 0.10. Under log-loss (N25) the residual-total ratios were 1.007 and 0.994, and the own-row ratios 1.092 and 0.993 against P2's 1.065 and 0.984. What this supports: with unequal clusters and own-row fold means, the Nadeau–Bengio ratio at $G = 21$ is set mainly by the actual sizes and partitions (P2), not by the first-order limit, which is approached at $G = 84$, at $K = 10$ only loosely. P2's own misses (+0.013 at $K = 5$ under 0–1 loss, +0.027 under log-loss) are of the size of the terms it omits.

Exploratory results. $K = 3$, which the pilot had flagged, is not predicted well. In S21f the own-row ratio was 1.573 (SE 0.009) under 0–1 loss, against 1.555 for P2 and 1.586 for P1, and 1.151 (SE 0.006) under log-loss, against 1.075 for P2. The partition part that P2 omits was 0.044 and 0.041 there, against 0.009 and 0.008 at $K = 5$. Under log-loss even the residual-total ratio, which the partition part does not enter, was 1.030 (SE 0.005), 0.029 above P2. A second-order surrogate for log-loss, which adds the test cluster's own score term and curvature and was computed before the run, gave

1.163, 1.088 and 0.983 at $K = 3$, 5 and 10, against 1.151, 1.092 and 0.993 observed. *The partition and normalization split.* In S21f at $K = 5$ the own-row minus residual-total difference was 0.032 (SE 0.004) under 0–1 loss, of which 0.009 was the partition part and 0.023 the normalization part, and 0.084 (SE 0.004) under log-loss (0.008 and 0.076). At $K = 10$ the normalization part under 0–1 loss was -0.047, so own-row means lowered the ratio. In the A2 designs the normalization part was 0.178 in both and the partition part 0.014 and 0.025, so normalization accounts for nearly all of the 0.19 to 0.20 that own-row means add, and the residual-total ratios, 1.213 and 1.274, are closer to the P1 estimates of 1.270 and 1.275. At $G = 84$ the whole difference was 0.008 (S84r, 0–1 loss, $K = 5$). *The variance ratio.* Under 0–1 loss $\mathrm{Var}(\widehat{\mathrm{Err}}_{\mathrm{CV}} - \mathrm{Err}_{\mathrm{new}})/\overline{\hat{V}_{\mathrm{CR}}}$ was 0.992 in S21f (P2: 0.991), and 1.028 and 0.948 in the A2 designs with the target's Monte Carlo variance removed. Under log-loss with the 21 real sizes it was 1.200 (P2: 1.047), so the mean of $\hat{V}_{\mathrm{CR}}$ was 0.83 of the variance it estimates, about 17% low; at $G = 84$ it was 1.035 (P2: 0.999). $G\,\mathrm{Var}(\widehat{\mathrm{Err}}_{\mathrm{CV}} - \mathrm{Err}_{\mathrm{new}})$ was 0.0619 there, 28% above P2's 0.0482; the second-order surrogate gives 0.0580.

Real data. The A2 normalization effect does not carry over to the real data. Over 40 random event partitions, the own-row minus residual-total ratios of the four headline models were +0.088, +0.055, +0.039 and −0.030, below A2's mean but within its per-data-set spread (SD 0.23). Splitting the 40 partitions into four groups of 10 gives a noise variance of about 0.054 for one $R = 10$ ratio, about the variance across the four models' reported values in Table 9 (0.050). At $R = 40$ the ratios were 0.93 to 1.36; removing the noise leaves a between-model SD of about 0.17, so the models do differ, but at $R = 10$ the noise is as large as the differences. The plug-in is not reliable for the real models, which have dozens of features against 21 events: in-sample residual totals carry only 28% to 53% of the cross-validated residual sum of squares, and with cross-validated residual totals and a centred $\hat{\beta}$ its predictions (1.15 to 1.30) missed the $R = 40$ ratios by −1.1 to +4.8 partition SDs. We claim no quantitative real-data check. These numbers are from the theory note, Section 6g (scratch scripts; no CSV in `results/`).

An adversarial review of Proposition 11 and the pilot by an automated reviewer (3 November 2026, before registration) made three corrections that are folded in above: P2 omits the pooled estimate's variation across partitions; (c) holds in expectation, not in probability; and an expansion of the normalization effect in the fold-size deviation, accurate for small imbalance, fails at the real sizes. Part 5's ratios (Table 4; $J = 50$, $\hat{V}_{\mathrm{CR}}$ averaged over 10 nested repetitions) are not like-for-like with these designs, and we do not claim to explain them.

8. Recommendations

1. *Hold out whole clusters.* This fixes what is estimated, the error on new clusters (Lemma 1); no interval repairs random-row CV when the question concerns new people.
2. *Use the cluster-robust SE with t quantiles on $G - 1$ (CR2 when clusters are unequal, or t on $G^* - 1$), or the wild cluster bootstrap-t.* To first order this is the right variance for $\mathrm{Err}_{\mathrm{new}}$ (Proposition 8(i)). With a precisely scored target these intervals covered 0.853 to 0.915 in every design (Parts 6b and 7); normal quantiles covered 0.823 to 0.897 in Part 7.
3. *Treat the Nadeau–Bengio correction on grouped folds as conservative by a predictable factor*, $\frac{J+K-1}{J-1}\,\mathrm{Var}(a - \beta/(K-1))/\,\mathrm{Var}(a)$ relative to $\hat{V}_{\mathrm{CR}}$ (Proposition 8(ii)), predicted out of sample to within 0.053; on random-row folds it fails by the design effect (Proposition 6; N7). With unequal clusters and each fold's own mean, the factor also carries a fold-size normalization effect (Proposition 11, Section 7); fold residual totals divided by n/K remove that part.
4. *Do not use the published nested CV with random-row folds on clustered data* (N2). The cluster-nested version is unbiased in mean (Proposition 3) but too noisy at $K = 5$ (N1, N5); the floor covers (N3) but is conservative by construction.
5. *In coverage studies, redraw the clusters in every replicate and score the target precisely*, subtracting its Monte Carlo variance (Proposition 7; A2's 0.992 against 0.874 with events redrawn; Parts 6b and 7), and state whether the target is $\mathrm{Err}_{\mathrm{new}}$ or the average error.

6. *For the AUC, do not pool held-out scores across fold models when clusters differ in outcome rate.* Use BCWF-pooled with the Obuchowski SE and t_{G-1} quantiles (mean coverage 0.899 for $G \geq 40$ in Part 9, but down to 0.823 at $G = 20$), and choose the target to match the use (Section 6). For the AUC the Nadeau–Bengio correction is not conservative (ratio about 1).

9. What is new and what is known

The team's literature check (`proposals/07`) searched papers and package code, but not Google Scholar or theses.

Known.

- Proposition 6 at $J = K$ is the random group method of survey sampling [31, Chapter 2], including the rule that groups be formed from whole clusters.
- Applied with clusters as units, the CV central limit theorem of Bayle et al. [6] gives Proposition 8(i) for the average of the fold models' errors, including 0–1 loss; their Wald interval with events as units is the cluster-robust interval without $G/(G-1)$.
- Wager [30] overlaps Proposition 8(i) and (iv): CV's leading error term does not depend on the model, and the conditional risk moves at lower order.
- Random-row CV is optimistic under grouping [27, 28, 26, 24], and naive CV standard errors are too small [29, 8, 22]; the seen-against-new-cluster distinction of Lemma 1 is in [26], without intervals. We found no paper that runs the nested CV of [5] on clustered data and reports coverage, so A2 and N2 are a small confirming demonstration.
- Propositions 1 to 3 are clusters-as-units extensions of Theorem 3 of [5], of [8] and of [6], not new theory; the design effect is Kish's [16]. Grouped CV with influence curves summed within clusters [17], resampling whole subjects [10] and internal–external CV [13] are published. Proposition 7 is the classical design-based against model-based distinction.
- For the AUC, the pooling bias of pooled K-fold and leave-one-out estimates is known [25, 15, 1, 2], as are the remedies: averaging over folds, and leave-pair-out CV [1, 2, 21]. The closest work on dependent data, spatial leave-pair-out CV [3], corrects both the pooling bias and a spatial bias; BCWF is its grouped analogue.
- For unequal clusters, the ratio-estimator form of Proposition 11(b) [12], Kish's design effect with $\sum_h m_h^2 / \sum_h m_h$ [16], and the leverage bias of the CR1 variance, which CR2 removes under working independence [7, 19], are classical.

Plausibly new.

- The factor $(J + K - 1)/(J - 1)$ linking the corrected repeated K-fold variance to the random-group (cluster-robust) variance for R repeated partitions, and the reading of Proposition 6 for grouped against row folds: the corrected t handles clustering only on grouped folds (confirmed by N7). We found no statement of either.
- The Nadeau–Bengio limit $\frac{J+K-1}{J-1} \operatorname{Var}(a - \beta/(K-1))$ of Proposition 8(ii), and its quantitative check: predicted and observed ratios agreed within the bootstrap interval at all five G in Part 6b, and out of sample at $K = 3$ and 10 (N14, N15).
- The cautions for coverage studies of CV error bars: redraw the clusters (Proposition 7, A2 against N4), and score the target precisely (Parts 6b and 7).
- For the AUC, the clustered-data expansion of the pooling bias (Proposition 9(iii)): order $1/G$ rather than $1/n$, a leading term C_{mix}/G that does not depend on K with a next term $c_2/((K-1)G)$, computable constants, and its test (N16 to N21); the same-cluster-pair bias of the fold-averaged AUC; and the closed form of Proposition 10 for independent rows ($c_2 = C_{\text{mix}}/2$), which under clustering is only an approximation (the review's counterexamples). The literature check found none of these, but its Part 9 search was brief, and it asks for a further search (for example on leave-site-out AUC pooling) and a full reading of [3] before any novelty is claimed.

- For unequal clusters, the training-set weights ω_h of Proposition 11 and the statement that the Nadeau–Bengio limit survives unequal folds in expectation, the measurement of the order-$1/G$ terms at $G = 21$ (N22 to N25), and the per-data-set plug-in; these await a literature check.

We do not claim the cluster-nested, bias-adjusted interval that the literature check first proposed as the contribution: N1 failed, and N5 showed its under-coverage once events were redrawn.

10. Limitations and open questions

The dyadic case. Every person attends one event, so holding out events with event clustering covers all shared-person dependence here, and Aronow, Samii and Assenova [4] found dyadic-robust SEs barely larger than subject-clustered ones on these data. The two-way problem matters for within-event targets and for data where people appear in several clusters. The theory note proves the bias of the two-way estimator [11] for one direction of a balanced rater-by-ratee array; the two-direction case and the finite-sample bias of the dyadic-robust estimator [14, 4] are open. In the exploratory dyadic design, within-rater CV was biased by +0.024 for the full-data error (theory note), so the intervals missed that target. We claim nothing here.

Unbalanced clusters. Propositions 1, 2, 6 and 8 assume balance. Proposition 11 covers the pooled estimate, the unbalanced cluster-robust variance and the Nadeau–Bengio ratio to first order, for i.i.d. cluster sizes and the row-weighted target, and the ratio's limit only in expectation; it does not cover the cluster-weighted target. What remains open: P2 is a heuristic that omits order-$1/G$ terms of the same size as some it keeps (the pooled estimate's variation across partitions, the test cluster's gradient deviation, curvature and flip noise), and it misses at $K = 3$; under log-loss with the real sizes $\hat{V}_{\mathrm{CR}}$ ran about 17% low at $G = 21$ ($\mathrm{Var}(\widehat{\mathrm{Err}}_{\mathrm{CV}} - \mathrm{Err}_{\mathrm{new}})/\overline{\hat{V}_{\mathrm{CR}}} = 1.20$), and the second-order surrogate explains only part of the excess; and the real-data plug-in is not reliable with high-dimensional models at $G = 21$.

Unexplained parts. (a) Proposition 7 is consistent with A2's over-coverage but does not explain it quantitatively. (b) The finite-population effect at $G = 10$ does not match the factor $1 - 10/21$. (c) The $K = 2$ predictions miss by 3.4% to 6.6%. (d) Proposition 8(i) is not confirmed at the registered precision (N8, N8b). (e) $\bar{c}$ cannot be estimated from one CV run, since the cross-fold products are determined by the cluster-robust sum; a direct estimate needs variation across fold assignments. Separating $\mathrm{Var}(\beta)$ and $\mathrm{Cov}(a, \beta)$ with $K = 5$ and $K = 2$ was tried and was too noisy to use. (f) Proposition 11 gives the first-order limit for unequal events and P2 a finite-G heuristic, both checked in simulation (Part 11), but the plug-in for the real models is not reliable and partition noise at $R = 10$ is as large as the differences between models, so the real-data ratios of Section 5 remain descriptive.

The AUC. Proposition 9(iii) is a formal expansion with no remainder bound; it failed its registered out-of-sample test at $K = 2$ under strong clustering (N18), and the larger rerun that shrinks the miss to 5.8% is exploratory (`scripts/sim_cv_auc_k2check.py`, `results/sim_cv_auc_k2check.csv`). The reviewer could not exclude a small residual shortfall at large G for $K = 2$. Proposition 10's identity $c_2 = C_{\mathrm{mix}}/2$ holds for independent rows and a correct model; with clustering it is false in general, and the Lemma above is the review's (`proposals/09`). Part 9 used balanced synthetic clusters; the real events are unbalanced.

Scope. All simulations fit logistic regression; the real-data error bars cover logistic regression and LDA.

Review. The readings of Proposition 2 and Propositions 6 to 8 were reviewed adversarially once, Propositions 9 and 10 were reviewed adversarially by an automated reviewer (`proposals/09`, 16 October 2026), and Proposition 11 was reviewed adversarially by an automated reviewer on 3 November 2026 (theory note, Section 6g); the corrections of all three reviews are folded in above. None of the theory, Proposition 11 included, has yet been checked by the team's mathematical lead, Jenna Yang; that review is still to come, and until then the theory is the team's working result.

References

[1] Airola et al., JMLR W&CP **8**, 2010. [Pooling bias; leave-pair-out CV.]

[2] Airola et al., Comput. Statist. Data Anal., 2011. [Leave-pair-out CV.]
[3] Airola, Pohjankukka, Torppa, Middleton, Nykanen, Heikkonen and Pahikkala, Data Mining and Knowledge Discovery, 2019. [Spatial leave-pair-out CV.]
[4] Aronow, Samii and Assenova, *Political Analysis*, 2015.
[5] Bates, Hastie and Tibshirani, *Cross-validation: what does it estimate and how well does it do it?*, J. Amer. Statist. Assoc. **119** (2024), no. 546; arXiv:2104.00673.
[6] Bayle, Bayle, Janson and Mackey, NeurIPS, 2020. [Central limit theorem for CV.]
[7] Bell and McCaffrey, Survey Methodology **28** (2002), no. 2. [Bias-reduced linearization (CR2).]
[8] Bengio and Grandvalet, *No unbiased estimator of the variance of K-fold cross-validation*, J. Mach. Learn. Res., 2004.
[9] Bouckaert and Frank, 2004. [Corrected resampled t in repeated K-fold CV.]
[10] Cai, Luo, Guo et al., *Bootstrapping the cross-validation estimate*, arXiv:2307.00260v2.
[11] Cameron, Gelbach and Miller, *Robust inference with multiway clustering*, J. Bus. Econom. Statist., 2011.
[12] Cochran, *Sampling Techniques*, 3rd ed., Wiley, 1977. [Ratio estimators.]
[13] de Jong et al., Stat. Med., 2021. [Internal–external CV.]
[14] Fafchamps and Gubert, J. Development Economics, 2007. [Dyadic-robust variance.]
[15] Forman and Scholz, SIGKDD Explorations **12** (2010), no. 1. [Pooling bias of CV AUC.]
[16] Kish, *Survey Sampling*, Wiley, 1965.
[17] LeDell, Petersen and van der Laan, Electron. J. Statist. **9** (2015), 1583–1607.
[18] Li and Duan, 1989. [Single-index models with elliptical covariates.]
[19] MacKinnon and Webb, 2017. [Cluster-robust inference with unequal cluster sizes.]
[20] Markatou et al., 2005, Table 6. [Variance of CV estimates.]
[21] Montoya Perez et al., Statistical Methods in Medical Research, 2019. [Tournament leave-pair-out CV.]
[22] Nadeau and Bengio, Machine Learning, 2003. [Corrected resampled t.]
[23] Obuchowski, 1997. [Clustered placement-value variance of the AUC.]
[24] Park and Marcotte, Nature Methods, 2012.
[25] Parker, Gunter and Bedo, BMC Bioinformatics **8** (2007), 326. ["Stratification bias" of pooled CV AUC.]
[26] Rabinowicz and Rosset, J. Amer. Statist. Assoc. **117** (2022), no. 538.
[27] Roberts et al., *Cross-validation strategies for data with temporal, spatial, hierarchical, or phylogenetic structure*, Ecography, 2017.
[28] Saeb et al., GigaScience, 2017. [Random-row CV under grouping.]
[29] Varoquaux, NeuroImage, 2018. [Naive CV error bars.]
[30] Wager, arXiv:1909.11696, 2020. [CV error and conditional risk.]
[31] K. M. Wolter, *Introduction to Variance Estimation*, 2nd ed., Springer, New York, 2007, Chapter 2, "The Method of Random Groups", pp. 21–106.