



Search

Return to Search Results

My Tools ▾

Search History

Marked List



Save to EndNote online

Add to Marked List

1 of 2

Approximate Solution of Kuramoto-Sivashinsky Equation using Reduced Differential Transform Method

By: Acan, O (Acan, Omer)^[1]; Keskin, Y (Keskin, Yildiray)^[1]

Edited by: Simos, TE; Tsitouras, C

PROCEEDINGS OF THE INTERNATIONAL CONFERENCE OF NUMERICAL ANALYSIS AND APPLIED MATHEMATICS 2014 (ICNAAM-2014)

Book Series: AIP Conference Proceedings

Volume: 1648

Article Number: UNSP 470003

DOI: 10.1063/1.4912680

Published: 2015

Conference

Conference: International Conference on Numerical Analysis and Applied Mathematics (ICNAAM)

Location: Rhodes, GREECE

Date: SEP 22-28, 2014

Abstract

In this study, approximate solution of Kuramoto-Sivashinsky Equation, by the reduced differential transform method, are presented. We apply this method to an example. Thus, we have obtained numerical solution Kuramoto-Sivashinsky equation. Comparisons are made between the exact solution and the reduced differential transform method. The results show that this method is very effective and simple.

Keywords

Author Keywords: Reduced differential transform method (RDTM); Kuramoto-Sivashinsky(KS) Equations; Approximate solution

KeyWords Plus: VARIATIONAL ITERATION METHOD; HOMOTOPY ANALYSIS METHOD; NONLINEAR PROBLEMS

Author Information

Reprint Address: Acan, O (reprint author)

+ Selcuk Univ, Fac Sci, Dept Math, TR-42003 Konya, Turkey.

Addresses:

+ [1] Selcuk Univ, Fac Sci, Dept Math, TR-42003 Konya, Turkey

E-mail Addresses: acan_omer@selcuk.edu.tr; yildiraykeskin@yahoo.com

Publisher

AMER INST PHYSICS, 2 HUNTINGTON QUADRANGLE, STE 1N01, MELVILLE, NY 11747-4501
USA

Categories / Classification

Research Areas: Mathematics; Physics

Web of Science Categories: Mathematics, Applied; Physics, Applied

Document Information

Document Type: Proceedings Paper

Language: English

Accession Number: WOS:000355339702089

ISBN: 978-0-7354-1287-3

Citation Network

0 Times Cited

24 Cited References

View Related Records

View Citation Map

Create Citation Alert

(data from Web of Science™ Core Collection)

All Times Cited Counts

0 in All Databases

0 in Web of Science Core Collection

0 in BIOSIS Citation Index

0 in Chinese Science Citation Database

0 in Data Citation Index

0 in Russian Science Citation Index

0 in SciELO Citation Index

Usage Count

Last 180 Days: 3

Since 2013: 3

Learn more

This record is from:
Web of Science™ Core Collection

Suggest a correction

If you would like to improve the quality of the data in this record, please [suggest a correction](#).

Approximate solution of Kuramoto–Sivashinsky equation using reduced differential transform method

Omer Acan and Yildiray Keskin

Citation: [AIP Conference Proceedings](#) **1648**, 470003 (2015); doi: 10.1063/1.4912680

View online: <http://dx.doi.org/10.1063/1.4912680>

View Table of Contents: <http://scitation.aip.org/content/aip/proceeding/aipcp/1648?ver=pdfcov>

Published by the [AIP Publishing](#)

Articles you may be interested in

[Travelling wave solution of the Kuramoto-Sivashinsky equation: A computational study](#)

[AIP Conf. Proc.](#) **1479**, 777 (2012); 10.1063/1.4756252

[Hopping behavior in the Kuramoto–Sivashinsky equation](#)

[Chaos](#) **15**, 013706 (2005); 10.1063/1.1848311

[Scale and space localization in the Kuramoto–Sivashinsky equation](#)

[Chaos](#) **9**, 452 (1999); 10.1063/1.166419

[Spatiotemporal chaos in the damped Kuramoto-Sivashinsky equation](#)

[AIP Conf. Proc.](#) **342**, 702 (1995); 10.1063/1.48763

[The concept of minimal attractor and maximal attractors of partial differential equations of the Kuramoto–Sivashinsky type](#)

[Chaos](#) **1**, 168 (1991); 10.1063/1.165825

Approximate Solution of Kuramoto–Sivashinsky Equation using Reduced Differential Transform Method

Omer Acan^{1,a} and Yıldıray Keskin^{1,b}

Department of Mathematics, Science Faculty, Selcuk University, Konya 42003, Turkey.

^aCorresponding author: acan_omer@selcuk.edu.tr

^byildiraykeskin@yahoo.com

Abstract. In this study, approximate solution of Kuramoto–Sivashinsky Equation, by the reduced differential transform method, are presented. We apply this method to an example. Thus, we have obtained numerical solution Kuramoto–Sivashinsky equation. Comparisons are made between the exact solution and the reduced differential transform method. The results show that this method is very effective and simple.

Keywords: Reduced differential transform method (RDTM), Kuramoto–Sivashinsky(KS) Equations, Approximate solution.

PACS: 02.60.Cb.

INTRODUCTION

Partial differential equations (PDEs) have numerous essential applications in various fields of science and engineering such as fluid mechanic, thermodynamic, heat transfer, physics [10].

Most new nonlinear PDEs do not have a precise analytic solution. So, numerical methods have largely been used to handle these equations. It is difficult to handle nonlinear part of these equations. Although most of scientists applied numerical methods to find the solution of these equations, solving such equations analytically is of fundamental importance since the existent numerical methods which approximate the solution of PDEs don't result in such an exact and analytical solution which is obtained by analytical methods.

Many researchers have paid attention by studying to the solutions of nonlinear PDEs by various methods [11-19]. RDTM [1-4], devised by Yıldıray Keskin in 2009, is a numerical method to obtain approximate solutions of various types of nonlinear partial differential equations. It has received much attention since it has applied to solve a wide variety of problems by many authors [20-24].

In this study, RDTM is used to obtain approximate solution of KS equation. The Generalized Kuramoto–Sivashinsky equation is given [5,6] by

$$u_t + \alpha u^\beta u_x + \gamma u^\tau u_{xx} + \lambda u_{xxxx} = 0, \quad (1)$$

Where $\alpha, \beta, \gamma, \lambda, \tau \neq 0$ and $\alpha, \beta, \gamma, \lambda, \tau \in R$.

When $\alpha = \beta = 1$ and $\tau = 0$, eq. (1) reduces to original KS equation of the form [6-8]

$$u_t + uu_x + \gamma u_{xx} + \lambda u_{xxxx} = 0. \quad (2)$$

In the past several decades, many researchers have used various methods to solve KS equation. We apply the RDTM to solve KS equation of the form (2).

ANALYSIS OF THE RDTM

The basic definitions of RDTM were already given in [1].

For the purpose of illustration of the methodology to the proposed method, we write the KS equation in the standard operator form

$$L_t(u(x,t)) + L_x(u(x,t)) + N(u(x,t)) = 0 \quad (3)$$

with initial condition

$$u(x,0) = f(x) \quad (4)$$

where $L_t(u(x,t)) = u_t$ and $L_x(u(x,t)) = \gamma u_{xx} + \lambda u_{xxxx}$ are linear operators which have partial derivatives, $N(u(x,t)) = uu_x$ is a nonlinear term.

According to the RDTM, we can construct the following iteration formula:

$$(k+1)U_{k+1}(x) = -\gamma \frac{\partial^2}{\partial x^2} U_k(x) - \lambda \frac{\partial^4}{\partial x^4} U_k(x) - \sum_{r=0}^k U_{k-r}(x) \left(\frac{\partial^2}{\partial x^2} U_r(x) \right). \quad (5)$$

From initial condition (4), we write

$$U_0(x) = f(x) \quad (6)$$

substituting (6) into (5) and by a straight forward iterative calculations, we get the following $U_k(x)$ values. Then the inverse transformation of the set of values $\{U_k(x)\}_{k=0}^n$ gives approximation solution as,

$$\tilde{u}_n(x,t) = \sum_{k=0}^n U_k(x) t^k \quad (7)$$

where n is order of approximation solution.

Therefore, the exact solution of problem is given by

$$u(x,t) = \lim_{n \rightarrow \infty} \tilde{u}_n(x,t). \quad (8)$$




FIGURE 1. The surface shows the solution $u(x,t)$ for eq.(9) when $c = 0.1$, $k = (\sqrt{11/19})/4$, $x_0 = -30$ (a) exact solution (11), (b) 2th order of approximate solution for eq.(9) and (c) the absolute error between exact and numerical solution.

NUMERICAL APPLICATIONS

In this section, we test the RDTM for KS equation. We consider the following KS equation [6,9]:

$$u_t + uu_x + u_{xx} + u_{xxx} = 0 \quad (9)$$

subject to initial condition

$$u(x, 0) = c + \frac{5}{19} \sqrt{\frac{11}{19}} \left(11 \tanh^3(k(x - x_0)) - 9 \tanh(k(x - x_0)) \right). \quad (10)$$

In [6,9], the exact solution of (9) is given as

$$u(x, t) = c + \frac{5}{19} \sqrt{\frac{11}{19}} \left(11 \tanh^3(k(x - ct - x_0)) - 9 \tanh(k(x - ct - x_0)) \right). \quad (11)$$

RESULTS AND DISCUSSION

When analysed the solution of the above an example by RDTM, the following results are obtained:

TABLE 1. The exact and 2th order of approximate solution for(9) and absolute error between exact and numerical solutions

| x | t | RDTM Solution | Exact Solution | Abs-Error |
|-----|-----|---------------|----------------|-------------------------------|
| 0.0 | 0.0 | 0.5003600908 | 0.500360093 | $0.5420022964 \times 10^{-9}$ |
| 0.0 | 0.5 | 0.5003685212 | 0.500358052 | $0.1046850860 \times 10^{-4}$ |
| 0.0 | 1.0 | 0.5003762240 | 0.500355975 | $0.2024957166 \times 10^{-4}$ |
| 0.5 | 0.0 | 0.5003784827 | 0.500378483 | $0.6041276398 \times 10^{-9}$ |
| 0.5 | 0.5 | 0.5003854531 | 0.500376798 | $0.8655395022 \times 10^{-5}$ |
| 0.5 | 1.0 | 0.5003918215 | 0.500375080 | $0.1674210244 \times 10^{-4}$ |
| 1.0 | 0.0 | 0.5003936888 | 0.500393690 | $0.5954475010 \times 10^{-9}$ |
| 1.0 | 0.5 | 0.5003799452 | 0.500392296 | $0.7156111486 \times 10^{-5}$ |
| 1.0 | 1.0 | 0.5004047174 | 0.500390875 | $0.1384174699 \times 10^{-4}$ |

In this study, reduced differential transform method has been successfully applied to KSequation. This example shows that the results of the presented method are in excellent agreement with those of absolute errors and the obtained solutions shown in table.

REFERENCES

1. Y. Keskin, G. Oturanc, Reduced differential transform method for partial differential equations, *Int. J. Nonlinear Sci. Numer. Simul.* 10 (6), 2009, pp. 741–749.
2. Y. Keskin, G. Oturanc, The reduced differential transform method: a new approach to fractional partial differential equations, *Nonlinear Sci. Lett. A* 1 (2), 2010, pp. 207–217.
3. Y. Keskin, G. Oturanc, Reduced differential transform method for generalized KdV equations, *Math. Comput. Applic.* 15 (3), 2010, 382–393.
4. Y. Keskin, Ph.D Thesis, Selcuk University, 2010 (in Turkish).
5. C. Li, G. Chen and S. Zhao, Exact Travelling wave solution to the generalized Kuramoto-Sivashinsky equation, *Latin American Appl. Research*, 34, 2004, pp. 64–68.
6. M. Kurulay, A. Secer and M. A. Akinlar, A New Approximate Analytical Solution of Kuramoto – Sivashinsky Equation Using Homotopy Analysis Method, *Appl. Math. & Inf. Sci.* 7 (1), 2013, pp.267–271.
7. G. I. Sivashinsky, Nonlinear Analysis of hydrodynamic instability in laminar flames, Part I, Derivation of Basic Equations, *Acta Astronautica*, 4, 1977, pp. 1177–1206.
8. G. I. Sivashinsky, On flame propagation under conditions of stoichiometry, *SIAM J. Appl. Math.*, 39, 1980, pp. 67–82.
9. M. G. Porshokouhi and B. Ghanbari, Application of Hesvariational iteration method for solution of the family of Kuramoto-Sivashinsky equations Original Research Article, *J. King Saud Univ. Sci.*, 23 (4), 2011, pp. 407–41.
10. L. Debnath, Nonlinear Partial Differential Equations for Scientist and Engineers, Birkhauser, Boston, 1997.
11. G. Adomian, A new approach to nonlinear partial differential equations, *J. Math. Anal. Applic.* 102, 1984 pp. 420–434.
12. A.M. Wazwaz, Partial differential equations: methods and applications, The Netherlands: Balkema Publishers, 2002.

13. J.H. He, Variational iteration method-a kind of non-linear analytical technique: Some *examples Int. J. Non-Linear Mechanics*, 34 (4), 1999, pp. 699–708.
14. S.J. Liao, Homotopy analysis method: a new analytic method for nonlinear problems, *Appl. Math. And Mechanics*, 19 (10), 1998, pp. 957–962.
15. S.J. Liao, On the homotopy analysis method for nonlinear problems, *Appl. Math. Comput.* 147 (2), 2004, pp. 499–513.
16. J.K. Zhou, Differential Transformation and its Application for Electrical Circuits, Huazhong University Press, Wuhan, China, 1986.
17. N. Bildik, A. Konuralp, The use of variational iteration method, differential transform method and Adomian decomposition method for solving different types of nonlinear partial differential equations, *Int. J. Nonlinear Sci. Numer. Simul.* 7 (1), 2006, pp. 65–70.
18. V.O. Vakhnenko, E.J. Parkes, A.J. Morrison, A Bäcklund transformation and the inverse scattering transform method for the generalized Vakhnenko equation, *Chaos Solitons Fractals* 17 (4), 2003, pp. 683–692.
19. L. Brugnano, G. Carreras, K. Burrage, On the Convergence of LMF-type Methods for SODEs, *Mediterr. J. Math.* 1(3), 2004, pp. 297–313.
20. A. Saravanan, N. Magesh, A comparison between the reduced differential transform method and the Adomian decomposition method for the Newell–Whitehead–Segel equation, *J. Egypt. Math. Soc.* 21 (3), 2013, pp. 259–265.
21. R. Abazari, M. Abazari, Numerical study of Burgers–Huxley equations via reduced differential transform method, *Comput. Appl. Math.* 32 (1), 2013, pp. 1–17.
22. B. Bis, M. Bayram, Approximate solutions for some nonlinear evolutions equations by using the reduced differential transform method, *Int. J. Appl. Math. Res.* 1 (3), 2012, pp. 288–302.
23. R. Abazari, B. Soltanalizadeh, Reduced differential transform method and its application on Kawahara equations, *Thai J. Math.* 11 (1), 2013, pp. 199–216.
24. M.A. Abdou, A.A. Soliman, Numerical simulations of nonlinear evolution equations in mathematical physics, *Int. J. Nonlinear Sci.* 12 (2) 2011, pp. 131–139.