

# Forecasting: principles and practice

**Rob J Hyndman**

2.2 Transformations

## #Variance stabilization

## ##Variance stabilization

If the data show different variation at different levels of the series, then a transformation can be useful.

Denote original observations as  $y_1, \dots, y_n$  and transformed observations as  $w_1, \dots, w_n$ .

### Mathematical transformations for stabilizing variation

Square root	$w_t = \sqrt{y_t}$	↓
Cube root	$w_t = \sqrt[3]{y_t}$	Increasing
Logarithm	$w_t = \log(y_t)$	strength

Logarithms, in particular, are useful because they are more interpretable: changes in a log value are relative (percent)

## #Variance stabilization

## ##Variance stabilization

If the data show different variation at different levels of the series, then a transformation can be useful.

Denote original observations as  $y_1, \dots, y_n$  and transformed observations as  $w_1, \dots, w_n$ .

### Mathematical transformations for stabilizing variation

Square root	$w_t = \sqrt{y_t}$	↓
Cube root	$w_t = \sqrt[3]{y_t}$	Increasing
Logarithm	$w_t = \log(y_t)$	strength

Logarithms, in particular, are useful because they are more interpretable: changes in a log value are relative (percent)

#Variance stabilization

##Variance stabilization

If the data show different variation at different levels of the series, then a transformation can be useful.

Denote original observations as  $y_1, \dots, y_n$  and transformed observations as  $w_1, \dots, w_n$ .

#### Mathematical transformations for stabilizing variation

Square root	$w_t = \sqrt{y_t}$	↓
Cube root	$w_t = \sqrt[3]{y_t}$	Increasing
Logarithm	$w_t = \log(y_t)$	strength

Logarithms, in particular, are useful because they are more interpretable: changes in a log value are relative (percent)

#Variance stabilization

##Variance stabilization

If the data show different variation at different levels of the series, then a transformation can be useful.

Denote original observations as  $y_1, \dots, y_n$  and transformed observations as  $w_1, \dots, w_n$ .

#### Mathematical transformations for stabilizing variation

Square root	$w_t = \sqrt{y_t}$	↓
Cube root	$w_t = \sqrt[3]{y_t}$	Increasing
Logarithm	$w_t = \log(y_t)$	strength

Logarithms, in particular, are useful because they are more interpretable: changes in a log value are **relative (percent)**

# Box-Cox transformations

## ##Box-Cox transformations

Each of these transformations is close to a member of the family of **Box-Cox transformations**:

$$w_t = \begin{cases} \log(y_t), & \lambda = 0; \\ (y_t^\lambda - 1)/\lambda, & \lambda \neq 0. \end{cases}$$

- $\lambda = 1$ : (No substantive transformation)
- $\lambda = \frac{1}{2}$ : (Square root plus linear transformation)
- $\lambda = 0$ : (Natural logarithm)
- $\lambda = -1$ : (Inverse plus 1)

## ##Box-Cox transformations

# Box-Cox transformations

## ##Box-Cox transformations

Each of these transformations is close to a member of the family of **Box-Cox transformations**:

$$w_t = \begin{cases} \log(y_t), & \lambda = 0; \\ (y_t^\lambda - 1)/\lambda, & \lambda \neq 0. \end{cases}$$

- $\lambda = 1$ : (No substantive transformation)
- $\lambda = \frac{1}{2}$ : (Square root plus linear transformation)
- $\lambda = 0$ : (Natural logarithm)
- $\lambda = -1$ : (Inverse plus 1)

## ##Box-Cox transformations

# Box-Cox transformations

## ##Box-Cox transformations

Each of these transformations is close to a member of the family of **Box-Cox transformations**:

$$w_t = \begin{cases} \log(y_t), & \lambda = 0; \\ (y_t^\lambda - 1)/\lambda, & \lambda \neq 0. \end{cases}$$

- $\lambda = 1$ : (No substantive transformation)
- $\lambda = \frac{1}{2}$ : (Square root plus linear transformation)
- $\lambda = 0$ : (Natural logarithm)
- $\lambda = -1$ : (Inverse plus 1)

## ##Box-Cox transformations



# Box-Cox transformations

## ##Box-Cox transformations

Each of these transformations is close to a member of the family of **Box-Cox transformations**:

$$w_t = \begin{cases} \log(y_t), & \lambda = 0; \\ (y_t^\lambda - 1)/\lambda, & \lambda \neq 0. \end{cases}$$

- $\lambda = 1$ : (No substantive transformation)
- $\lambda = \frac{1}{2}$ : (Square root plus linear transformation)
- $\lambda = 0$ : (Natural logarithm)
- $\lambda = -1$ : (Inverse plus 1)

## ##Box-Cox transformations

# Box-Cox transformations

## ##Box-Cox transformations

Each of these transformations is close to a member of the family of **Box-Cox transformations**:

$$w_t = \begin{cases} \log(y_t), & \lambda = 0; \\ (y_t^\lambda - 1)/\lambda, & \lambda \neq 0. \end{cases}$$

- $\lambda = 1$ : (No substantive transformation)
- $\lambda = \frac{1}{2}$ : (Square root plus linear transformation)
- $\lambda = 0$ : (Natural logarithm)
- $\lambda = -1$ : (Inverse plus 1)

## ##Box-Cox transformations