



Forecasting: principles and practice

Rob J Hyndman

2.2 Transformations

#Variance stabilization

##Variance stabilization

If the data show different variation at different levels of the series, then a transformation can be useful.

#Variance stabilization

##Variance stabilization

If the data show different variation at different levels of the series, then a transformation can be useful.

Denote original observations as y_1, \dots, y_n and transformed observations as w_1, \dots, w_n .

#Variance stabilization

##Variance stabilization

If the data show different variation at different levels of the series, then a transformation can be useful.

Denote original observations as y_1, \dots, y_n and transformed observations as w_1, \dots, w_n .

Mathematical transformations for stabilizing variation

Square root	$w_t = \sqrt{y_t}$	\downarrow
Cube root	$w_t = \sqrt[3]{y_t}$	Increasing
Logarithm	$w_t = \log(y_t)$	strength

#Variance stabilization

##Variance stabilization

If the data show different variation at different levels of the series, then a transformation can be useful.

Denote original observations as y_1, \dots, y_n and transformed observations as w_1, \dots, w_n .

Mathematical transformations for stabilizing variation

Square root	$w_t = \sqrt{y_t}$	\downarrow
Cube root	$w_t = \sqrt[3]{y_t}$	Increasing
Logarithm	$w_t = \log(y_t)$	strength

Logarithms, in particular, are useful because they are

Box-Cox transformations

##Box-Cox transformations

Each of these transformations is close to a member of the family of **Box-Cox transformations**:

$$w_t = \begin{cases} \log(y_t), & \lambda = 0; \\ (y_t^\lambda - 1)/\lambda, & \lambda \neq 0. \end{cases}$$

Box-Cox transformations

##Box-Cox transformations

Each of these transformations is close to a member of the family of **Box-Cox transformations**:

$$w_t = \begin{cases} \log(y_t), & \lambda = 0; \\ (y_t^\lambda - 1)/\lambda, & \lambda \neq 0. \end{cases}$$

- $\lambda = 1$: (No substantive transformation)
- $\lambda = \frac{1}{2}$: (Square root plus linear transformation)
- $\lambda = 0$: (Natural logarithm)
- $\lambda = -1$: (Inverse plus 1)

##Box-Cox transformations

Box-Cox transformations

##Box-Cox transformations

Each of these transformations is close to a member of the family of **Box-Cox transformations**:

$$w_t = \begin{cases} \log(y_t), & \lambda = 0; \\ (y_t^\lambda - 1)/\lambda, & \lambda \neq 0. \end{cases}$$

- $\lambda = 1$: (No substantive transformation)
- $\lambda = \frac{1}{2}$: (Square root plus linear transformation)
- $\lambda = 0$: (Natural logarithm)
- $\lambda = -1$: (Inverse plus 1)

##Box-Cox transformations

Box-Cox transformations

##Box-Cox transformations

Each of these transformations is close to a member of the family of **Box-Cox transformations**:

$$w_t = \begin{cases} \log(y_t), & \lambda = 0; \\ (y_t^\lambda - 1)/\lambda, & \lambda \neq 0. \end{cases}$$

- $\lambda = 1$: (No substantive transformation)
- $\lambda = \frac{1}{2}$: (Square root plus linear transformation)
- $\lambda = 0$: (Natural logarithm)
- $\lambda = -1$: (Inverse plus 1)

##Box-Cox transformations

Box-Cox transformations

##Box-Cox transformations

Each of these transformations is close to a member of the family of **Box-Cox transformations**:

$$w_t = \begin{cases} \log(y_t), & \lambda = 0; \\ (y_t^\lambda - 1)/\lambda, & \lambda \neq 0. \end{cases}$$

- $\lambda = 1$: (No substantive transformation)
- $\lambda = \frac{1}{2}$: (Square root plus linear transformation)
- $\lambda = 0$: (Natural logarithm)
- $\lambda = -1$: (Inverse plus 1)

##Box-Cox transformations