



# Forecasting: principles and practice

Rob J Hyndman

1.4 Seasonality and trends

# Time series components

## ##Time series patterns

**Trend** pattern exists when there is a long-term increase or decrease in the data.

**Seasonal** pattern exists when a series is influenced by seasonal factors (e.g., the quarter of the year, the month, or day of the week).

**Cyclic** pattern exists when data exhibit rises and falls that are *not of fixed period* (duration usually of at least 2 years).

## ##Time series decomposition

# Time series components

## ##Time series patterns

**Trend** pattern exists when there is a long-term increase or decrease in the data.

**Seasonal** pattern exists when a series is influenced by seasonal factors (e.g., the quarter of the year, the month, or day of the week).

**Cyclic** pattern exists when data exhibit rises and falls that are *not of fixed period* (duration usually of at least 2 years).

## ##Time series decomposition

# Time series components

## ##Time series patterns

**Trend** pattern exists when there is a long-term increase or decrease in the data.

**Seasonal** pattern exists when a series is influenced by seasonal factors (e.g., the quarter of the year, the month, or day of the week).

**Cyclic** pattern exists when data exhibit rises and falls that are *not of fixed period* (duration usually of at least 2 years).

## ##Time series decomposition

# STL decomposition

## ##STL decomposition

- STL: “Seasonal and Trend decomposition using Loess”,
- Very versatile and robust.
- Unlike X-12-ARIMA, STL will handle any type of seasonality.
- Seasonal component allowed to change over time, and rate of change controlled by user.
- Smoothness of trend-cycle also controlled by user.
- Robust to outliers

# Forecasting and decomposition

## ##Forecasting and decomposition

- Forecast seasonal component by repeating the last year
- Forecast seasonally adjusted data using non-seasonal time series method. E.g.,
  - Holt's method — next topic
  - Random walk with drift model
- Combine forecasts of seasonal component with forecasts of seasonally adjusted data to get forecasts of original data.
- Sometimes a decomposition is useful just for understanding the data before building a