## **ASSIGNMENT - 1**

## EE24BTECH11019 - Dwarak A

## SECTION B — JEE MAIN/AIEEE

- 1) A parabola has the origin as its focus and the line x = 2 as the directrix. Then the vertex of the parabola is at (2008)
  - (a) (0,2)
  - (b) (1,0)
  - (c) (0,1)
  - (d) (2,0)
- 2) The ellipse  $x^2 + 4y^2 = 4$  is inscribed in a rectangle aligned with the coordinate axes, which in turn is inscribed in another ellipse that passes through the point (4,0). Then the equation of the ellipse is: (2009)
  - (a)  $x^2 + 12y^2 = 16$
  - (b)  $4x^2 + 48y^2 = 48$
  - (c)  $4x^2 + 64y^2 = 48$
  - (d)  $x^2 + 16y^2 = 16$
- 3) If two tangents drawn from a point **P** to the parabola  $y^2 = 4x$  are at right angles, then the locus of **P** is (2010)
  - (a) 2x + 1 = 0
  - (b) x = -1
  - (c) 2x 1 = 0
  - (d) x = 1
- 4) Equation of the ellipse whose axes are the axes of coordinates and which passes through the point (-3, 1) and has eccentricity  $\sqrt{\frac{2}{5}}$  is (2011)
  - (a)  $5x^2 + 3y^2 48 = 0$
  - (b)  $3x^2 + 5y^2 15 = 0$
  - (c)  $5x^2 + 3y^2 32 = 0$
  - (d)  $3x^2 + 5y^2 32 = 0$
- 5) Statement-1: An equation of a common tangent to the parabola  $y^2 = 16 \sqrt{3}x$  and the ellipse  $2x^2 + y^2 = 4$  is  $y = 2x + 2\sqrt{3}$

- Statement-2: If the line  $y = mx + \frac{4\sqrt{3}}{m}$ ,  $(m \ne 0)$  is a common tangent to the parabola  $y^2 = 16\sqrt{3}x$  and the ellipse  $2x^2 + y^2 = 4$ , then m satisfies  $m^4 + 2m^2 = 24$  (2012)
- (a) Statement-1 is false, Statement-2 is true.
- (b) Statement-1 is true, Statement-2 is true; Statement-2 is a correct explanation for Statement-1.
- (c) Statement-1 is true, Statement-2 is true; Statement-2 is **not** a correct explanation for Statement-1.
- (d) Statement-1 is true, Statement-2 is false.
- 6) An ellipse is drawn by taking a diameter of the circle  $(x-1)^2 + y^2 = 1$  as its semi-minor axis and a diameter of the circle  $x^2 + (y-2)^2 = 4$  is semi-major axis. If the centre of the ellipse is at the origin and its axes are the coordinate axes, then the equation of the ellipse is:

(2012)

1

- (a)  $4x^2 + y^2 = 4$
- (b)  $x^2 + 4y^2 = 8$
- (c)  $4x^2 + y^2 = 8$
- (d)  $x^2 + 4y^2 = 16$
- 7) The equation of the circle passing through the foci of the ellipse  $\frac{x^2}{16} + \frac{y^2}{9} = 1$ , and have a center at (0,3) is

(JEE M 2013)

- (a)  $x^2 + y^2 6y 7 = 0$
- (b)  $x^2 + y^2 6y + 7 = 0$
- (c)  $x^2 + y^2 6y 5 = 0$
- (d)  $x^2 + y^2 6y + 5 = 0$
- 8) Given : A circle,  $2x^2 + 2y^2 = 5$  and a parabola,  $y^2 = 4\sqrt{5}x$ .

Statement-1 : An equation of a common tangent to these curves is  $y = x + \sqrt{5}$ .

Statement-2: If the line,  $y = mx + \frac{\sqrt{5}}{m}(m \neq 0)$  is their common tangent, then m satisfies  $m^4 - 3m^2 + 2 = 0$ .

(JEE M 2016)

(JEE M 2013)

- (a) Statement-1 is true; Statement-2 is true; Statement-2 is a correct explanation for Statement-1
- (b) Statement-1 is true; Statement-2 is true; Statement-2 is **not** a correct explanation for Statement-1
- (c) Statement-1 is true; Statement-2 is false.
- (d) Statement-1 is false; Statement-2 is true.
- 9) The locus of the foot of perpendicular drawn from the centre of the ellipse  $x^2 + 3y^2 = 6$  on any tangent to it is

(JEE M 2014)

(a) 
$$(x^2 + y^2)^2 = 6x^2 + 2y^2$$
  
(b)  $(x^2 + y^2)^2 = 6x^2 - 2y^2$ 

(b) 
$$(x^2 + y^2)^2 = 6x^2 - 2y^2$$

(c) 
$$(x^2 - y^2)^2 = 6x^2 + 2y^2$$

(d) 
$$(x^2 - y^2)^2 = 6x^2 - 2y^2$$

10) The slope of the line touching both the parabolas  $y^2 = 4x$  and  $x^2 = -32y$  is

(JEE M 2014)

- (a)  $\frac{1}{8}$ (b)  $\frac{2}{3}$ (c)  $\frac{1}{2}$ (d)  $\frac{3}{2}$

- 11) Let **O** be the vertex and **Q** be any point on the parabola,  $x^2 = 8y$ . If the point **P** divides the line segment **OQ** internally in the ratio 1:3, then locus of P is: (JEE M 2015)
  - (a)  $y^2 = 2x$
  - (b)  $x^2 = 2y$
  - (c)  $x^2 = y$
  - (d)  $y^2 = x$
- 12) The normal to the curve,  $x^2 + 2xy 3y^2 = 0$ , at (1, 1)(JEE M 2015)
  - (a) meets the curve again in the third quadrant.
  - (b) meets the curve again in the fourth quadrant.
  - (c) does not meet the curve again.
  - (d) meets the curve again in the second quadrant.
- 13) The area (in sq. units) of the quadrilateral formed by the tangents at the end points of

the latera recta to the ellipse  $\frac{x^2}{9} + \frac{y^2}{5} = 1$ , is:

- (a)  $\frac{27}{2}$  (b) 27
- (c)  $\frac{27}{4}$  (d) 18
- 14) Let **P** be the point on the parabola,  $y^2 = 8x$ which is at a minimum distance from the centre C of the circle,  $x^2 + (y + 6)^2 = 1$ . Then the equation of the circle, passing through C and

having its centre at **P** is :  
(a) 
$$x^2 + y^2 - \frac{x}{4} + 2y - 24 = 0$$

(a) 
$$x^2 + y^2 - \frac{x}{4} + 2y - 24 = 0$$
  
(b)  $x^2 + y^2 - 4x + 9y + 18 = 0$ 

(c) 
$$x^2 + y^2 - 4x + 8y + 12 = 0$$

- (d)  $x^2 + y^2 x + 4y 12 = 0$
- 15) The eccentricity of the hyperbola whose length of the latus rectum is equal to 8 and the length of its conjugate axis is equal to half of the distance between its foci, is: (JEE M 2016)
  - (a)  $\frac{2}{\sqrt{3}}$
  - (b)  $\sqrt{3}$

  - (c)  $\frac{4}{3}$  (d)  $\frac{4}{\sqrt{3}}$