## TEXT CORRECTIONS for DSM-5 (10/31/13) \*Desk Reference not affected

Name of Disorder and	Currently printed as	Correct as follows:	Affected DSM-5
Type of Change			page
Sexual Masochism Disorder In "Differential Diagnosis," last word: Change "orientation" to "interest."	Differential Diagnosis Many of the conditions that could be differential diagnoses for sexual masochism disorder (e.g., transvestic fetishism, sexual sadism disorder, hypersexuality, alcohol and substance use disorders) sometimes occur also as comorbid diagnoses. Therefore, it is necessary to carefully evaluate the evidence for sexual masochism disorder, keeping the possibility of other paraphilias or other mental disorders as part of the differential diagnosis. Sexual masochism in the absence of distress (i.e., no disorder) is also included in the differential, as individuals who conduct the behaviors may be satisfied with their masochistic orientation.	Differential Diagnosis Many of the conditions that could be differential diagnoses for sexual masochism disorder (e.g., transvestic fetishism, sexual sadism disorder, hypersexuality, alcohol and substance use disorders) sometimes occur also as comorbid diagnoses. Therefore, it is necessary to carefully evaluate the evidence for sexual masochism disorder, keeping the possibility of other paraphilias or other mental disorders as part of the differential diagnosis. Sexual masochism in the absence of distress (i.e., no disorder) is also included in the differential, as individuals who conduct the behaviors may be satisfied with their masochistic interest.	695
Sexual Sadism Disorder In "Development and Course," sentence beginning "One study reported": Change "orientation" to "interest."	Development and Course Individuals with sexual sadism in forensic samples are almost exclusively male, but a representative sample of the population in Australia reported that 2.2% of men and 1.3% of women said they had been involved in bondage and discipline, "sadomasochism," or dominance and submission in the previous year. Information on the development and course of sexual sadism disorder is extremely limited. One study reported that females became aware of their sadomasochistic orientation as young adults, and another reported that the mean age at onset of sadism in a group of males was 19.4 years. Whereas sexual sadism per se is probably a lifelong characteristic, sexual sadism disorder may fluctuate according to the individual's subjective distress or his or her propensity to harm nonconsenting others. Advancing age is likely to have the same reducing effect on this disorder as it has on other paraphilic or normophilic sexual behavior.	Development and Course Individuals with sexual sadism in forensic samples are almost exclusively male, but a representative sample of the population in Australia reported that 2.2% of men and 1.3% of women said they had been involved in bondage and discipline, "sadomasochism," or dominance and submission in the previous year. Information on the development and course of sexual sadism disorder is extremely limited. One study reported that females became aware of their sadomasochistic interest as young adults, and another reported that the mean age at onset of sadism in a group of males was 19.4 years. Whereas sexual sadism per se is probably a lifelong characteristic, sexual sadism disorder may fluctuate according to the individual's subjective distress or his or her propensity to harm nonconsenting others. Advancing age is likely to have the same reducing effect on this disorder as it has on other paraphilic or normophilic sexual behavior.	697
Pedophilic Disorder In "Diagnostic Features," first paragraph, last sentence: Change "orientation" to "interest."	Diagnostic Features The diagnostic criteria for pedophilic disorder are intended to apply both to individuals who freely disclose this paraphilia and to individuals who deny any sexual attraction to prepubertal children (generally age 13 years or younger), despite substantial objective evidence to the contrary. Examples of disclosing this	Diagnostic Features The diagnostic criteria for pedophilic disorder are intended to apply both to individuals who freely disclose this paraphilia and to individuals who deny any sexual attraction to prepubertal children (generally age 13 years or younger), despite substantial objective evidence to the contrary. Examples of disclosing this	698

	paraphilia include candidly acknowledging an intense sexual interest in children and indicating that sexual interest in children is greater than or equal to sexual interest in physically mature individuals. If individuals also complain that their sexual attractions or preferences for children are causing psychosocial difficulties, they may be diagnosed with pedophilic disorder. However, if they report an absence of feelings of guilt, shame, or anxiety about these impulses and are not functionally limited by their paraphilic impulses (according to self-report, objective assessment, or both), and their self-reported and legally recorded histories indicate that they have never acted on their impulses, then these individuals have a pedophilic sexual orientation but not pedophilic disorder.	paraphilia include candidly acknowledging an intense sexual interest in children and indicating that sexual interest in children is greater than or equal to sexual interest in physically mature individuals. If individuals also complain that their sexual attractions or preferences for children are causing psychosocial difficulties, they may be diagnosed with pedophilic disorder. However, if they report an absence of feelings of guilt, shame, or anxiety about these impulses and are not functionally limited by their paraphilic impulses (according to self-report, objective assessment, or both), and their self-reported and legally recorded histories indicate that they have never acted on their impulses, then these individuals have a pedophilic sexual interest but not pedophilic disorder.	
Pedophilic Disorder In "Risk and Prognostic Factors," "Genetic and physiological," last word: Change "orientation" to "interest."	Genetic and physiological. Since pedophilia is a necessary condition for pedophilic disorder, any factor that increases the probability of pedophilia also increases the risk of pedophilic disorder. There is some evidence that neurodevelopmental perturbation in utero increases the probability of development of a pedophilic orientation.	Genetic and physiological. Since pedophilia is a necessary condition for pedophilic disorder, any factor that increases the probability of pedophilia also increases the risk of pedophilic disorder. There is some evidence that neurodevelopmental perturbation in utero increases the probability of development of a pedophilic interest.	699