

# Association Between Brain Structure and Alcohol Use Behaviors in Adults A Mendelian Randomization and Multiomics Study

Lucas A. Mavromatis, ScB; Daniel B. Rosoff, AB, ScB; Renata B. Cupertino, PhD; Hugh Garavan, PhD; Scott Mackey, PhD; Falk W. Lohoff, MD

[+ Supplemental content](#)

**IMPORTANCE** Past studies have identified associations between brain macrostructure and alcohol use behaviors. However, identifying directional associations between these phenotypes is difficult due to the limitations of observational studies.

**OBJECTIVE** To use mendelian randomization (MR) to identify directional associations between brain structure and alcohol use and elucidate the transcriptomic and cellular underpinnings of identified associations.

**DESIGN, SETTING, AND PARTICIPANTS** The main source data comprised summary statistics from population-based and case-control genome-wide association studies (GWAS) of neuroimaging, behavioral, and clinical phenotypes (N = 763 874). Using these data, bidirectional and multivariable MR was performed analyzing associations between brain macrostructure and alcohol use. Downstream transcriptome-wide association studies (TWAS) and cell-type enrichment analyses investigated the biology underlying identified associations. The study approach was data driven and did not test any a priori hypotheses. Data were analyzed August 2021 to May 2022.

**MAIN OUTCOMES AND MEASURES** Brain structure phenotypes (global cortical thickness [GCT] and global cortical surface area [GCSA] in 33 709 individuals and left-right subcortical volumes in 19 629 individuals) and alcohol use behaviors (alcoholic drinks per week [DPW] in 537 349 individuals, binge drinking frequency in 143 685 individuals, and alcohol use disorder in 8845 individuals vs 20 657 control individuals [total of 29 502]).

**RESULTS** The main bidirectional MR analyses were performed in samples totaling 763 874 individuals, among whom more than 94% were of European ancestry, 52% to 54% were female, and the mean cohort ages were 40 to 63 years. Negative associations were identified between genetically predicted GCT and binge drinking ( $\beta$ , -2.52; 95% CI, -4.13 to -0.91) and DPW ( $\beta$ , -0.88; 95% CI, -1.37 to -0.40) at a false discovery rate (FDR) of 0.05. These associations remained significant in multivariable MR models that accounted for neuropsychiatric phenotypes, substance use, trauma, and neurodegeneration. TWAS of GCT and alcohol use behaviors identified 5 genes at the 17q21.31 locus oppositely associated with GCT and binge drinking or DPW (FDR = 0.05). Cell-type enrichment analyses implicated glutamatergic cortical neurons in alcohol use behaviors.

**CONCLUSIONS AND RELEVANCE** The findings in this study show that the associations between GCT and alcohol use may reflect a predispositional influence of GCT and that 17q21.31 genes and glutamatergic cortical neurons may play a role in this association. While replication studies are needed, these findings should enhance the understanding of associations between brain structure and alcohol use.

**Author Affiliations:** Section on Clinical Genomics and Experimental Therapeutics, National Institute on Alcohol Abuse and Alcoholism, National Institutes of Health, Bethesda, Maryland (Mavromatis, Rosoff, Lohoff); National Institutes of Health–Oxford–Cambridge Scholars Program; Nuffield Department of Population Health, University of Oxford, Oxford, United Kingdom (Rosoff); Department of Psychiatry, University of Vermont College of Medicine, Burlington (Cupertino, Garavan, Mackey).

**Corresponding Author:** Falk W. Lohoff, MD, Section on Clinical Genomics and Experimental Therapeutics, National Institute on Alcohol Abuse and Alcoholism, National Institutes of Health, 10 Center Dr (10CRC/2-2352), Bethesda, MD 20892-1540 (falk.lohoff@nih.gov).

JAMA Psychiatry. doi:10.1001/jamapsychiatry.2022.2196  
Published online August 10, 2022.

Alcohol misuse causes large health and economic burdens globally<sup>1,2</sup> and is a leading risk factor for premature death and disability in individuals aged 15 to 49 years.<sup>1</sup> Heavy alcohol consumption impairs the nervous system,<sup>3-6</sup> which may lead to neurological, cognitive, and psychiatric health ramifications.<sup>7</sup> Altered macroscale brain structure is associated with psychopathology<sup>8</sup> and could represent a mechanistic link between alcohol-associated neurotoxicity and health outcomes. Studies have consistently associated greater alcohol use and alcohol misuse with lower cortical and subcortical volumes.<sup>9-16</sup> However, the directionality of these associations remains unclear, with some studies suggesting a predispositional impact of brain anatomy on alcohol use,<sup>9,11,13,17</sup> challenging the notion that brain structure changes as a result of alcohol exposure.<sup>14,18-21</sup>

Robert et al<sup>13</sup> analyzed longitudinal adolescent brain imaging data among 726 individuals and concluded that a greater rate of gray matter atrophy in frontal and temporal regions may lead to greater frequency of drunkenness. Similarly, a 2021 latent causal variable analysis<sup>11</sup> suggested that greater pars opercularis volume, greater cuneus thickness, and lower basal forebrain volume were associated with increased alcohol misuse. By contrast, other studies continue to suggest that alcohol use alters neuroanatomy. For example, a 2021 co-twin study<sup>18</sup> among 436 individuals found that alcohol exposure and genetic predisposition to alcohol use decreased thickness in multiple cortical regions. These studies highlight the ongoing debate regarding the directionality of associations between brain structure and alcohol use. Randomized clinical trials conducted to infer causality cannot be ethically or practically applied to study these associations and tens of thousands of participants may be required to identify replicable associations between brain magnetic resonance imaging (MRI) measures and behavioral traits.<sup>22</sup> Alternative approaches using large data sets are needed to characterize associations between brain structure and alcohol use.

Recently developed genomics methods, including latent causal variable analysis and mendelian randomization (MR), facilitate the identification of directional associations between genetically influenced variables from population-based observational data and have been underapplied to questions regarding alcohol use and brain structure.<sup>18,23-25</sup> Latent causal variable analysis only evaluates 2 phenotypes<sup>26</sup> and does not explicitly test bidirectional associations.<sup>27</sup> By contrast, MR is frequently used to evaluate directionality in neuropsychiatry.<sup>28</sup> The multivariable extension of MR (MVMR) enables the assessment of multiple exposures to identify the direct association of each exposure with an outcome,<sup>29</sup> which could help clarify the associations between brain structure and alcohol consumption accounting for potential mediating or confounding phenotypes.

In this study, we investigated associations between brain anatomy and alcohol use using summary-level genome-wide association study (GWAS) data for brain MRI measures and alcohol-related phenotypes. Our primary MR associated genetically predicted global cortical thickness (GCT) with alcohol use. We investigated whether GCT broadly associates with substance use by evaluating its association with smoking. Given

## Key Points

**Question** Are there directional associations between cortical or subcortical macrostructure and alcohol use?

**Findings** This mendelian randomization study including 763 874 participants in UK Biobank, Enhancing Neuroimaging Genetics through Meta Analysis (ENIGMA), Psychiatric Genomics Consortium (PGC), and GWAS & Sequencing Consortium of Alcohol and Nicotine use (GSCAN) studies identified a significant negative association between genetically predicted global cortical thickness and alcohol consumption and binge drinking. Downstream multiomic analyses indicate that 17q21.31 genes and glutamatergic cortical neurons contribute to this association.

**Meaning** The results from this study support emerging literature suggesting that cortical structure is associated with alcohol use and identify transcriptomic and cellular associations between these phenotypes that warrant further investigation.

differences in alcohol use patterns between men and women,<sup>30</sup> we examined sex-specific associations between GCT and alcohol use. Next, we used MVMR accounting for confounding or mediating phenotypes to test the robustness of our GCT findings and performed multiomic analyses, including transcriptomic imputation<sup>31</sup> and cell-type enrichment analysis,<sup>32</sup> to describe the biological underpinnings of GCT-alcohol use associations.

## Methods

Figure 1 presents a study overview. This study is reported in accordance with the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) reporting guideline (eTable 1 in Supplement 1).<sup>33</sup> This study uses deidentified publicly available data, so no ethical approval from an institutional review board was required. The study protocol was not preregistered.


## Data Sources

Summary-level data were obtained from GWAS. Included GWAS have existing ethical permissions from their respective institutional review boards and include participant informed consent with rigorous quality control. Participants with missing phenotypic data were excluded from source GWAS. Exact data for sex and ethnicity were not available from all sources and are reported here as approximate percentages.

## Cortical and Subcortical Structure

Our primary analyses evaluated 19 measures of cortical and subcortical brain structure sampled with MRI. We analyzed measures of global cortical thickness (GCT) and global cortical surface area (GCSA) from a recent GWAS of T1-weighted MR images from 1.5-3 T scans ( $n = 33\,709$ ).<sup>34</sup> We performed secondary analyses on 34 regional cortical measures (eMethods in Supplement 1), but emphasized global averages rather than regional phenotypes because global measures may be less impacted by interindividual neuroanatomical variability,<sup>35</sup> the

Figure 1. Study Overview



GWAS indicates genome-wide association study; SNV, single-nucleotide variant.

limited functional relevance of gyral-based atlases like the Desikan-Killiany atlas used by Grasby et al,<sup>34,36</sup> and the multiple-testing burden associated with a hypothesis-free regional analysis. GCSA was measured at the gray-white matter boundary and GCT was defined as the average distance between white matter and pial surfaces across both cortical hemispheres.<sup>34</sup> Images were processed using FreeSurfer.<sup>34,37</sup>

Additionally, because a previous mega-analysis identified left-right hemispheric asymmetry in associations between subcortical structure and alcohol use,<sup>12</sup> we focused our subcortical analysis on left-right volumes ( $n = 19\,629$ ).<sup>38</sup> We evaluated 17 left-right volumes derived from T1-weighted MR images from 1.5-3 T scans: amygdala, hippocampus, accumbens, putamen, pallidum, thalamus, insula, caudate, and brainstem (combined volume). Mindboggle 101 atlases were used to label subcortical structures.<sup>38</sup> MRI data were processed using advanced normalization tools.<sup>38</sup> In exploratory bidirectional analyses, we analyzed phenotypes from a GWAS of overall subcortical volumes.<sup>39</sup> We also investigated the association of genetically predicted alcohol consumption with longitudinal changes in brain structure (eMethods in Supplement 1).

#### Alcohol Use Behaviors

We used 3 GWAS of alcohol use behaviors in predominantly EA samples: a meta-analysis GWAS<sup>40</sup> of alcoholic drinks consumed per week (DPW) ( $n = 537\,349$ ), a GWAS<sup>41</sup> of binge drinking frequency among participants in the UK Biobank ( $n = 143\,685$ ), and a case-control GWAS<sup>42</sup> of alcohol use disorder

(AUD) (8845 individuals with AUD and 20 657 control individuals; total  $n = 29\,502$ ). Further description of these studies can be found in the eMethods in Supplement 1.

#### MR

##### MR Instruments

The eMethods in Supplement 1 provides detailed methodology for instrument clumping, evaluations of instrument strength, sample overlap, procedures for missing instrument data, and testing of MR assumptions (eFigure 1 in Supplement 1). Our DPW and global cortical structure instruments included all genome-wide significant (GWS) single-nucleotide variants (SNVs) at a threshold of  $P < 5 \times 10^{-8}$ . Like previous neuropsychiatric MR studies evaluating exposures with few GWS SNVs,<sup>43,44</sup> we used a  $P$  value threshold of  $5 \times 10^{-6}$  to select AUD, binge drinking, subcortical structure, and regional thickness instruments (eTable 3 in Supplement 1 and eTables 4-6 in Supplement 2). We performed MR using SNVs within the *ADH1B* (alcohol dehydrogenase 1B) locus, a primary enzyme in alcohol metabolism,<sup>45</sup> as sensitivity analyses further assessing relationships of alcohol use on brain structure (eMethods in Supplement 1).

We evaluated our main GCT findings with additional MR. First, we tested the association of 34 regional cortical thickness exposures<sup>34</sup> with alcohol use. Next, we evaluated the associations of GCT and smoking behaviors and examined potential sex-specific associations of GCT and alcohol use using sex-specific alcohol use GWAS data from the UK Biobank.

Finally, we performed 11 MVMR analyses (eFigures 2 and 3 in [Supplement 1](#)) accounting for neuropsychiatric phenotypes, substance use, trauma, and neurodegeneration. We concatenated, extracted, and harmonized the independent instrument sets for GCT and the controlled-for exposure with each alcohol use behavior using standard MVMR methods (eMethods in [Supplement 1](#) and eTables 7-17 in [Supplement 2](#)).<sup>46</sup> We also performed leave-1-out MR<sup>47</sup> and investigated the biological function of the GCT instrument with a gene-set enrichment analysis (eMethods in [Supplement 1](#)).

#### MR Statistical Analysis

We used the conventional inverse-variance weighted estimator (IVW) as our primary MR method. We supplemented IVW MR with MR-Egger, weighted median, weighted mode, and simple mode estimators, which rely on different assumptions than IVW.<sup>48-50</sup> Evaluating multiple estimators facilitates the assessment of the robustness of MR estimates and is important in MR studies evaluating complex traits (ie, GCT) where the biological function of the instruments is unknown.<sup>51</sup> Additionally, we used the Cochran Q heterogeneity test to evaluate heterogeneity in instrument effects, as heterogeneity may indicate violations of IVW assumptions.<sup>52</sup> We used MR-PRESSO<sup>53</sup> and MR-Lasso<sup>54</sup> to obtain MR estimates with heterogenous SNVs removed. Finally, we used the MR Steiger directionality test for reverse causality in our bidirectional MR.<sup>55</sup>

We conservatively defined significance at a false discovery rate (FDR) of 0.05 for each MR analysis (eMethods in [Supplement 1](#)). We also discuss nominally significant results ( $P < .05$ ). We report MR estimates as  $\beta$  values representing a change in outcome units per change in exposure unit. The unit for DPW was log-transformed,<sup>40</sup> AUD was a binary measure, and binge drinking frequency was a categorical measure quantified as (0) never (1) less than monthly (2) monthly (3) weekly (4) daily/almost daily. Brain structure phenotypes were quantified as cortical thickness (mm); cortical surface area (mm<sup>2</sup>), and subcortical volumes (cm<sup>3</sup>) (eMethods in [Supplement 1](#)).

#### Transcriptome-Wide Association Studies (TWAS)

We used the FUSION method<sup>31</sup> to identify gene transcript-level associations with the alcohol use and brain structure phenotypes. To perform TWAS, we integrated alcohol use and GCT GWAS summary statistics with cortical RNA sequence reference panels from the CommonMind Consortium<sup>56</sup> and the Genotype-Tissue Expression Consortium<sup>57</sup> (eMethods in [Supplement 1](#)).

#### Cell-Type Enrichment Analyses

We used Cell-Type Expression-Specific Integration for Complex Traits<sup>32</sup> with default parameters to perform cell-type enrichment analyses using the alcohol-associated GWAS data and single-cell RNA sequencing data of 120 cortical cell types (56 excitatory neurons, 54 inhibitory neurons, and 10 nonneuronal cells) from the Allen Brain Map Human Multiple Cortical Areas SMART-sequence data set<sup>58</sup> (eMethods in [Supplement 1](#)).

## Results

### Bidirectional MR Reveals Negative Association Between Genetically Predicted Global Cortical Thickness and Alcohol Use Behaviors

The main bidirectional MR analyses included 763 874 individuals who were predominantly of European ancestry (more than 94%). Cohorts had mean ages between 40 and 63 years, and 52% to 54% of included individuals were female (eTable 2 in [Supplement 1](#)). Analyses revealed significant associations of GCT with alcohol use at FDR = 0.05. These associations were unidirectional. The MR analyses failed to find any nominally significant associations between genetically predicted alcohol use and GCT (eTables 18 and 19 in [Supplement 2](#)). Conversely, we found that genetically predicted GCT has a negative association with DPW and binge drinking frequency (DPW  $\beta$ , -0.88; 95% CI -1.36 to -0.40;  $P = 3.58 \times 10^{-4}$ ; binge drinking  $\beta$ , -2.52, CI; -4.13 to -0.91;  $P = .002$ ) ([Figure 2](#); IVW estimators are presented unless otherwise specified). The associations between GCT, DPW, and binge drinking remained significant using weighted median and MR-Lasso estimators, supporting the validity of the IVW estimate ([Table 1](#)). Regarding associations between global cortical surface area (GCSA) and alcohol phenotypes, one finding suggested genetically predisposed GCSA was positively associated with DPW ( $\beta$ ,  $3.87 \times 10^{-6}$ ; 95% CI,  $1.16 \times 10^{-6}$  to  $6.59 \times 10^{-6}$ ;  $P = .005$ ); however, other MR methods did not corroborate this association. Additionally, unlike GCT, GCSA was not associated with binge drinking.

Our tests of left-right subcortical volumes on alcohol behaviors yielded 3 significant results using secondary MR methods (FDR = 0.05) (eTable 20 in [Supplement 2](#)). Greater right pallidum volume was unidirectionally associated with increased binge drinking (IVW-PRESSO  $\beta$ , 0.063; 95% CI, 0.023 to 0.10;  $P = .002$ ; LASSO  $\beta$ , 0.06; 95% CI, 0.03 to 0.09;  $P = 3.69 \times 10^{-4}$ ) and greater risk for AUD (LASSO  $\beta$ , 0.25; 95% CI, 0.09 to 0.40;  $P = .002$ ) ([Figure 2](#)). Genetically predicted right pallidum volume was nominally associated with binge drinking and AUD (binge  $\beta$ , 0.08; 95% CI, 0.030 to 0.13;  $P = .001$ ; AUD  $\beta$ , 0.18; 95% CI, 0.017 to 0.35;  $P = .03$ ). The other nominally significant unidirectional associations between subcortical volumes and alcohol use were left pallidum with binge drinking ( $\beta$ , 0.06; 95% CI, 0.01 to 0.10;  $P = .01$ ), brain stem with binge drinking ( $\beta$ , -0.04, 95% CI, -0.07 to -0.00;  $P = .03$ ), and brain stem with AUD ( $\beta$ , -0.17; 95% CI, -0.33 to -.00;  $P = .047$ ). The right pallidum was the subcortical structure most strongly associated with alcohol use in this study. The genetically predicted volume of this region was positively associated with alcohol use across methodologies and alcohol use behaviors. Beyond the right pallidum, we failed to robustly connect subcortical volumes with alcohol use. Additionally, unlike our GCT finding, our pallidum finding was not corroborated by weighted median analysis or a significant association with DPW.

MR identified nominally significant unidirectional associations between binge drinking and right amygdala volume ( $\beta$ , -0.19; 95% CI, -0.35 to -0.04;  $P = .01$ ) and between AUD

Figure 2. Bidirectional Mendelian Randomization of Brain Structures and Alcohol Use Behaviors




Table 1. Mendelian Randomization (MR) of Global Cortical Thickness (GCT), Alcoholic Drinks Per Week (DPW), and Binge Drinking Frequency Using Complementary MR Estimators

Exposure	Outcome	MR estimator	No. of instrument SNVs	Effect estimate, $\beta$ (95% CI)	P value
GCT	DPW	IVW	5	-0.88 (-1.36 to -0.40)	$3.58 \times 10^{-4}$
		MR Egger	5	-2.26 (-4.12 to -0.40)	.098
		MR Lasso	5	-0.88 (-1.36 to -0.40)	$3.58 \times 10^{-4}$
		Weighted median	5	-0.80 (-1.33 to -0.28)	.003
		Simple mode	5	-0.77 (-1.51 to -0.03)	.11
		Weighted mode	5	-0.78 (-1.54 to -0.03)	.11
	Binge drinking frequency	IVW	6	-2.52 (-4.13 to -0.91)	.002
		Post-PRESSO IVW	5	-1.64 (-2.66 to -0.63)	.002
		MR Egger	6	-6.91 (-14.01 to 0.19)	.13
		MR Lasso	5	-1.64 (-2.66 to -0.63)	.002
		Weighted median	6	-2.13 (-3.36 to -0.90)	$6.75 \times 10^{-4}$
		Simple mode	6	-1.83 (-3.82 to 0.17)	.13
		Weighted mode	6	-1.57 (-3.31 to 0.17)	.14

Abbreviations: IVW, inverse-variance weighted estimator; SNV, single-nucleotide variant.

and right putamen volume ( $\beta$ , -0.04; 95% CI, 0.00 to 0.08;  $P = .04$ ) (Figure 2; eTable 21 in Supplement 2). However, exploratory bidirectional results using overall subcortical volumes were null (eTables 22 and 23 in Supplement 2). MR estimates using *cis-ADH1B* instruments were also null for both cortical and subcortical structures (eTables 24-26 in Supplement 2), as were exploratory estimates of the associations of alcohol use with age-independent and age-dependent longitudinal changes in brain structure (eTables 27-30 in Supplement 2). Ultimately, our most robust finding was an association between genetically predicted GCT and alcohol use, motivating the focus of our downstream analyses.

### MR Testing the Robustness of GCT-Alcohol Consumption Associations

To investigate whether specific cortical regions underlie the association between GCT and alcohol use, we performed MR using regional thickness phenotypes as exposures and DPW, binge drinking, and AUD as outcomes. No results approached FDR significance (eResults in Supplement 1 and eTables 31, 32, and 33 in Supplement 2). Additionally, we failed to find an association between genetically predicted GCT and smoking, and exploratory MR identified no sex differences in GCT-alcohol consumption associations (eResults in Supplement 1 and eTables 34 and 35 in Supplement 2). Leave-1-out analyses found



**Table 2. FUSION<sup>31</sup> Transcriptome-Wide Association Studies (TWAS): Protein Coding Genes Oppositely Associated With Global Cortical Thickness (GCT) and Alcohol Consumption**

Gene	Locus	Z score: thickness	Reference panel: thickness	Significant alcohol use associations	Z score(s): alcohol use	Top reference panel(s): alcohol use
<i>PLEKHM1</i>	17q21.31	5.59	GTEx cortex	Binge drinking, DPW	-5.77, -5.34	GTEx cortex, GTEx cortex
<i>CRHR1</i>	17q21.31	-5.16	CMC DLPFC: splicing	Binge drinking, DPW	5.09, 4.75	CMC DLPFC: splicing, CMC DLPFC: splicing
<i>ARHGAP27</i>	17q21.31	4.89	CMC DLPFC	Binge drinking, DPW	-4.87, -4.50	CMC DLPFC, CMC DLPFC
<i>LRRC37A2</i>	17q21.31	5.30	GTEx cortex	Binge drinking	-6.13	GTEx cortex
<i>LRRC37A</i>	17q21.31	3.75	GTEx cortex	Binge drinking	-4.21	GTEx cortex
<i>ACTR1B</i>	2q11.2	6.11	CMC DLPFC: splicing	DPW	3.06	CMC DLPFC: splicing
<i>WNT3</i>	17q21.31	4.62	GTEx cortex	DPW	3.11	GTEx cortex
<i>RTN1</i>	14q23.1	4.15	CMC DLPFC: splicing	DPW	2.97	CMC DLPFC: splicing

Abbreviations: CMC, CommonMind Consortium; DLPFC, dorsolateral prefrontal cortex; DPW, alcoholic drinks per week; GTEx, Genotype-Tissue Expression Consortium.

no evidence of high influence variants among GCT instrument SNVs (eFigure 4 and eTable 36 in [Supplement 1](#)). In 11 MVMR models jointly assessing GCT and possible confounding exposures on alcohol use, GCT retained robust, statistically significant associations with DPW and binge drinking (eMethods in [Supplement 1](#) and eTable 37 in [Supplement 2](#)). The GCT MVMR estimates were similar in magnitude and direction to the corresponding single-variable estimates except in the MVMR model accounting for cognition. In this model, the GCT estimate on DPW was reduced by 29.5% (from  $\beta$ , -0.88 to  $\beta$ , -0.62) relative to the single-variable estimate, while the estimate on binge drinking was reduced by 46.4% (from  $\beta$ , -2.52 to  $\beta$ , -1.35). MVMR-IVW estimates were broadly consistent with MVMR-Egger sensitivity analyses; however, the MVMR-Egger estimates were substantially less precise. MVMR-Egger intercept analysis did not indicate horizontal pleiotropy (eTable 37 in [Supplement 2](#)).

### Genes at the 17q21.31 Locus Were Oppositely Associated With GCT and Alcohol Use

We performed TWAS on GCT and alcohol consumption using 3 cortical transcriptomic expression and splicing reference panels<sup>56,57</sup> (Table 2). eTables 38-41 in [Supplement 2](#) contain the full TWAS results. We identified 8 protein-coding genes associated with both GCT and 1 or more alcohol use behaviors: *ACTR1B*, *PLEKHM1*, *LRRC37A2*, *CRHR1*, *ARHGAP27*, *WNT3*, *RTN1*, and *LRRC37A*. Five of these genes, *PLEKHM1*, *LRRC37A2*, *CRHR1*, *ARHGAP27*, and *LRRC37A*, were oppositely associated with GCT and alcohol consumption (eTable 42 in [Supplement 1](#)). All 5 of these genes are contained within the 17q21.13 locus.

### Cell-Type Enrichment Analysis—Associated Glutamatergic Cortical Neurons With Alcohol Consumption

eTable 43 in [Supplement 2](#) contains full results from our cell-type enrichment analysis. We found a total of 31 nominally significant associations between a cell type and a alcohol use behavior representing 30 distinct cell types, 27 of which are excitatory glutamatergic cells and 3 of which are inhibitory GABAergic cells. Twelve excitatory cell types remained significant at FDR = 0.05, including 10 cells associated with DPW and 2 associated with binge drinking (Figure 3).


## Discussion

This MR study used large population-based data on the genetic architecture of cortical and subcortical structure,<sup>34,38</sup> MR, and novel multiomic methods to identify directional and biological associations between human brain structure and alcohol use. Our large sample sizes (between 19 629 and 537 349 participants<sup>34,38,40-42</sup>) increased statistical power relative to previous brain structure-alcohol consumption studies.<sup>13,18</sup> Our findings suggest that a predisposition toward lower GCT may be associated with greater alcohol consumption and binge drinking. Conversely, we failed to find strong evidence that a genetic predisposition for alcohol use was associated with brain structure or its longitudinal plasticity.

More modestly, our study suggests genetically predicted right pallidum volume was positively associated with alcohol consumption. This finding was not replicated in either our MR of overall subcortical volumes or a recent MR by Logtenberg et al<sup>59</sup> investigating substance use and overall subcortical volumes. Additionally, while Logtenberg et al<sup>59</sup> associated alcohol dependence with reduced overall amygdala and hippocampal volumes, after multiple testing corrections, we failed to associate a genetic liability for binge drinking, DPW, AUD, or an *ADH1B* instrument with these regions using hemispheric, overall, and longitudinal subcortical outcomes. Discrepancies between our studies may have resulted from differences in statistical methodology, power, or the specific phenotypes evaluated (eDiscussion in [Supplement 1](#)).

Our consistent identification of an association between genetically predicted GCT and alcohol use behaviors across MR methods and sex-specific analyses implicates the cortex as a potential driver of vulnerability to alcohol consumption and binge drinking. Interestingly, GCT had no association with smoking, suggesting its association with alcohol consumption may not reflect a broader association with substance use. Our failure to identify specific cortical regions associated with alcohol use may mean larger data sets are needed to characterize such associations. Importantly, GCT estimates from MVMR analyses remained significant when accounting for 11 potential mediators or confounders. The reduction in GCT effect estimates in MVMR models accounting for cognition

Figure 3. Cell-Type Expression-Specific Integration for Complex Traits–Cell-Type Enrichment Analyses



suggests mediation of the GCT-alcohol use associations, especially given the significant MVMR estimates for cognition on alcohol use (eTable 37 in Supplement 2). Additionally, while we failed to find evidence for alcohol-associated cortical thinning in a population of adults with a mean (SD) age of 40 (8) years, alcohol use could cause cortical thinning in younger adults and adolescents due to increased cortical plasticity during these developmental stages.<sup>19</sup> For instance, recent work analyzing young adults showed that alcohol use predisposition leads to decreased thickness of cortical control and salience networks.<sup>18</sup> While participant age and other methodological particularities may influence the results of studies investigating alcohol-brain structure interactions, we found that for a middle-aged population, alcohol use primarily followed cortical anatomy.

Our investigation of the transcriptomic relationship between GCT and alcohol use identified 5 protein coding genes oppositely associated with GCT and alcohol use behavior: *PLEKHM1*, *LRRC37A2*, *CRHR1*, *ARHGAP27*, and *LRRC37A*. These 5 genes could contribute to the negative association between GCT and alcohol use. All 5 are located at 17q21.31. This locus, characterized by extensive linkage disequilibrium,<sup>60</sup> is the site of 2 haplotypes: the inverted H2 haplotype (found in approximately 20% of individuals of European ancestry), and the H1 haplotype.<sup>61</sup> Comparing our imputed transcriptomes with limited cortical RNA-sequence data and past association studies suggests that lower GCT and greater alcohol use may be associated with the H1

haplotype.<sup>62</sup> Notably, *CRHR1* encodes a G-protein coupled receptor that binds corticotropin-releasing hormone and its agonists. In line with our findings, *CRHR1* upregulation in the amygdala<sup>63,64</sup> and cortex<sup>63</sup> have been associated with greater alcohol consumption and dependence. Additionally, previous studies have suggested that *CRHR1* modulates the behavioral and cognitive outcomes associated with stress.<sup>65,66</sup> *CRHR1* may also affect cortical macrostructure, as previous studies indicate *CRHR1* overexpression may be associated with early life stress-induced neuroanatomical changes and dendritic spine loss,<sup>67,68</sup> suggesting a potential mechanism whereby early life stress interacts with *CRHR1* to impact cortical structure, leading to behavioral adaptations and harmful alcohol use. We present this hypothesis cautiously due to *CRHR1*'s location in a linkage disequilibrium block containing genes like *MAPT*, which may be involved in neurodegenerative diseases and cortical anatomy.<sup>69</sup>

Our cell-type analysis also found that excitatory neurons may underlie GCT's association with alcohol use. These data support the notion that glutamatergic transmission plays an important role in alcohol misuse.<sup>70,71</sup> Interestingly, *CRHR1* is expressed in glutamatergic, but not GABAergic, cortical neurons.<sup>72</sup> Activation of *CRHR1* in the forebrain is associated with alteration in glutamatergic neurotransmission and increased behavioral susceptibility to stress in mice.<sup>72</sup> Therefore, our single-cell findings support our hypothesis associating cortical *CRHR1* expression with increased stress susceptibility, cortical thinning, and alcohol misuse.

## Limitations

This study has several limitations. First, MR instrumentation in neuropsychiatry remains challenging due to the complexity of the phenotypes and frequent uncertainty of genetic variants' biological functions.<sup>28</sup> For several of our alcohol use phenotypes, we used a relaxed *P* value threshold due to the limited number of variants at GWS, in line with previous psychiatric MR studies.<sup>43,44</sup> While these relaxed thresholds could introduce weak instrument bias or increase the possibility of horizontal pleiotropy, all instrument SNVs had *F* statistics exceeding 10, the conventional cutoff for designating strong instruments.<sup>73</sup> To protect our MR estimates from the influence of invalid instruments and violations of MR's core assumptions, we used sensitivity analyses (eg, Steiger directionality test, *ADH1B* instrument, and leave-1-out), MR estimators with relaxed assumptions (eg, weighted median and post-Lasso IVW), and MVMR accounting for possible confounding or mediating traits, which yielded largely consistent results and suggested minimal violations of MR's assumptions.<sup>51</sup> However, causal inference requires triangulating evidence,<sup>74</sup> and

we emphasize that our results should be interpreted in the context of other studies investigating similar questions with different methodologies.<sup>9,11,13,17</sup> Furthermore, we recognize the neuroanatomical phenotypes we analyzed may not fully encapsulate brain damage and caution that our null findings do not imply that alcohol does not affect brain health.

## Conclusions

The results of this study provide evidence that genetically predicted GCT was associated with alcoholic drinks consumed per week and binge drinking frequency after accounting for neuropsychiatric phenotypes, substance use, trauma, and neurodegeneration. We also found that several genes located at 17q21.31 and glutamatergic cortical neurons may be biological mechanisms associating GCT with alcohol consumption. These findings should be replicated in larger samples to better understand the interactions between brain structure and alcohol use.

## ARTICLE INFORMATION

**Accepted for Publication:** June 15, 2022.

**Published Online:** August 10, 2022.

doi:10.1001/jamapsychiatry.2022.2196

**Open Access:** This is an open access article distributed under the terms of the [CC-BY License](#). © 2022 Mavromatis LA et al. *JAMA Psychiatry*.

**Author Contributions:** Messrs Mavromatis and Rosoff had full access to all of the data in this study and take responsibility for the integrity of the data and the accuracy of the data analysis.

**Concept and design:** Mavromatis, Rosoff, Cupertino, Lohoff.

**Acquisition, analysis, or interpretation of data:** Mavromatis, Rosoff, Mackey, Garavan, Lohoff.

**Drafting of the manuscript:** Mavromatis, Rosoff, Lohoff.

**Critical revision of the manuscript for important intellectual content:** All authors.

**Statistical analysis:** Mavromatis, Rosoff.

**Obtained funding:** Garavan, Lohoff.

**Administrative, technical, or material support:** Garavan, Lohoff.

**Supervision:** Lohoff.

**Conflict of Interest Disclosures:** None were reported.

**Funding/Support:** This research was supported by the Medical Research Council Integrative Epidemiology Unit (MRC-IEU, University of Bristol, UK) UK Biobank GWAS Pipeline and National Institutes of Health intramural funding (ZIA-AA000242 to Dr Lohoff), Division of Intramural Clinical and Biological Research of the National Institute on Alcohol Abuse and Alcoholism.

**Role of the Funder/Sponsor:** The funders had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

**Additional Contributions:** We thank the participants and investigators involved in the studies. We also acknowledge the work of the

Enhancing Neuroimaging Genetics Through Meta-Analysis (ENIGMA) consortium and the contributions of the participants and investigators involved in these studies.

**Additional Information:** The mendelian randomization instruments and harmonized datasets required to replicate the study findings are presented in eTable 3 in [Supplement 1](#) and eTables 4-17 in [Supplement 2](#). Code is available from the authors upon request. This study used publicly available analytic methods available at their respective repositories: 2-sample mendelian randomization (<https://mrcieu.github.io/TwoSampleMR/>); mendelian randomization (<https://cran.r-project.org/web/packages/MendelianRandomization/index.html>), FUSION ([https://github.com/gusevlab/fusion\\_twas](https://github.com/gusevlab/fusion_twas)), Collect (<https://github.com/perslab/collect>).

## REFERENCES

- World Health Organization. Global status report on alcohol and health 2014. Accessed August 1, 2021. <https://www.afro.who.int/publications/global-status-report-alcohol-and-health-2014>
- Baumberg B. The global economic burden of alcohol: a review and some suggestions. *Drug Alcohol Rev*. 2006;25(6):537-551. doi:10.1080/09595230600944479
- Crews FT, Sarkar DK, Qin L, Zou J, Boyadjieva N, Vetreno RP. Neuroimmune function and the consequences of alcohol exposure. *Alcohol Res*. 2015;37(2):331-341, 344-351.
- Pfefferbaum A, Rosenbloom MJ, Chu W, et al. White matter microstructural recovery with abstinence and decline with relapse in alcohol dependence interacts with normal ageing: a controlled longitudinal DTI study. *Lancet Psychiatry*. 2014;1(3):202-212. doi:10.1016/S2215-0366(14)70301-3
- Qin L, Crews FT. NADPH oxidase and reactive oxygen species contribute to alcohol-induced microglial activation and neurodegeneration. *J Neuroinflammation*. 2012;9:5. doi:10.1186/1742-2094-9-5
- Marshall SA, Geil CR, Nixon K. Prior binge ethanol exposure potentiates the microglial response in a model of alcohol-induced neurodegeneration. *Brain Sci*. 2016;6(2):16. doi:10.3390/brainsci6020016
- Hammoud N, Jimenez-Shahed J. Chronic neurologic effects of alcohol. *Clin Liver Dis*. 2019;23(1):141-155. doi:10.1016/j.cld.2018.09.010
- Taber KH, Black DN, Porrino LJ, Hurley RA. Neuroanatomy of dopamine: reward and addiction. *J Neuropsychiatry Clin Neurosci*. 2012;24(1):1-4. doi:10.1176/appi.neuropsych.24.1.1
- Baranger DAA, Demers CH, Elsayed NM, et al. Convergent evidence for predispositional effects of brain gray matter volume on alcohol consumption. *Biol Psychiatry*. 2020;87(7):645-655. doi:10.1016/j.biopsych.2019.08.029
- Chen W-JA, Maier SE, Parnell SE, West JR. Alcohol and the developing brain: neuroanatomical studies. *Alcohol Res Health*. 2003;27(2):174-180.
- Hatoum AS, Johnson EC, Agrawal A, Bogdan R. Brain structure and problematic alcohol use: a test of plausible causation using latent causal variable analysis. *Brain Imaging Behav*. 2021;15(6):2741-2745. doi:10.1007/s11682-021-00482-z
- Mackey S, Allgaier N, Chaurani B, et al; ENIGMA Addiction Working Group. Mega-analysis of gray matter volume in substance dependence: general and substance-specific regional effects. *Am J Psychiatry*. 2019;176(2):119-128. doi:10.1176/appi.ajp.2018.17040415
- Robert GH, Luo Q, Yu T, et al; IMAGEN Consortium. Association of gray matter and personality development with increased drunkenness frequency during adolescence. *JAMA Psychiatry*. 2020;77(4):409-419. doi:10.1001/jamapsychiatry.2019.4063
- Schmidt T, Roser P, Ze O, Juckel G, Suchan B, Thoma P. Cortical thickness and trait empathy in patients and people at high risk for alcohol use disorders. *Psychopharmacology (Berl)*. 2017;234(23-24):3521-3533. doi:10.1007/s00213-017-4741-3



15. Thayer RE, YorkWilliams S, Karoly HC, et al. Structural neuroimaging correlates of alcohol and cannabis use in adolescents and adults. *Addiction*. 2017;112(12):2144-2154. doi:10.1111/add.13923
16. Yang X, Tian F, Zhang H, et al. Cortical and subcortical gray matter shrinkage in alcohol-use disorders: a voxel-based meta-analysis. *Neurosci Biobehav Rev*. 2016;66:92-103. doi:10.1016/j.neubiorev.2016.03.034
17. Dager AD, McKay DR, Kent JW Jr, et al. Shared genetic factors influence amygdala volumes and risk for alcoholism. *Neuropsychopharmacology*. 2015;40(2):412-420. doi:10.1038/npp.2014.187
18. Harper J, Malone SM, Wilson S, Hunt RH, Thomas KM, Iacono WG. The effects of alcohol and cannabis use on the cortical thickness of cognitive control and salience brain networks in emerging adulthood: a co-twin control study. *Biol Psychiatry*. 2021;89(10):1012-1022. doi:10.1016/j.biopsych.2021.01.006
19. Luciana M, Collins PF, Muetzel RL, Lim KO. Effects of alcohol use initiation on brain structure in typically developing adolescents. *Am J Drug Alcohol Abuse*. 2013;39(6):345-355. doi:10.3109/00952990.2013.837057
20. Taffe MA, Kotzebue RW, Crean RD, Crawford EF, Edwards S, Mandym CD. Long-lasting reduction in hippocampal neurogenesis by alcohol consumption in adolescent nonhuman primates. *Proc Natl Acad Sci U S A*. 2010;107(24):11104-11109. doi:10.1073/pnas.0912810107
21. Zou X, Durazzo TC, Meyerhoff DJ. Regional brain volume changes in alcohol-dependent individuals during short-term and long-term abstinence. *Alcohol Clin Exp Res*. 2018;42(6):1062-1072. doi:10.1111/acer.13757
22. Marek S, Tervo-Clemmens B, Calabro FJ, et al. Reproducible brain-wide association studies require thousands of individuals. *Nature*. 2022;603(7902):654-660. doi:10.1038/s41586-022-04492-9
23. Davey Smith G, Hemani G. Mendelian randomization: genetic anchors for causal inference in epidemiological studies. *Hum Mol Genet*. 2014;23(R1):R89-R98. doi:10.1093/hmg/ddu328
24. Davey Smith G. Use of genetic markers and gene-diet interactions for interrogating population-level causal influences of diet on health. *Genes Nutr*. 2011;6(1):27-43. doi:10.1007/s12263-010-0181-y
25. Smith GD, Ebrahim S. 'Mendelian randomization': can genetic epidemiology contribute to understanding environmental determinants of disease? *Int J Epidemiol*. 2003;32(1):1-22. doi:10.1093/ije/dyg070
26. Walker VM, Zheng J, Gaunt TR, Smith GD. Phenotypic causal inference using genome-wide association study data: mendelian randomization and beyond. *Annu Rev Biomed Data Sci*. Published online April 1, 2022. doi:10.1146/annurev-biodatasci-122120-024910
27. Treur JL, Demontis D, Smith GD, et al. Investigating causality between liability to ADHD and substance use, and liability to substance use and ADHD risk, using Mendelian randomization. *Addict Biol*. 2021;26(1):e12849. doi:10.1111/adb.12849
28. Wootton RE, Jones HJ, Sallis HM. Mendelian randomisation for psychiatry: how does it work, and what can it tell us? *Mol Psychiatry*. 2022;27(1):53-57. doi:10.1038/s41380-021-01173-3
29. Burgess S, Thompson SG. Multivariable mendelian randomization: the use of pleiotropic genetic variants to estimate causal effects. *Am J Epidemiol*. 2015;181(4):251-260. doi:10.1093/aje/kwu283
30. Wilsnack RW, Wilsnack SC, Kristjansson AF, Vogeltanz-Holm ND, Gmel G. Gender and alcohol consumption: patterns from the multinational GENACIS project. *Addiction*. 2009;104(9):1487-1500. doi:10.1111/j.1360-0443.2009.02696.x
31. Gusev A, Ko A, Shi H, et al. Integrative approaches for large-scale transcriptome-wide association studies. *Nat Genet*. 2016;48(3):245-252. doi:10.1038/ng.3506
32. Timshel PN, Thompson JJ, Pers TH. Genetic mapping of etiologic brain cell types for obesity. *Elife*. 2020;9:e55851. doi:10.7554/eLife.55851
33. Skrivankova VW, Richmond RC, Woolf BAR, et al. Strengthening the Reporting of Observational Studies in Epidemiology using mendelian randomization: the STROBE-MR statement. *JAMA*. 2021;326(16):1614-1621. doi:10.1001/jama.2021.18236
34. Grasby KL, Jahanshad N, Painter JN, et al; Alzheimer's Disease Neuroimaging Initiative; CHARGE Consortium; EPiGen Consortium; IMAGEN Consortium; SYS Consortium; Parkinson's Progression Markers Initiative; Enhancing Neuroimaging Genetics through Meta-Analysis Consortium (ENIGMA)—Genetics working group. The genetic architecture of the human cerebral cortex. *Science*. 2020;367(6484):eaay6690. doi:10.1126/science.aay6690
35. Mueller S, Wang D, Fox MD, et al. Individual variability in functional connectivity architecture of the human brain. *Neuron*. 2013;77(3):586-595. doi:10.1016/j.neuron.2012.12.028
36. Chen C-H, Fiecas M, Gutiérrez ED, et al. Genetic topography of brain morphology. *Proc Natl Acad Sci U S A*. 2013;110(42):17089-17094. doi:10.1073/pnas.1308091110
37. Fischl B. FreeSurfer. *NeuroImage*. 2012;62(2):774-781. doi:10.1016/j.neuroimage.2012.01.021
38. Zhao B, Luo T, Li T, et al; Alzheimer's Disease Neuroimaging Initiative; Pediatric Imaging, Neurocognition and Genetics. Genome-wide association analysis of 19,629 individuals identifies variants influencing regional brain volumes and refines their genetic co-architecture with cognitive and mental health traits. *Nat Genet*. 2019;51(11):1637-1644. doi:10.1038/s41588-019-0516-6
39. Satizabal CL, Adams HHH, Hibar DP, et al. Genetic architecture of subcortical brain structures in 38,851 individuals. *Nat Genet*. 2019;51(11):1624-1636. doi:10.1038/s41588-019-0511-y
40. Liu M, Jiang Y, Wedow R, et al; 23andMe Research Team; HUNT All-In Psychiatry. Association studies of up to 1.2 million individuals yield new insights into the genetic etiology of tobacco and alcohol use. *Nat Genet*. 2019;51(2):237-244. doi:10.1038/s41588-018-0307-5
41. Bycroft C, Freeman C, Petkova D, et al. The UK Biobank resource with deep phenotyping and genomic data. *Nature*. 2018;562(7726):203-209. doi:10.1038/s41586-018-0579-z
42. Walters RK, Polimanti R, Johnson EC, et al; 23andMe Research Team. Transancestral GWAS of alcohol dependence reveals common genetic underpinnings with psychiatric disorders. *Nat Neurosci*. 2018;21(12):1656-1669. doi:10.1038/s41593-018-0275-1
43. Gage SH, Bowden J, Davey Smith G, Munafò MR. Investigating causality in associations between education and smoking: a two-sample Mendelian randomization study. *Int J Epidemiol*. 2018;47(4):1131-1140. doi:10.1093/ije/dyy131
44. Rosoff DB, Smith GD, Lohoff FW. Prescription opioid use and risk for major depressive disorder and anxiety and stress-related disorders: a multivariable mendelian randomization analysis. *JAMA Psychiatry*. 2021;78(2):151-160. doi:10.1001/jamapsychiatry.2020.3554
45. Edenberg HJ. The genetics of alcohol metabolism: role of alcohol dehydrogenase and aldehyde dehydrogenase variants. *Alcohol Res Health*. 2007;30(1):5-13.
46. Abifadel M, Varret M, Rabès J-P, et al. Mutations in PCSK9 cause autosomal dominant hypercholesterolemia. *Nat Genet*. 2003;34(2):154-156. doi:10.1038/ng1161
47. Hemani G, Bowden J, Davey Smith G. Evaluating the potential role of pleiotropy in mendelian randomization studies. *Hum Mol Genet*. 2018;27(R2):R195-R208. doi:10.1093/hmg/ddy163
48. Bowden J, Davey Smith G, Burgess S. Mendelian randomization with invalid instruments: effect estimation and bias detection through Egger regression. *Int J Epidemiol*. 2015;44(2):512-525. doi:10.1093/ije/dyv080
49. Bowden J, Davey Smith G, Haycock PC, Burgess S. Consistent estimation in mendelian randomization with some invalid instruments using a weighted median estimator. *Genet Epidemiol*. 2016;40(4):304-314. doi:10.1002/gepi.21965
50. Hartwig FP, Davey Smith G, Bowden J. Robust inference in summary data mendelian randomization via the zero modal pleiotropy assumption. *Int J Epidemiol*. 2017;46(6):1985-1998. doi:10.1093/ije/dyx102
51. Sanderson E, Glymour MM, Holmes MV, et al. Mendelian randomization. *Nat Rev Methods Primers*. 2022;2(1):6. doi:10.1038/s43586-021-00092-5
52. Bowden J, Del Greco M F, Minelli C, et al. Improving the accuracy of two-sample summary-data mendelian randomization: moving beyond the NOME assumption. *Int J Epidemiol*. 2019;48(3):728-742. doi:10.1093/ije/dyy258
53. Verbanck M, Chen CY, Neale B, Do R. Detection of widespread horizontal pleiotropy in causal relationships inferred from mendelian randomization between complex traits and diseases. *Nat Genet*. 2018;50(5):693-698. doi:10.1038/s41588-018-0099-7
54. Rees JMB, Wood AM, Dudbridge F, Burgess S. Robust methods in mendelian randomization via penalization of heterogeneous causal estimates. *PLoS One*. 2019;14(9):e0222362. doi:10.1371/journal.pone.0222362
55. Hemani G, Tilling K, Davey Smith G. Orienting the causal relationship between imprecisely measured traits using GWAS summary data. *PLoS Genet*. 2017;13(11):e1007081. doi:10.1371/journal.pgen.1007081
56. Fromer M, Roussos P, Sieberts SK, et al. Gene expression elucidates functional impact of polygenic risk for schizophrenia. *Nat Neurosci*. 2016;19(11):1442-1453. doi:10.1038/nn.4399

57. Consortium G; GTEx Consortium. Human genomics. the Genotype-Tissue Expression (GTEx) pilot analysis: multitissue gene regulation in humans. *Science*. 2015;348(6235):648-660. doi:10.1126/science.1262110
58. Map AB. Human Multiple Cortical Areas SMART-seq. Accessed May 5, 2022. <https://portal.brain-map.org/atlas-and-data/maseq/human-multiple-cortical-areas-smart-seq>
59. Logtenberg E, Overbeek MF, Pasman JA, et al. Investigating the causal nature of the relationship of subcortical brain volume with smoking and alcohol use. *Br J Psychiatry*. 2022;221(1):377-385.
60. Nelson EC, Agrawal A, Pergadia ML, et al. H2 haplotype at chromosome 17q21.31 protects against childhood sexual abuse-associated risk for alcohol consumption and dependence. *Addict Biol*. 2010;15(1):1-11. doi:10.1111/j.1369-1600.2009.00181.x
61. Stefansson H, Helgason A, Thorleifsson G, et al. A common inversion under selection in Europeans. *Nat Genet*. 2005;37(2):129-137. doi:10.1038/ng1508
62. de Jong S, Chepelev I, Janson E, et al. Common inversion polymorphism at 17q21.31 affects expression of multiple genes in tissue-specific manner. *BMC Genomics*. 2012;13(1):458. doi:10.1186/1471-2164-13-458
63. Hansson AC, Cipitelli A, Sommer WH, et al. Variation at the rat Crhr1 locus and sensitivity to relapse into alcohol seeking induced by environmental stress. *Proc Natl Acad Sci U S A*. 2006;103(41):15236-15241. doi:10.1073/pnas.0604419103
64. Heilig M, Koob GF. A key role for corticotropin-releasing factor in alcohol dependence. *Trends Neurosci*. 2007;30(8):399-406. doi:10.1016/j.tins.2007.06.006
65. Glaser YG, Zubieta J-K, Hsu DT, et al. Indirect effect of corticotropin-releasing hormone receptor 1 gene variation on negative emotionality and alcohol use via right ventrolateral prefrontal cortex. *J Neurosci*. 2014;34(11):4099-4107. doi:10.1523/JNEUROSCI.3672-13.2014
66. Uribe-Mariño A, Gassen NC, Wiesbeck MF, et al. Prefrontal cortex corticotropin-releasing factor receptor 1 conveys acute stress-induced executive dysfunction. *Biol Psychiatry*. 2016;80(10):743-753. doi:10.1016/j.biopsych.2016.03.2106
67. Ivy AS, Rex CS, Chen Y, et al. Hippocampal dysfunction and cognitive impairments provoked by chronic early-life stress involve excessive activation of CRH receptors. *J Neurosci*. 2010;30(39):13005-13015. doi:10.1523/JNEUROSCI.1784-10.2010
68. Wang X-D, Su Y-A, Wagner KV, et al. Nectin-3 links CRHR1 signaling to stress-induced memory deficits and spine loss. *Nat Neurosci*. 2013;16(6):706-713. doi:10.1038/nn.3395
69. Zhang C-C, Xing A, Tan M-S, Tan L, Yu J-T. The role of MAPT in neurodegenerative diseases: genetics, mechanisms and therapy. *Mol Neurobiol*. 2016;53(7):4893-4904. doi:10.1007/s12035-015-9415-8
70. Quednow BB, Herdener M. Human pharmacology for addiction medicine: from evidence to clinical recommendations. *Prog Brain Res*. 2016;224:227-250. doi:10.1016/bs.pbr.2015.07.017
71. Rao PSS, Bell RL, Engleman EA, Sari Y. Targeting glutamate uptake to treat alcohol use disorders. *Front Neurosci*. 2015;9:144. doi:10.3389/fnins.2015.00144
72. Refojo D, Schweizer M, Kuehne C, et al. Glutamatergic and dopaminergic neurons mediate anxiogenic and anxiolytic effects of CRHR1. *Science*. 2011;333(6051):1903-1907. doi:10.1126/science.1202107
73. Burgess S, Davies NM, Thompson SG. Bias due to participant overlap in two-sample mendelian randomization. *Genet Epidemiol*. 2016;40(7):597-608. doi:10.1002/gepi.21998
74. Hammerton G, Munafò MR. Causal inference with observational data: the need for triangulation of evidence. *Psychol Med*. 2021;51(4):563-578. doi:10.1017/S0033291720005127