



AMIDD 2023 Lecture 5: Five key questions of drug discovery




Medical Need




**Target(s) &
modality**



PK/PD



Benefit/risk



**Patient
stratification**

Dr. Jitao David Zhang, Computational Biologist

¹ *Pharmaceutical Sciences, Pharma Research and Early Development, Roche Innovation Center Basel, F. Hoffmann-La Roche*

² *Department of Mathematics and Informatics, University of Basel*

Today's goals

1. Five questions and a *meta*-question for drug discovery and development
2. Case study: mRNA vaccines for coronavirus

Interests and concerns of companies working on drug discovery: summary of our previous discussions

Interests

- Return of Investment
 - Commercial potential
 - Cycle time
- Good reputation
 - Efficacy of the drug
 - Safety of the drug
 - Market access
- Environmental, social, and governance (e.g. fighting internal corruption, diversity of board members).

Concerns

- Low or no return of investment
 - Lack of efficacy of drugs
 - Unfavorable benefit/risk profiles of the drug
 - No approval from agency
 - Cost, time, effectiveness of R&D
 - Competitor
 - Poor targets or disease models due to lack of reproducibility of published data
 - Companion diagnostic
- Intellectual property
- Idea and knowledge management
- Acceptance by doctors and patients
- Legal of concern

Five key questions to ask before we start and along the way

1. What is the unmet medical need to be addressed?
2. What are the target(s) and what is the modality of our drug?
3. Where should the drug go in patient's body, what does body do to the drug, and what does the drug do to the body?
4. What is the safety profile of the drug in light of its benefits?
5. Who are responsive to the drug, or susceptible to adverse events?

The *meta*-question: What knowledge, data, and tools do we have to address these questions?

Right target

- Strong link between target and disease
- Differentiated efficacy
- Available and predictive biomarkers

Right tissue

- Adequate bioavailability and tissue exposure
- Definition of PD biomarkers
- Clear understanding of preclinical and clinical PK/PD
- Understanding of drug–drug interactions

Right safety

- Differentiated and clear safety margins
- Understanding of secondary pharmacology risk
- Understanding of reactive metabolites, genotoxicity and drug–drug interactions
- Understanding of target liability

Right patient


- Identification of the most responsive patient population
- Definition of risk–benefit for a given population

Right commercial potential

- Differentiated value proposition versus future standard of care
- Focus on market access, payer and provider
- Personalized health-care strategy, including diagnostics and biomarkers

Morgan, P. et al. [Impact of a five-dimensional framework on R&D productivity at AstraZeneca](#). *Nature Reviews Drug Discovery* 17, 167–181 (2018).


Chevron diagrams as a pipeline view of drug discovery and development



FDA: US Food and Drug Administration. Top: Wagner, J. A. et al. [Application of a Dynamic Map for Learning, Communicating, Navigating, and Improving Therapeutic Development](#). Clinical and Translational Science 11, 166–174 (2018). Bottom: Adapted from Paul et al. [How to Improve R&D Productivity: The Pharmaceutical Industry's Grand Challenge](#). Nature Reviews Drug Discovery, 2010.


A dynamic map for drug discovery, development, and deployment

1. **Basic science research and target identification.** *What causes the disease? What do we want to achieve? Which protein can I target with which modality?*
2. **Target pharmacology and biomarker development.** *What is the effect of targeting the protein? What we can measure to confirm that the protein is properly targeted?*
3. **Lead identification.** *How can we find a starting point of a new drug?*
4. **Lead optimization and clinical candidate selection.** *What are criteria to define a good drug? How can I improve the starting material?*
5. **Clinical research and development.** *Does it work in human? How about efficacy and safety profiles?*
6. **Regulatory review.** *Should we approve the drug?*
7. **Post marketing.** *How does the drug work in real world?*



Wagner, J. et al. [A dynamic map for learning, communicating, navigating and improving therapeutic development.](#) Nat Rev Drug Discov 17, 150–150 (2018).

The multiscale modelling view of drug discovery



Drug Discovery Today


Zhang, Jitao David, Lisa Sach-Peltason, Christian Kramer, Ken Wang, and Martin Ebeling. 2020. "Multiscale Modelling of Drug Mechanism and Safety." *Drug Discovery Today* 25 (3): 519–34. <https://doi.org/10.1016/j.drudis.2019.12.009>.

Nobel Prize in Physiology or Medicine 2023 was awarded to Katalin Karikó and Drew Weissman for “their discoveries concerning nucleoside base modifications that enabled the development of effective mRNA vaccines against COVID-19”



Methods for vaccine production before the COVID-19 pandemic.

Issues: large-scale cell culture required, which limits the possibilities for rapid production in response to pandemics.




© The Nobel Committee for Physiology or Medicine. III. Mattias Karlén

Vaccine mimics viral infection to activate the immune system to protect body from future infections

Vaccine mimics a viral infection to activate innate and adaptive immune system, while minimizing the pathogenic effects.


Key players in the game:

1. Viral proteins as *antigens*
2. Antigen-presenting cells (e.g. dendritic cells)
3. T cells (T comes from Thymus, because they mature there)
4. B cells (B comes from bone marrow).



Pollard, A. J. & Bijker, E. M. A guide to vaccinology: from basic principles to new developments. Nature Reviews Immunology 21, 83–100 (2021).

Virus is special




- The three-domain model of *cellular* life: (eu-)bacteria, archaeobacteria, and eukaryotes.
- The two-domain model: bacteria as one branch, archaea and eukaryotes as the other.
- Virus fits in no domain of neither models.

1. Woese, C. R., Kandler, O. & Wheelis, M. L. Towards a natural system of organisms: proposal for the domains Archaea, Bacteria, and Eucarya. Proc Natl Acad Sci U S A 87, 4576–4579 (1990). Figure from [Wikimedia](#), reused with CC-AS license.


1. Hug, L. A. et al. A new view of the tree of life. Nat Microbiol 1, 1–6 (2016). Figure from [Wikimedia](#), CC-AS license.

1. Forterre, P., Krupovic, M. & Prangishvili, D. Cellular domains and viral lineages. Trends in Microbiology 22, 554–558 (2014).




Coronavirus is a RNA virus infecting human and other species

Viral particle




The coronavirus virion consists of structural proteins, namely spike (S), envelope (E), membrane (M), nucleocapsid (N)



Phylogenetic relationships of representative members of the species Severe acute respiratory syndrome-related coronavirus



V'kovski, P., Kratzel, A., Steiner, S., Stalder, H. & Thiel, V. Coronavirus biology and replication: implications for SARS-CoV-2. Nat Rev Microbiol 19, 155–170 (2021).

Sequence of the spike protein is largely conserved between corona and related viruses



Andersen *et al.*, [The Proximal Origin of SARS-CoV-2](#), Nature Medicine, 2020

Spike protein of coronavirus is responsible for viral entry into human cells





Huang, Y., Yang, C., Xu, X., Xu, W. & Liu, S. Structural and functional properties of SARS-CoV-2 spike protein: potential antivirus drug development for COVID-19. *Acta Pharmacol Sin* 41, 1141–1149 (2020).

DNA or RNA encodes genetic information of all life forms that we know, including viruses



Photo 51, X-ray diffraction image of DNA

Franklin R, Gosling RG (1953)
"Molecular Configuration in Sodium Thymonucleate". *Nature* 171: 740–741.



From the textbook OpenStax Anatomy and Physiology, discovered through Wikimedia, reused under the CC license.

RNA is transcribed from DNA and translated into protein






Figure: https://commons.wikimedia.org/wiki/File:DNA_transcriptie.svg and https://en.m.wikipedia.org/wiki/File%3AHAR1F_RF00635_rna_secondary_structure.jpg. Original work by wikipedia user: OrgreBot and user:Ppgardne. Used under CC-SA 3.0 license.

A summary of what we have learned so far in the context of coronavirus


1. What is the unmet medical need to be addressed? **We need a vaccine to prevent a large population of individuals from being infected by coronavirus, which have severe consequences.**
2. What are the target(s) of our drug? **Spike protein is conserved: immune reaction is desired.**
3. Where should the drug go in patient's body, what does body do to the drug, and what does the drug do to the body? **Due to time constraints, classical vaccine may not meet the need. How about mRNA vaccines?**
4. What is the safety profile of the drug in light of its benefits? **To be investigated.**
5. Who are responsive to the drug, or susceptible to adverse events? **To be investigated.**

Three essential challenges for mRNA-based therapies: delivery, stability, and *unwanted* immune responses



Key challenges:

- mRNAs are too large and charged to pass lipid bilayers.
- mRNAs are readily degraded by ribonucleases.
- *Exogeneous* mRNAs cause *Immunogenicity*.




Left: Cell membrane, copyright of Encyclopædia Britannica, Inc. Right: The four-billion-year-old barrier to RNA therapeutic


Lipid NanoParticles (LNP) helps delivering RNAs into cells

- Lipid nanoparticles can take mRNA vaccines as cargos, and deliver them into human cells.
- In the cell, mRNA encoding the part of the spike protein sequence is translated into proteins with the human protein translation mechanism.
- Synthesized proteins will be degraded and exposed on cell surface, which will be recognized by antigen presenting cells.

Salleh, Mohd Zulkifli *et al.* "[Immunogenicity Mechanism of mRNA Vaccines and Their Limitations in Promoting Adaptive Protection against SARS-CoV-2.](#)" PeerJ 10 (March 9, 2022)



RNAs are degraded by ribonucleases (RNAases)



- mRNAs are too large and charged to pass lipid bilayers.
- **mRNAs are degradable by ribonucleases (RNases).**
- mRNAs are rapidly cleared from liver and kidney.
- *Exogeneous* mRNAs induce *immunogenicity*.

		<u>1</u>		<u>10</u>		<u>20</u>				<u>40</u>		
RNase 1	1	---	KESRAKKE	FQR	QH	MDSDSSPSSS	STY	C	NQ	MRRRN	MTQGRCKPV	
RNase 2	2	KPPQ	FTWAQW	FET	QH	INMTSQ	-----	Q	CTNA	QVINNY	QRCKNQ	
RNase 3	3	RPPQ	FTWAQW	FAI	QH	ISLNPP	-----	R	CTIA	MRAINNY	RWRCKNQ	
RNase 4	4	--	<QDGM	YQRF	LR	QH	VHPEET	-	GGSD	RYCNLM	QRRKMTLYHCKRF	
RNase 5	5	--	<QDNS	RYTH	FL	TO	HYDAKPQ	-	GRDD	RYCESI	MRRRGLTS-PCKDI	
RNase 6	6	WPKRL	TKAHW	FET	QH	IQPSPL	-----	Q	CNRA	MSGINNY	TOHCKHQ	
RNase 7	7	KPKGM	TSSQW	FKI	QH	MQPSQ	-----	A	CNSA	MKNINKH	TKCKDL	
RNase 8	8	KPKDM	TSSQW	FET	QH	VQPSQ	-----	A	CNSA	MSIINKY	TERCKDL	
							↓				↓	
		<u>50</u>		<u>60</u>		<u>70</u>		<u>80</u>		<u>90</u>		
RNase 1	1	NTTF	VHEPL	VDVQ	NVCF	QEKV	TCK	NGQ	--	GNCY	KSNSSMHITDCRLTNGS	
RNase 2	2	NTFLL	TTFAN	VVNV	CGN	PNMT	CPS	NKTR	KNCH	HSGSQV	PLIHCCNLTTPS	
RNase 3	3	NTFL	RTTF	ANVV	NVCG	NQSIR	CPH	NRTL	NNCH	RSRFRV	PLLHCCOLINPG	
RNase 4	4	NTFI	HEDI	WNIR	SIC	STTN	IQC	KNGK	--	MNCHE	GV--VKVTD CRTDGSS	
RNase 5	5	NTFI	HGNKR	SIAI	CEN	KNG	--	NPHRE	-	NLRISK	SSFQVTTCKLHGGG	
RNase 6	6	NTFL	HDSF	QNVAA	VCD	LLSIV	CKN	RNR	--	HNCH	QSSKPVNMTDCRLTSG-	
RNase 7	7	NTFL	HEPF	SSVA	ATC	QTPK	IAC	KNGD	--	KNCH	QSHGVPVSLTMCKLTSG-	
RNase 8	8	NTFL	HEPF	SSVA	ITC	QTPN	IAC	KNSC	--	KNCH	QSHGPMSTLTMGELTSG-	
									▲		▲	
		<u>100</u>		<u>110</u>		<u>120</u>						
RNase 1	1	--	RYPN	CA	YRTS	PKER	HI	I	VACE	-----	GSPYV	VPVHFDASVEDST
RNase 2	2	PQNIS	NCRY	AQT	PANM	FYIV	ACD	NRD	QRRD	PPQY	PVVPVHLDRII----	
RNase 3	3	AQNIS	NCRY	AD	PRGR	FRFY	VVAC	DNR	DPR	-	DS	PRYPVVPVHLDITTI----
RNase 4	4	--	RAPN	CRY	RAI	ASTR	RV	VIACE	-----	GNPQV	VPVHFDG-----	
RNase 5	5	--	PWPP	CQY	RAT	AGFR	NV	VVACE	-----	NG---	LPVHLDQSIFRRP	
RNase 6	6	--	KYPQ	CRY	SAA	AQYK	FFIV	ACD	-	PPQK	SDPP-YKLVPVHLDQSIL----	
RNase 7	7	--	KYPN	CRY	KEK	RQNK	SYV	VACK	-	PPQK	KDSQFHLVPVHLDRLV----	
RNase 8	8	--	KYPN	CRY	KEK	HLN	TPY	IVACD	-	PPQQ	GDGP-YPLVPVHLDKVV----	
												↓


Left: Structure of PDB [7RSA](#). Right: alignment of protein sequences of 8 canonical human RNases (ribonuclease A family). [Sorrentino FEBS Letters, 2010](#).

Unmodified RNA induces unwanted immune reactions: modified RNAs instead have reduced or little unwanted effect

Exogeneous mRNAs induce Immunogenicity. All RNAs is synthesize from four ribonucleotides: ATP (adenosine triphosphate), CTP (cytidine triphosphate), UTP (uridine triphosphate), and GTP (guanosine triphosphate).


When unmodified RNAs are delivered into cells, they induce unwanted immune reaction, e.g. the release of the tumor necrosis factor alpha (TNF-alpha), by activating the Toll-like receptors (TLRs), which defend against bacterial and viral infections.

Some type of RNA, however, does not induce immunogenicity, for instance human *tRNA*. This finding by Karikó and Drewman made major contributions to successful development of SARS-CoV-2 mRNA vaccines.





Karikó, K., Buckstein, M., Ni, H. & Weissman, D. Suppression of RNA Recognition by Toll-like Receptors: The Impact of Nucleoside Modification and the Evolutionary Origin of RNA. *Immunity* 23, 165–175 (2005).


tRNAs transfer amino-acids to ribosome for protein translation, and human tRNA contains *pseudouridine*, a modified uridine



Left: Role of tRNA in protein translation. Top right: tRNA structure, with the TΨC loop highlighted in the blue ellipse. Below: structures of uridine and pseudouridine.




Uridine (U)




Pseudouridine (psi, Ψ)

By Boumphreyfr vector conversion by Glrx - File:Peptide syn.png, CC BY-SA 3.0, <https://commons.wikimedia.org/w/index.php?curid=101457889>. By Yikrazuul, CC BY-SA 3.0, via Wikimedia Commons.

Further modification (N1-methyl-Ψ) combined with LNP delivery lead to the currently used vaccines




mRNA vaccines against human SARS-Cov-2 viruses, developed in 2020 by Pfizer-BioNTech and Moderna Therapeutics (comirnaty® and spikevax®, respectively), reached clinical efficacies higher than 90%. Both used modified RNA and LNP. Curevac mRNA vaccine (CVnCoV), which used LNP but not modified RNA, reached an efficacy of 48%.



Morais, P., Adachi, H. & Yu, Y.-T. The Critical Contribution of Pseudouridine to mRNA COVID-19 Vaccines. Front Cell Dev Biol 9, 789427 (2021).

Coding sequence of the spike protein along is not enough: mRNA transcription depends on 5'-UTR and 3'-UTR, too




The structure of a gene and the control of gene expression in eukaryotes:

1. RNA polymerase, an enzyme, binds to the promoter region and reads the DNA from the 5' UTR to the 3' UTR.
2. pre-mRNA is synthesized and receives a cap of a modified nucleotide (7-methylguanosine triphosphate) to the 5' end, and a repeated adenine sequence (a poly-A tail) to the 3' end.
3. pre-mRNA is spliced to remove introns. Mature mRNA contains the 5' cap, 5'-untranslated region (5'-UTR), coding sequence, 3'-untranslated region (3'-UTR), and a poly-A tail.
4. mRNAs are transported from the nucleus to the cytoplasm for translation.

Zerbini, F., Da Silva, F., Urquiza, G. & Basso, M. F. [Transgenic Plants](#). in *Biotechnology and Plant Breeding: Applications and Approaches for Developing Improved Cultivars* 179–199 (2014).


5'-UTR of human hemoglobin is a good choice to make sure that the vaccine sequence is stable and highly translated



	-MVLSPADKTNVKAAGWGRGAHAGEYGAEALERMFSLFPTTKTYFPHPFD-----LSHGSG	53	P69905	HBA_HUMAN
1	MVHLTPEEKSAVTALWGKVNV--DEVGGEALGRLLVVYPWTQRFFESFGDLSTPDAVMGN	58	P68871	HBB_HUMAN
1	MVHLTPEEKTAVNALWGKVNV--DAVGGEALGRLLVVYPWTQRFFESFGDSSPDAMVGN	58	P02042	HBD_HUMAN
	: *: * : *: *. * ***** . * . *** * :: : * * : * *			
54	AQVKGHGGKKVADALTNAVAVHDMPNALSALSDLHAHKLRVDPVNFKLLSHCLLVILAAH	113	P69905	HBA_HUMAN
59	PKVKAHGKKVGLGFSDGLAHLNLKGTFATLSELHC DKLHVDPENFRLLGNVLVCVLAHH	118	P68871	HBB_HUMAN
59	PKVKAHGKKVGLGFSDGLAHLNLKGTFSQLSE LHC DKLHVDPENFRLLGNVLVCVLARN	118	P02042	HBD_HUMAN
	::*.***** *: :::*: :::: :: **:*:.****** ****.. *: . ** .			
114	LPAEFTPAVASLDKFLASVSVLTISKYR	142	P69905	HBA_HUMAN
119	FGKEFTPVPQAAYQKVVAGVANALAHKYH	147	P68871	HBB_HUMAN
119	FGKEFTPQM QAAYQKVVAGVANALAHKYH	147	P02042	HBD_HUMAN
	: **** : : : : *: : . * : : : :			

- Hemoglobin (left) is a iron-containing protein that transports oxygen.
- The protein present in erythrocytes of almost all vertebrates.
- The protein is essential, therefore the mRNA is relatively stable and highly translated.

LNP, modified RNA, and 5'-UTR of HBA are all essential to make effective *and safe* vaccines against coronavirus



References: Heinz, Franz X., and Karin Stiasny. "Distinguishing Features of Current COVID-19 Vaccines: Knowns and Unknowns of Antigen Presentation and Modes of Action." *Npj Vaccines* 6, no. 1 (August 16, 2021): 1–13. <https://doi.org/10.1038/s41541-021-00369-6>; [Assemblies of putative SARS-CoV2-spike-encoding mRNA sequences for vaccines BNT-162b2 and mRNA-1273](https://github.com/NAalytics/Assemblies_of_putative_SARS-CoV2-spike-encoding_mRNA_sequences_for_vaccines_BNT-162b2_and_mRNA-1273) (github.com/NAalytics); Xia, Xuhua. "Detailed Dissection and Critical Evaluation of the Pfizer/BioNTech and Moderna MRNA Vaccines." *Vaccines* 9, no. 7 (July 3, 2021): 734. <https://doi.org/10.3390/vaccines9070734>.

A summary of what we have learned so far in the context of coronavirus

1. What is the unmet medical need to be addressed? [We need a vaccine to prevent a large population of individuals from being infected by coronavirus, which have severe consequences.](#)
2. What are the target(s) of our drug? [Spike protein is conserved: immune reaction is desired.](#)
3. Where should the drug go in patient's body, what does body do to the drug, and what does the drug do to the body? [Thanks to LNP, N1-mythel-Ψ, and 5'-UTR of HBA1, the mRNA vaccination can enter cells with minimal side effects. In cells, spike protein RNA is synthesized into proteins, which are digested, presented, and elicit immune response.](#)
4. What is the safety profile of the drug in light of its benefits? [Initial study: Polack, F. P. et al. Safety and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine. New England Journal of Medicine 383, 2603–2615 \(2020\), and watch \[this video\]\(#\).](#)
5. Who are responsive to the drug, or susceptible to adverse events? [Updated regularly by regulatory agencies, for instance \[European Medicines Agency\]\(#\).](#)

Conclusions

1. Drug discovery is an interdisciplinary effort to solve medical and technical challenges.
2. Biological understanding, including sequence analysis, is key for indication and target selection.
3. Modern drug discovery needs to address five key questions:
 - a. **unmet medical need**
 - b. **target(s) and modalities**
 - c. **pharmacokinetics** (what body does to the drug) **and pharmacodynamics** (what the drug does to the body)
 - d. **safety** (risk/benefit assessment)
 - e. **patient enrichment/stratification**