

# Variational form and iterative methods

$$\rho c(T)T_t = \nabla \cdot (k(T)\nabla T) \quad \mathbf{x} \in \Omega \quad (1)$$

With Robin boundary condition:

$$-k(T)\frac{\partial T}{\partial n} = h(T)(T - T_s(t)), \quad \mathbf{x} \in \partial\Omega \quad (2)$$

## (a) The variational form

We apply a Backward Euler discretization in time:

$$\begin{aligned} \rho c(u)^n \frac{u^n - u^{n-1}}{\Delta t} &= [\nabla \cdot (k(u)\nabla u)]^n \\ \rho c(u^n)u^n &= \rho c(u^n)u^{n-1} + \Delta t[\nabla \cdot (k(u^n)\nabla u^n)] \end{aligned} \quad (3)$$

The residual reads:

$$R = \rho c(u^n)u^n - \rho c(u^n)u^{n-1} - \Delta t[\nabla \cdot (k(u^n)\nabla u^n)] \quad (4)$$

We use the Galerkin method to derive the variational form:

$$\begin{aligned} (R, v) &= 0 \\ (R, v) &= (\rho c(u^n)u^n, v) - (\rho c(u^n)u^{n-1}, v) - \Delta t(\nabla \cdot (k(u^n)\nabla u^n), v) \end{aligned} \quad (5)$$

We reformulate the last term using integration by parts and the boundary condition

$$\begin{aligned} (\nabla \cdot (k(u^n)\nabla u^n), v) &= \int_{\Omega} \nabla \cdot (k(u^n)\nabla u^n)v \, dx \\ &= \int_{\partial\Omega} (k(u^n)\nabla u^n)v \cdot \mathbf{n} \, dx - \int_{\Omega} (k(u^n)\nabla u^n) \cdot \nabla v \, dx \\ &= - \int_{\partial\Omega} h(u^n)(u^n - T_s(t))v \, dx - \int_{\Omega} (k(u^n)\nabla u^n) \cdot \nabla v \, dx \end{aligned} \quad (6)$$

The variational form now looks as follows:

$$\int_{\Omega} \rho c(u^n)u^n v \, dx = \int_{\Omega} \rho c(u^n)u^{n-1}v \, dx - \Delta t \int_{\Omega} (k(u^n)\nabla u^n) \cdot \nabla v \, dx - \Delta t \int_{\partial\Omega} h(u^n)(u^n - T_s(t))v \, dx \quad (7)$$

### (b) Picard Iteration

We use  $u^-$  as a guess for  $u^n$  in all the nonlinear terms. To simplify the notation we use  $u^n = u$  and  $u^{n-1} = u_1$ . The linear system we need to solve is then:

$$\int_{\Omega} \rho c(u^-) u v + \Delta t \int_{\Omega} k(u^-) \nabla u \cdot \nabla v \, dx + \quad (8)$$

$$\Delta t \int_{\partial\Omega} h(u^-) (u - T_s(t)) v \, dx = \int_{\Omega} \rho c(u^-) u_1 v \, dx \quad (9)$$

$u$  can be expressed in terms of the basisfunctions  $u = \sum c_j \phi_j$ :

$$\begin{aligned} & \sum c_j \int_{\Omega} \rho c(u^-) \phi_j \phi_i \, dx \\ & + \Delta t \sum c_j \int_{\Omega} k(u^-) \nabla \phi_j \cdot \nabla \phi_i \, dx \\ & + \Delta t \sum c_j \int_{\partial\Omega} h(u^-) \phi_j \phi_i \, dx \\ & = \int_{\Omega} \rho c(u^-) u_1 \phi_i \, dx + \Delta t \int_{\partial\Omega} h(u^-) T_s(t) \phi_i \, dx \end{aligned} \quad (10)$$

### (c) Newtons Method on the variational form

$$\begin{aligned} F(u) &= \int_{\Omega} \rho c(u^n) u^n v \\ &\quad - \int_{\Omega} \rho c(u^n) u^{n-1} v \, dx \\ &\quad + \Delta t \int_{\Omega} (k(u^n) \nabla u^n) \cdot \nabla v \, dx \\ &\quad + \Delta t \int_{\partial\Omega} h(u^n) (u^n - T_s(t)) v \, dx \end{aligned}$$

We use that  $u = \sum c_j \phi_j$ , and  $u^n = u$ ,  $u^{n-1} = u_1$ :

$$\begin{aligned}
F_i(c) = & \sum c_j \int_{\Omega} \rho c (\sum c_k \phi_k) \phi_j \phi_i \, dx \\
& + \Delta t \sum c_j \int_{\Omega} k (\sum c_k \phi_k) \nabla \phi_j \cdot \nabla \phi_i \, dx \\
& + \Delta t \sum c_j \int_{\partial\Omega} h (\sum c_k \phi_k) \phi_j \phi_i \, dx \\
& - \int_{\Omega} \rho c (\sum c_k \phi_k) u_1 \phi_i \, dx \\
& - \Delta t \int_{\partial\Omega} h (\sum c_k \phi_k) T_s(t) \phi_i \, dx
\end{aligned} \tag{11}$$

We need the Jacobian  $J_{i,j} = \frac{\partial F_i}{\partial c_j}$ :

$$\begin{aligned}
J_{i,j} = & \int_{\Omega} \rho c (\sum c_k \phi_k) \phi_j \phi_i \, dx + \sum c_m \int_{\Omega} \rho c' (\sum c_k \phi_k) \phi_j \phi_m \phi_i \, dx \\
& + \Delta t \int_{\Omega} k (\sum c_k \phi_k) \nabla \phi_j \cdot \nabla \phi_i \, dx + \Delta t \sum c_m \int_{\Omega} k' (\sum c_k \phi_k) \phi_j \nabla \phi_m \cdot \nabla \phi_i \, dx \\
& + \Delta t \int_{\partial\Omega} h (\sum c_k \phi_k) \phi_j \phi_i \, dx + \Delta t \sum c_m \int_{\partial\Omega} h' (\sum c_k \phi_k) \phi_j \phi_m \phi_i \, dx \\
& - \int_{\Omega} \rho c' (\sum c_k \phi_k) \phi_j u_1 \phi_i \, dx \\
& - \Delta t \int_{\partial\Omega} h' (\sum c_k \phi_k) \phi_j T_s(t) \phi_i \, dx
\end{aligned} \tag{12}$$

In the iteration we use  $J(u^-)$

$$\begin{aligned}
J(u^-) = & \int_{\Omega} \rho c(u^-) \phi_j \phi_i \, dx + \int_{\Omega} \rho c'(u^-) u^- \phi_j \phi_i \, dx \\
& + \Delta t \int_{\Omega} k(u^-) \nabla \phi_j \cdot \nabla \phi_i \, dx + \Delta t \int_{\Omega} k'(u^-) \phi_j \nabla u^- \cdot \nabla \phi_i \, dx \\
& + \Delta t \int_{\partial\Omega} h(u^-) \phi_j \phi_i \, dx + \Delta t \int_{\partial\Omega} h'(u^-) u^- \phi_j \phi_i \, dx \\
& - \int_{\Omega} \rho c'(u^-) \phi_j u_1 \phi_i \, dx \\
& - \Delta t \int_{\partial\Omega} h'(u^-) \phi_j T_s(t) \phi_i \, dx
\end{aligned} \tag{13}$$

One can now solve  $J(u^-) \delta u = -F(u^-)$

**(d) Newtons Method at PDE level**

$$F(u) = \rho c(u)u - \rho c(u)u_1 - \Delta t[\nabla \cdot (k(u)\nabla u)] \quad (14)$$

We insert  $u = u^- + \delta u$  in  $F(u)$ :

$$F(u^- + \delta u) = \rho c(u^- + \delta u)(u^- + \delta u) - \rho c(u^- + \delta u)u_1 - \Delta t[\nabla \cdot (k(u^- + \delta u)\nabla(u^- + \delta u))] \quad (15)$$

We Tayloexpand the functions around  $u^-$

$$\begin{aligned} c(u^- + \delta u) &= c(u^-) + c'(u^-)\delta u + O(\delta u^2) \\ k(u^- + \delta u) &= k(u^-) + k'(u^-)\delta u + O(\delta u^2) \end{aligned} \quad (16)$$

We keep terms up to order  $\delta u$ :

$$\begin{aligned} F(u^- + \delta u) &= \rho c(u^-)(u^- + \delta u) + \rho c'(u^-)\delta u u^- - \rho(c(u^-) + c'(u^-)\delta u)u_1 \\ &\quad - \Delta t[\nabla \cdot (k(u^-)\nabla(u^- + \delta u) + k'(u^-)\delta u \nabla u^-)] \end{aligned} \quad (17)$$

This gives a linear system for  $\delta u$

$$\begin{aligned} &\rho[c(u^-) + c'(u^-)u^- - c'(u^-)u_1]\delta u \\ &\quad - \Delta t[\nabla \cdot (k(u^-)\nabla \delta u + k'(u^-)\nabla u^- \delta u)] \\ &= -\rho c(u^-)u^- + \rho c(u^-)u_1 + \Delta t[\nabla \cdot (k(u^-)\nabla u^-)] \end{aligned} \quad (18)$$

Variational form:

$$\begin{aligned} &\rho([c(u^-) + c'(u^-)u^- - c'(u^-)u_1]\delta u, v) \\ &\quad + \Delta t([(k(u^-)\nabla \delta u + k'(u^-)\nabla u^- \delta u)], \nabla v) + B.T \\ &= -\rho(c(u^-)(u^- + u_1), v) - \Delta t(k(u^-)\nabla u^-, \nabla v) + B.T \end{aligned} \quad (19)$$