

1 GFN2-xTB E^Γ

$$E^\Gamma = \frac{1}{3} \sum_A \sum_{\mu \in A} (q_{A,\mu})^3 \Gamma_{A,\mu} \quad (1)$$

where $q_{A,\nu} = \sum_B \sum_{\nu \in B} P_{\mu\nu} S_{\mu\nu}$ is the partial charge of shell μ associated with atom A . P, S are the density and overlap matrices. $\Gamma_{A,\mu} = \Gamma_A K_\mu$ is just the product of an element specific constant and a shell specific constant, for our purposes the element is always carbon and the shell is either the first or second in GFN2 thus we have 2 numbers $\Gamma_{\text{Carbon},1(2)}$ henceforth referred to as $\Gamma_{1(2)}$.

Let us first rewrite the inner expression a bit given our new definition and knowledge of the atoms we are working with.

$$\sum_{\mu \in A} (q_{A,\mu})^3 \Gamma_{A,\mu} = \sum_{\mu=1}^2 (q_{A,\mu})^3 \Gamma_\mu \quad (2)$$

We now need a unitary which computes this function on a given state $|q_{A,\mu}\rangle |\Gamma_\mu\rangle |0\rangle \rightarrow |q_{A,\mu}\rangle |\Gamma_\mu\rangle |(q_{A,\mu})^3 \Gamma_\mu\rangle$. Consider having access to the following addition (multiplication) unitary $|A\rangle |B\rangle \rightarrow |A\rangle |A + (*)B\rangle$ as well as the unitary taking $|A\rangle \rightarrow |A^2\rangle$. We can now apply the following to get our desired unitary.

$$\text{MULT}_{x,y}^\dagger \text{SQR}_x^\dagger \text{MULT}_{x,y}^\dagger \text{ADD}_{y,z} \text{MULT}_{x,y} \text{SQR}_x \text{MULT}_{x,y} |q_{A,\mu}\rangle_x |\Gamma_\mu\rangle_y |0\rangle_z \quad (3)$$

$$= \text{MULT}_{x,y}^\dagger \text{SQR}_x^\dagger \text{MULT}_{x,y}^\dagger \text{ADD}_{y,z} \text{MULT}_{x,y} \text{SQR}_x |q_{A,\mu}\rangle_x |\Gamma_\mu q_{A,\mu}\rangle_y |0\rangle_z \quad (4)$$

$$= \text{MULT}_{x,y}^\dagger \text{SQR}_x^\dagger \text{MULT}_{x,y}^\dagger \text{ADD}_{y,z} \text{MULT}_{x,y} |(q_{A,\mu})^2\rangle_x |\Gamma_\mu q_{A,\mu}\rangle_y |0\rangle_z \quad (5)$$

$$= \text{MULT}_{x,y}^\dagger \text{SQR}_x^\dagger \text{MULT}_{x,y}^\dagger \text{ADD}_{y,z} |(q_{A,\mu})^2\rangle_x |(q_{A,\mu})^3 \Gamma_\mu\rangle_y |0\rangle_z \quad (6)$$

$$= \text{MULT}_{x,y}^\dagger \text{SQR}_x^\dagger \text{MULT}_{x,y}^\dagger |(q_{A,\mu})^2\rangle_x |(q_{A,\mu})^3 \Gamma_\mu\rangle_y |(q_{A,\mu})^3 \Gamma_\mu\rangle_z \quad (7)$$

$$= \text{MULT}_{x,y}^\dagger \text{SQR}_x^\dagger |(q_{A,\mu})^2\rangle_x |\Gamma_\mu q_{A,\mu}\rangle_y |(q_{A,\mu})^3 \Gamma_\mu\rangle_z \quad (8)$$

$$= \text{MULT}_{x,y}^\dagger |q_{A,\mu}\rangle_x |\Gamma_\mu q_{A,\mu}\rangle_y |(q_{A,\mu})^3 \Gamma_\mu\rangle_z \quad (9)$$

$$= |q_{A,\mu}\rangle_x |\Gamma_\mu\rangle_y |(q_{A,\mu})^3 \Gamma_\mu\rangle_z \quad (10)$$

$$(11)$$

This circuit to compute the inner sum in E^Γ could be called $E_i^\Gamma(x,y,z)$. Let us say we are given the following state

$$\bigotimes_A (|q_{A,1}\rangle_a |q_{A,2}\rangle_b) \otimes (|\Gamma_1\rangle_c |\Gamma_2\rangle_d |0\rangle_e)$$

we can now apply $\prod_a E_i^\Gamma(a,c,e) \prod_b E_i^\Gamma(b,d,e)$ to get

$$\bigotimes_A (|q_{A,1}\rangle_a |q_{A,2}\rangle_b) \otimes (|\Gamma_1\rangle_c |\Gamma_2\rangle_d |0\rangle_e) \rightarrow \bigotimes_A (|q_{A,1}\rangle_a |q_{A,2}\rangle_b) \otimes (|\Gamma_1\rangle_c |\Gamma_2\rangle_d |E^\Gamma\rangle_e)$$