

Background, Problem Statement, Objectives and Research Questions

A Network-Informed Bayesian Shrinkage Model for Ordinal Biological Outcomes

Andrews Acheampong

Student, Department of Biochemistry and Biotechnology

Dr. Isabel Mensah

Supervisor, Department of Mathematics

Kwame Nkrumah University of Science and Technology

September 21, 2026

1. Background to the methodological problem.
2. Summary of the gap from the literature review.
3. Evidence-based problem statement.
4. General and specific research objectives.
5. Research questions derived from the objectives.

Background: Ordinal Biological Outcomes

Many biological and clinical outcomes are naturally ordered.

Stage 0 < Stage 1 < Stage 2 < Stage 3 < Stage 4

- The categories have a meaningful order.
- The distance between two neighbouring stages is usually unknown.
- Treating the outcome as continuous may impose an artificial scale.
- Collapsing the stages into two groups may discard useful information.

Bayesian models for ordered response data provide a principled way to model such outcomes and quantify uncertainty (Albert and Chib, 1993; Zhang and Archer, 2021).

Background: High-Dimensional Biological Predictors

Genomic and transcriptomic studies often measure many more predictors than observations.

$$p \gg n$$

- Ordinary regression becomes unstable when the number of predictors is very large.
- Shrinkage methods such as Lasso, Elastic Net and Bayesian Lasso address this problem by controlling the size of regression coefficients (Tibshirani, 1996; Zou and Hastie, 2005; Park and Casella, 2008).
- Biological predictors are often correlated because genes may belong to the same pathway, module or regulatory process.

Background: Why Gene Networks Matter

Biological predictors do not usually act independently.

- Genes may be connected through pathways, co-expression modules or regulatory relationships.
- Network-constrained regression uses these connections to guide variable selection and estimation (Li and Li, 2008; Pan et al., 2010).
- Bayesian network-informed methods extend this idea by placing network structure into the prior distribution of regression coefficients (Cao et al., 2021).

Gap Analysis Summary

The literature review shows that important parts of the problem have been studied separately.

- Shrinkage methods handle high-dimensional predictors, but they may ignore gene-network structure.
- Network-constrained methods use biological networks, but many are developed for continuous or binary outcomes.
- Bayesian ordinal methods handle ordered outcomes, but the reviewed approaches generally do not incorporate biological network information.
- Regularized ordinal regression software exists, but the network-informed Bayesian prior remains underdeveloped for ordinal biological prediction (Wurm et al., 2021; Zhang and Archer, 2021; Cao et al., 2021).

Problem Statement

The gap analysis shows four connected issues.

- High-dimensional biological datasets may contain ordered outcomes and thousands of correlated predictors.
- Shrinkage methods improve estimation and variable selection, but they may ignore biological relationships among genes (Tibshirani, 1996; Zou and Hastie, 2005; Park and Casella, 2008).
- Network-constrained methods incorporate biological graph structure, but the reviewed methods are mainly developed for continuous or binary outcomes (Li and Li, 2008; Pan et al., 2010; Cao et al., 2021).
- Bayesian and regularized ordinal methods exist, but the reviewed ordinal approaches do not fully integrate network-informed priors for prediction and uncertainty quantification (Zhang and Archer, 2021; Wurm et al., 2021).

Therefore, the methodological problem is that existing approaches do not adequately combine the following three requirements in one coherent framework:

1. modelling an ordered biological outcome;
2. shrinking and selecting high-dimensional predictors;
3. using biological network information to guide the prior distribution of gene effects.

To develop and evaluate, a network-informed Bayesian ordinal shrinkage model for high-dimensional biological data with ordered outcomes.


Specific Objectives Following SMART Qualities

1. To formulate a Bayesian cumulative-logit model with a graph-based shrinkage prior that incorporates gene-network information into the regression coefficients.
2. To implement posterior computation for estimating gene effects, credible intervals and posterior probabilities of ordered biological stages.
3. To evaluate the model through simulation under controlled high-dimensional settings, including predictor correlation, network quality and signal strength.
4. To validate the model using an existing biological dataset with an ordered outcome and compare its performance with network-unaware Bayesian ordinal shrinkage and penalized ordinal regression methods.

Research Questions

1. How can gene-network information be incorporated into a Bayesian shrinkage prior for high-dimensional ordinal biological outcomes?
2. How can posterior computation be used to estimate gene effects, quantify uncertainty and predict ordered biological stages?
3. Under simulated high-dimensional settings, does the network-informed Bayesian ordinal model improve prediction and variable selection compared with network-unaware alternatives?
4. When applied to an existing biological dataset, does the proposed model produce accurate, stable and biologically interpretable stage predictions?

Proposed Methodological Workflow



How Existing Data Will Be Used for Validation

The study will first develop and test the method using simulation, where the true signals are known.

An existing biological dataset with an ordered outcome will then be used to assess practical performance.

Validation will focus on whether the proposed model can:

- predict ordered stages accurately;
- produce calibrated posterior stage probabilities;
- identify stable predictors across resampling or cross-validation;
- recover biologically interpretable genes or pathways;
- perform better than or competitively with comparison methods.

References

- Albert, J. H. and Chib, S. (1993). Bayesian analysis of binary and polychotomous response data. *Journal of the American Statistical Association*, 88(422):669–679.
- Cao, M., Fan, Y., and Peng, Q. (2021). Bayesian gene selection based on pathway information and network-constrained regularization. *Computational and Mathematical Methods in Medicine*, 2021:7471516.
- Li, C. and Li, H. (2008). Network-constrained regularization and variable selection for analysis of genomic data. *Bioinformatics*, 24(9):1175–1182.
- Pan, W., Xie, B., and Shen, X. (2010). Incorporating predictor network in penalized regression with application to microarray data. *Biometrics*, 66(2):474–484.

- Park, T. and Casella, G. (2008). The bayesian lasso. *Journal of the American Statistical Association*, 103(482):681–686.
- Tibshirani, R. (1996). Regression shrinkage and selection via the lasso. *Journal of the Royal Statistical Society: Series B*, 58(1):267–288.
- Wurm, M. J., Rathouz, P. J., and Hanlon, B. M. (2021). Regularized ordinal regression and the ordinalnet r package. *Journal of Statistical Software*, 99(6):1–42.
- Zhang, Y. and Archer, K. J. (2021). Bayesian penalized cumulative logit model for high-dimensional data with an ordinal response. *Statistics in Medicine*, 40(6):1453–1481.
- Zou, H. and Hastie, T. (2005). Regularization and variable selection via the elastic net. *Journal of the Royal Statistical Society: Series B*, 67(2):301–320.

Questions and Suggestions