

Introduction

In this research paper, I will be describing how researchers measure "human disturbance" in regard to streamflow variability, the validity of the "human disturbance" index, and current cases of variability having an affect in the Southwest region of the United States. The "human disturbance" index is measurement tool used by the United States Environmental Protection Agency (USEPA) to measure the amount of disturbance present in watershed regions across the nation (Falcone, Carlisle, & Weber, 2010). The reason for needing the index is to use this measurement, along with other variables, is to track the behavior of these hydrological systems. Changes in hydrological systems in the Continental United States (CONUS) have caused some negative outcomes where one being an increase in floods in some regions (Rice, Emanuel, & Vose, 2016). A regional example of a negative outcome from increased streamflow variability is the increasing number of floods in California (Tucker, 2020). The scale of variability is still being studied since previous methods used datasets that contained redundancies, which decreased accuracy but Falcone et al. (2010) described the best approach to accurately classifying watershed disturbance. There is a relationship between streamflow variability and both water resource availability and management (Rice et al., 2016). For present and future studies in relation to hydrological systems, understanding the previously mentioned relationship is crucial to measure human activity and the impact of the side effects from such activity in order to positively manage water resources here in the CONUS and elsewhere in the world.

Data Analysis

To better understand the negative outcomes from increased streamflow variability and increasingly warming climate, the Mallakpour, Sadegh, and AghaKouchak (2018) study reported streamflow minimum and maximum annual streamflows mean ($\text{m}^3 \text{s}^{-1}$) comparing watershed in Northern California.

In 1, notice the minimum streamflow values (red-down arrows) in panels (D-F). In D-panel, there are more reports on negative trends occurring in streamflow values from

1950-2005 which means that inland streams are displaying a decreasing mean during the dry seasons and in G-panel and during the wet season, there is increase in the mean streamflow value. The increase in mean during the wet-season promotes more floods and decrease in mean during dry-season promotes more droughts (Mallakpour et al., 2018). Although the trends are minor from 1950-2005, the cumulative distribution for future highlight a negative trend. *Figure 2* highlights this trend which displays the cumulative distribution of three dataset – (1) 1950-2005 (2) RCP 4.5 2020-2099 (3) RCP 8.5 2020-2099. The left panel is Oroville Lake in California and the right panel is Shasta Lake. Notice the blue line's distribution is greater than the two project trends from RCP 4.5 and RCP 8.5. RCP 4.5 is a model of future streamflow using using a bias-corrected projection and RCP 8.5 uses data derived from global climate model (GCM) to simulate future streamflows. GCM data is known to have systematic error (biases) due to limited GIS spatial resolution which is why one (RCP 4.5) of the two future projections uses a bias correction projection to account for theses errors (biases).

But how does climate change? From Rice et al. (2016) study, atmospheric variables may be potential drivers in streamflow variability. Rice et al. (2016) results suggest that human activities may magnify or amplify the expression of changes. P_{mean} (Precipitation) and DI_{mean} (Dryness index) which were the important atmospheric variables that acted as drivers. There were more variables being used by United State Geological Survey (USGS) but watershed climatology, for instance was too varied amongst CONUS watersheds. *Figure 3* models streamflow trends across the CONUS (Rice et al., 2016).


Relevance

Results

Conclusion

References

- Falcone, J., Carlisle, D., & Weber, L. (2010). Quantifying human disturbance in watersheds: Variable selection and performance of a GIS-based disturbance index for predicting the biological condition of perennial streams. *Ecological Indicators*, *10*(2), 264–273.
- Mallakpour, I., Sadegh, M., & AghaKouchak, A. (2018). A new normal for streamflow in california in a warming climate: Wetter wet seasons and drier dry seasons. *Journal of Hydrology*, *567*, 203–211.
- Rice, J., Emanuel, R., & Vose, J. (2016). The influence of watershed characteristics on spatial patterns of trends in annual scale streamflow variability in the continental u.s. *Journal of Hydrology*, *540*, 250–260.
- Tucker, D. (2020, January 27). More rain and less snow means increased flood risk, stanford study reveals. Retrieved May 20, 2020, from <https://news.stanford.edu/2020/01/27/rain-less-snow-increases-flooding/>

*Figure 1*