

Assignment:- 2

AI1110: Probability and Random Variables

Indian Institute of Technology, Hyderabad


CS22BTECH11001

Aayush Adlakha

29 April, 2023

Exemplar 11.16.3.11 The accompanying Venn diagram shows three events, A, B, and C, and also the probabilities of the various intersections (for instance, $\Pr(AB) = .07$). Determine

- (a) $\Pr(A)$
- (b) $\Pr(BC')$
- (c) $\Pr(A + B)$
- (d) $\Pr(AB')$
- (e) $\Pr(BC)$
- (f) Probability of exactly one of the three occurs.



Solution.

- (a) Clearly,

$$\Pr(A) = 0.13 + 0.07 \quad (1)$$

$$= 0.20 \quad (2)$$

- (b) Clearly,

$$\Pr(B) = 0.10 + 0.07 + 0.15 \quad (3)$$

$$= 0.32 \quad (4)$$

Also,

$$A = A(B + B') = AB + AB' \quad (5)$$

$$[\because B + B' = 1] \quad (6)$$

$$\Pr(A) = \Pr(AB) + \Pr(AB') \quad (7)$$

$$[\because BB' = 0]$$

Using (7)

$$\Pr(BC') = \Pr(B) - \Pr(BC) \quad (8)$$

$$= 0.32 - 0.15 \quad (9)$$

$$= 0.17 \quad (10)$$

- (c) From Axioms of Probability

$$\Pr(A + B) = \Pr(A) + \Pr(B) - \Pr(AB) \quad (11)$$

$$= 0.20 + 0.32 - 0.07 \quad (12)$$

$$= 0.45 \quad (13)$$

- (d) Using (7)

$$\Pr(AB') = \Pr(A) - \Pr(AB) \quad (14)$$

$$= 0.20 - 0.07 \quad (15)$$

$$= 0.13 \quad (16)$$

- (e) Clearly,

$$\Pr(BC) = 0.15 \quad (17)$$

- (f) Let X be the event that exactly one of A , B or C occur.

Let Y be the event that at least one of A , B or C occur.

Using Boolean logic,

$$Y = A + B + C \quad (18)$$

Let Z be the event that at least two of A , B or C occur.

Writing down Z as at least one of AB , BC or AC occurring gives us.

$$Z = AB + BC + CA \quad (19)$$

We know that, all three events never occur simultaneously.

Therefore,

Z represents occurrence of exactly 2 of A , B and C .

X represents occurrence of exactly 1 of A,B and C.

Y can therefore be written as,

$$Y = X + Z \quad (20)$$

$$YZ' = (X + Z)Z' \quad (21)$$

$$YZ' = XZ' + ZZ' \quad (22)$$

$$YZ' = XZ' \quad (23)$$

Z' can be thought of as either none of the events occurring or exactly one of them occurring

XZ' would therefore be event that exactly one occurs.

$$XZ' = X \quad (24)$$

$$X = YZ' \quad (25)$$

$$= (A + B + C)(AB + BC + CA)' \quad (26)$$

$$= (A + B + C)(AB)'(BC)'(CA)' \quad (27)$$

$$= (A + B + C)(A' + B')(B' + C')(A' + C') \quad (28)$$

$$= (AA' + BA' + CA' + AB' + BB' + CB') \quad (29)$$

$$(B'A' + C'A' + B'C' + C'C') \quad (30)$$

$$= (BA' + CA' + AB' + CB')(A'B' + A'C' + B'C') \quad (31)$$

Now, X has been represented as a union of 3 mutually exclusive events.

As any 2 of them has 0 intersection due of presence of complements.

Therefore, by Axioms of Probability

$$\Pr(X) = \Pr(AB'C') + \Pr(A'BC') + \Pr(A'B'C) \quad (32)$$

Clearly, from the figure

$$\Pr(X) = 0.13 + 0.10 + 0.28 \quad (33)$$

$$= 0.51 \quad (34)$$