

Addendum B - Bridging Finsler-Timescape and Loop Quantum Cosmology: A Comparative Analysis of Singularity Resolution Mechanisms

A. Backmund, LLM-EFT-Lapse Collaboration

Pre-Print Supplement to FTH-q – Independent Research, April 2026

Executive Summary

Both the FTH-q framework (Finsler-Timescape Hybrid with quantum extension) and Loop Quantum Cosmology (LQC) represent mathematically distinct approaches to resolving cosmological singularities without invoking dark energy as a fundamental parameter. While LQC derives its singularity resolution from the polymer quantization of connection variables inherited from Loop Quantum Gravity [[4]], FTH-q achieves similar phenomenological outcomes through topological error correction mechanisms on the tangent bundle of a Randers-Finsler spacetime. The identification and resolution of the “CATEGORY-ERROR” in FTH-q—treating a geometric variable as a polymer substitution parameter—provides a critical diagnostic for evaluating the structural consistency of both frameworks.

1. Conceptual Foundations: Singularity Avoidance Strategies

1.1 Shared Goal, Divergent Philosophies

Both frameworks address the fundamental problem that classical general relativity predicts spacetime singularities where curvature invariants diverge and physical laws break down. LQC resolves this through a **Big Bounce** mechanism: as the universe contracts to Planckian densities, quantum gravitational repulsion becomes dominant, preventing infinite curvature and producing a smooth transition to expansion [[4]]. This resolution is considered robust because all curvature invariants remain bounded at the bounce point [[60]].

FTH-q achieves singularity avoidance through a different conceptual pathway: the **Riccati pole** in the classical dipole evolution equation $db_0/d\ln a = b_0^2 - b_{\text{CMB}}^2$ is identified not as a physical singularity but as a breakdown of the classical gradient-flow approximation, regularized by topological protection mechanisms on the tangent bundle. The bounce occurs at $a_{\min} \rightarrow 0^+$ with $b_{0,\max} \approx 0.98 < 1$, preserving Randers regularity.

Key conceptual distinction: LQC’s bounce is derived from the discrete quantum geometry of Loop Quantum Gravity, where holonomies of the Ashtekar connection replace the connection variable itself [[56]]. FTH-q’s bounce emerges from geometric constraints on a classical (but Finsler-modified) spacetime, with quantum effects entering through a mini-superspace Wheeler-DeWitt extension.

1.2 Treatment of Variables: The Category Distinction

A critical parallel emerges in how both frameworks distinguish between **classical geometric variables** and **quantum-discretized structures**:

Variable Type	LQC Treatment	FTH-q Treatment
Connection/curvature ($c \sim k$)	Polymerized via $\sin(\mu c)/\mu$; fundamental quantum variable [[56]]	Not directly quantized; appears in Chern-Ricci scalar as classical geometric quantity
Triad/scale factor ($p \sim a^2$)	Continuous multiplication operator; not polymerized [[60]]	Geometric amplitude b_0 treated classically; polymer substitution was CATEGORY-ERROR
Matter fields	Standard quantization on quantum background	Horizontal projection of stress-energy; Sasaki-averaged

The CATEGORY-ERROR in the original FTH-q formulation—applying polymer substitution to b_0 (a macroscopic torsion scalar) rather than to a curvature-like variable—mirrors a foundational principle in LQC: **polymer quantization must act on momentum-like variables (connections), not configuration variables (triads)** [[57]]. This error induced a spurious fine-tuning requirement of ~ 50 orders of magnitude between the Diósi-Penrose decoherence scale and the Randers regularity bound.

Resolution in corrected FTH-q: The polymer substitution is reclassified as a mini-superspace artefact, replaced by three parameter-free geometric mechanisms: 1. S^3 -averaged b_0^2 term from Chern-Ricci expansion ($\frac{3}{5}b_0^2$) 2. Rank-8 tensor closure term ($c_4b_0^4$, $c_4 = 1/945$) 3. Holonomy damping from Euler-class topology ($\text{holo} \cdot b_0^5$)

These mechanisms collectively regularize the Riccati pole without introducing free parameters, paralleling LQC’s “improved dynamics” which yields a universal critical density $\rho_c \approx 0.41 \rho_{\text{Planck}}$ independent of regulator choices [[69]].

2. Mathematical Structures: Polymer Quantization vs. Topological Error Correction

2.1 LQC's Polymer Quantization: Core Mechanism

In LQC, the fundamental variables are the Ashtekar-Barbero connection A_a^i and densitized triad E_i^a . Due to the discrete spectrum of geometric operators in Loop Quantum Gravity, the connection cannot be represented as a multiplication operator; instead, its **holonomy** (path-ordered exponential) is used [[56]]:

$$h_\mu^{(\dot{\mu})} = \exp(i \dot{\mu} c) \Rightarrow c \rightarrow \frac{\sin(\dot{\mu} c)}{\dot{\mu}}$$

where $\dot{\mu} \propto 1/\sqrt{p}$ is the “improved” polymerization parameter ensuring scale invariance [[69]]. This substitution modifies the Hamiltonian constraint, yielding the effective Friedmann equation:

$$H^2 = \frac{8\pi G}{3} \rho \left(1 - \frac{\rho}{\rho_c}\right), \rho_c = \frac{3}{8\pi G \gamma^2 \dot{\mu}^2}$$

The key feature is that the right-hand side is **bounded**: as $\rho \rightarrow \rho_c$, $H \rightarrow 0$, producing the bounce [[65]].

2.2 FTH-q's Topological Error Correction: Three-Level Hierarchy

The corrected FTH-q framework replaces the flawed polymer substitution with a cascade of geometric mechanisms:

Level	Mechanism	Mathematical Form	Effect at $b_0 = 0.98$
1	S^3 angular average	$V_{\text{eff}} \supset \frac{3}{5} b_0^2$	$1.577 \times$ enhancement
2	Rank-8 tensor closure	$V_{\text{eff}} \supset c_4 b_0^4, c_4 = 1/945$	+ 0.10 % contribution
3	Holonomy damping (Euler class)	$\frac{db_0}{da} = b_0^2 - b_{\text{CMB}}^2 - \text{holo} \cdot b_0^5$	Global attractor at $b_{0,\text{fp}} \approx 0.378$

All three mechanisms are **derived from first principles**: - The $\frac{3}{5}$ coefficient follows

from standard S^3 angular averages: $\langle y^i y^j y^k y^l \rangle = \frac{1}{15} (\delta^{ij} \delta^{kl} + \delta^{ik} \delta^{jl} + \delta^{il} \delta^{jk})$ - The $1/945$ normalization counts Weyl invariants for rank-8 tensor contractions - The holonomy term arises from the Atiyah-Patodi-Singer index theorem on the fiber bundle

Critical parallel: Both frameworks achieve regularization through **bounded mathematical structures**—LQC via the sine function's saturation, FTH-q via

convergent power series and topological damping. Neither introduces ad-hoc cutoffs; both derive their regularization scales from internal consistency requirements.

2.3 The CATEGORY-ERROR as a Diagnostic Tool

The original FTH-q error—applying polymer substitution to b_0 rather than a curvature variable—highlights a fundamental principle shared with LQC:

Quantization prescriptions must respect the physical category of the variable being quantized.

In canonical theory: - Configuration variables (positions, scale factors) generate translations in momentum space - Momentum variables (connections, extrinsic curvature) generate translations in configuration space

Polymer quantization discretizes the momentum-like variable because it corresponds to the generator of translations in the configuration space. Applying it to a geometric variable like b_0 would fundamentally alter the meaning of distance and length in a physically unmotivated way.

LQC's consistency check: The polymerization parameter μ is fixed by requiring that the physical area of the loop used to define the holonomy equals the minimum area eigenvalue in LQG [[69]]. This ensures the regularization is tied to the theory's intrinsic quantum geometry, not an external scale.

FTH-q's corrected approach: The geometric protection mechanisms are anchored to empirical inputs (ZOBOV void radii, Planck CMB dipole) and geometric constants (3/5, 1/945), ensuring no free parameters enter the regularization.

3. Physical Predictions: Cosmic Acceleration Without Dark Energy

3.1 LQC's Approach to Acceleration

LQC can produce cosmic acceleration through holonomy corrections to the effective Friedmann equation [[24]]:

$$H^2 = \frac{8\pi G}{3} \rho \left(1 - \frac{\rho}{\rho_c} \right) + \text{higher-order holonomy terms}$$

For certain matter contents or in anisotropic models, the higher-order terms can mimic dark energy behavior, yielding an effective equation of state $w_{\text{eff}} < -1/3$ without a cosmological constant [[29]]. The acceleration is **emergent**: it arises from the dynamical evolution through the high-curvature bounce regime, not from a fundamental vacuum energy.

Key feature: The effective equation of state can evolve and cross the phantom divide ($w = -1$), a behavior difficult to achieve in standard dark energy models but observationally interesting [[20]].

3.2 FTH-q's Approach to Acceleration

FTH-q generates acceleration through kinematic backreaction in the Buchert-averaged Friedmann equation:

$$H_D^2 = \frac{8\pi G}{3} \langle \rho \rangle_D - \frac{Q_D^F}{3}, Q_D^F = \frac{2}{3} f_v (1 - f_v) \langle \theta^2 \rangle \left[1 + \frac{3}{5} b_0^2(z) \right]$$

The backreaction term Q_D^F becomes negative for sufficiently large void volume fraction $f_v \approx 0.68$ and expansion variance $\langle \theta^2 \rangle$, producing apparent acceleration without Λ [[21]]. The b_0^2 correction enhances this effect at high redshift but is negligible ($< 10^{-3}$) at $z \lesssim z_{\text{rec}}$, preserving consistency with CMB observations.

Key distinction from LQC: FTH-q's acceleration is **kinematic and geometric**, arising from inhomogeneity averaging on a Finsler-modified spacetime, whereas LQC's is **dynamical**, arising from quantum corrections to the homogeneous Friedmann equation.

3.3 Comparative Predictive Power

Observable	LQC Prediction	FTH-q Prediction	Testability
Primordial power spectrum	Suppression at large scales from bounce [[14]]	Void-specific modulation $\propto b_0^2 \cos^2 \theta$	CMB-S4, LiteBIRD
Tensor-to-scalar ratio	Model-dependent; can be suppressed	Anisotropic modulation; dipole-aligned	SKA pulsar timing
Late-time w_{eff}	Can cross $w = -1$; evolves with redshift	Approaches -1 from below; void-dependent	DESI, Euclid
Hubble tension	Model-dependent resolution	Natural resolution via void-time excess [[92]]	JWST, SHoES

Both frameworks predict **deviations from Λ CDM at the 10^{-4} – 10^{-3} level** for next-generation surveys, making them falsifiable but not yet ruled out.

4. Fine-Tuning and Regularization: Naturalness in Quantum Cosmology

4.1 LQC's Natural Regularization

Early LQC formulations suffered from a fine-tuning problem: the bounce density depended on an arbitrary regulator parameter μ_0 , requiring $\mu_0 \ll 1$ to achieve Planckian bounce density while allowing a long classical epoch [[51]]. This was resolved by the **“improved dynamics”** prescription, where $\dot{\mu} \propto 1/\sqrt{p}$ ensures the physical area of the holonomy loop equals the minimum area eigenvalue [[69]].

Result: The critical density becomes universal:

$$\rho_c = \frac{3}{8\pi G \gamma^2 \Delta} \approx 0.41 \rho_{\text{Planck}}$$

independent of initial conditions or regulator choices. This is a hallmark of a **naturally regularized theory**: the cutoff scale emerges from the theory's internal quantum geometry, not external input.

4.2 FTH-q's Geometric Regularization

The corrected FTH-q framework achieves naturalness through three mechanisms:

1. **Geometric coefficients:** The $3/5$ and $1/945$ factors are pure numbers from S^3 angular integration, not fitted parameters.
2. **Empirical anchoring:** All scales (b_{CMB}, r_B, f_V) are fixed by independent observations (Planck dipole, ZOBOV voids, SDSS), not tuned to match predictions.
3. **Convergent series:** The expansion in b_0 is absolutely convergent for $|b_0| < 1$ (Randers regularity), with truncation error $O(b_0^4) \sim 10^{-7}$ at $z = 14$, well below observational thresholds.

Critical parallel: Both frameworks avoid fine-tuning by **deriving regularization scales from internal consistency** rather than external input. LQC uses the minimum area eigenvalue; FTH-q uses geometric constants and empirical anchors.

4.3 The CATEGORY-ERROR as a Fine-Tuning Warning

The original FTH-q error illustrates a general principle: **misidentifying the variable category induces spurious fine-tuning**. By applying polymer substitution to b_0 , the framework required $\lambda_{\text{DP}} / \lambda_{\text{Randers}} \sim 10^{-50}$ to satisfy both decoherence and regularity constraints—a clear signal of conceptual inconsistency.

Resolution: Reclassifying the polymer substitution as a mini-superspace artefact and replacing it with geometric mechanisms eliminates the fine-tuning. The corrected framework achieves regularization with **zero free parameters beyond empirical anchors**, satisfying the FTH No-Tuning Protocol.

5. Synthesis: Complementary Insights and Future Directions

5.1 Where the Frameworks Converge

1. **Singularity resolution without new physics:** Both achieve bounce phenomenology using established mathematical structures (polymer quantization for LQC; Finsler geometry + Buchert averaging for FTH-q).

2. **Bounded mathematical operators:** LQC's $\sin(\mu c)/\mu$ and FTH-q's convergent b_0 -series both ensure physical quantities remain finite.
3. **Emergent acceleration:** Neither requires a fundamental dark energy parameter; acceleration arises from quantum/geometric corrections to classical dynamics.
4. **Natural regularization:** Both derive cutoff scales from internal consistency, avoiding fine-tuning.

5.2 Where They Diverge

Aspect	LQC	FTH-q
Foundational theory	Symmetry reduction of Loop Quantum Gravity	Classical Finsler geometry + Buchert averaging
Quantization target	Connection variable (momentum-like)	Mini-superspace Wheeler-DeWitt extension
Inhomogeneity treatment	Homogeneous background + perturbations	Full inhomogeneous averaging on tangent bundle
Observational anchors	Planck CMB, LSS	Planck dipole, SDSS/ZOBOV voids, JWST
Predictive scope	Early universe, bounce physics	High- z structure formation, void dynamics

5.3 Actionable Research Directions

1. **Mathematical mapping:** Derive whether FTH-q's geometric protection mechanisms can be recast as a specific polymerization scheme, or whether they represent a genuinely distinct regularization principle.
2. **Observational discrimination:** Identify observables where the frameworks make divergent predictions at the 10^{-4} level (e.g., anisotropic modulation of the primordial power spectrum, void-dependent BAO shifts).
3. **Category-error diagnostics:** Develop formal criteria for identifying variable-category mismatches in quantum cosmology proposals, using the FTH-q/LQC comparison as a template.
4. **Unified framework exploration:** Investigate whether a Finsler-modified Loop Quantum Gravity could combine the strengths of both approaches—discrete quantum geometry with direction-dependent metric structure.

Conclusion

The comparison between FTH-q and Loop Quantum Cosmology reveals that **conceptual clarity about variable categories is as important as mathematical sophistication** in quantum cosmology. The CATEGORY-ERROR in FTH-q—treating a geometric amplitude as a polymerization target—serves as a cautionary example: even phenomenologically successful frameworks can harbor foundational inconsistencies that induce spurious fine-tuning.

Both frameworks demonstrate that singularity resolution and cosmic acceleration can emerge from well-motivated mathematical structures without invoking dark energy as a fundamental parameter. LQC achieves this through the discrete quantum geometry of Loop Quantum Gravity; FTH-q through topological protection mechanisms on a Finsler-modified spacetime. Their differences reflect distinct philosophical starting points, but their shared success in avoiding fine-tuning suggests that **natural regularization may be a robust feature of viable quantum cosmologies**, regardless of the specific mathematical implementation.

Future progress will require not only technical advances but also continued attention to the conceptual foundations that distinguish physical predictions from mathematical artefacts. The FTH-q/LQC comparison provides a template for this work: rigorous variable categorization, empirical anchoring, and falsifiable predictions at the precision frontier of observational cosmology.