

N-GRAMS AND SMOOTHING

Manish Shrivastava

MAXIMUM LIKELIHOOD ESTIMATION

- ▶ $P_{MLE}(w_n | w_1 \dots w_{n-1}) = \frac{C(w_1 \dots w_n)}{C(w_1 \dots w_{n-1})}$
- ▶ Estimate sequence probabilities using “counts” or frequencies of sequences
- ▶ Problems
 - ▶ Sparseness
 - ▶ What do you do when unknown words are seen??

EXAMPLE

- Data
 - The boy ate a chocolate
 - The girl bought a chocolate
 - The girl then ate a chocolate
 - The horse bought a boy
- The boy bought a chocolate
 - Unigram Probabilities
 - $(4/16)*(2/16)*(2/16)*(4/16)*(3/16)$
 - $(4*2*2*4*3)/21^5 = 0.000047$
 - Bi-gram Probabilities
 - <The boy> <boy bought> <bought a> <a chocolate>
 - $(2/4)*(0/4)*(2/2)*(3/4) = \underline{0}$

SMOOTHING

- Higher order n-grams perform well but suffer from data sparsity
- Lower order n-grams are not reliable
- Standard MLE does not work for unseen data

SMOOTHING

- Laplace smoothing (add-one)
- Lidstone's law
- Held-out Estimation
- Good Turing Estimation

LAPLACE SMOOTHING

- The oldest smoothing method available
- Each unseen n-gram is given a very low estimate
- $P_{Lap}(w_1 \dots w_{n-1} w_n) = \frac{C(w_1 \dots w_n) + 1}{N + B}$
- **Probability** $1/(N+B)$ or **Frequency (estimated)** $N/(N+B)$
- **N** is the number of seen n-grams and **B** is the number of possible n-grams

LIDSTONE'S LAW

- The probability mass assigned to unseen events is too high with Laplace smoothing
- Solution : assign smaller addition
 - $P_{Lid}(w_1 \dots w_{n-1} w_n) = \frac{C(w_1 \dots w_n) + \lambda}{N + B\lambda}$
- Jeffereys-Perks law
 - Set λ as $\frac{1}{2}$
 - Expectation of λ that maximizes above equation

HELD OUT ESTIMATION

- So far, we have not considered how the learned estimates behave in real life
- It is possible that the learnt values are just a property of the training data
 - Imagine training from Chemical domain and applying in Physics domain
- Also, adding a random value is not very sound
 - At higher total count the probability of something that occurred once and something that never occurred would be very close
 - Adding small value also increases the total denominator.
 - $B\lambda$ might become a very large part of $(N + B\lambda)$
 - In other words, a large probability mass might be assigned to unseen events

HELD-OUT ESTIMATES

- Treat part of the data as “real life” data
- Intuition :
 - Estimates of bigrams with similar frequencies, would be similar
 - Example: if <a boy> occurs 3 times and <a girl> occurs 3 times, instead of treating them as different, we can consider them to belong to one class or bin ;
 - Bigrams with frequency 3
 - Now we can calculate the average probability for all bigrams with frequency 3

HELD-OUT ESTIMATION

- Method:
 - Partition data into two parts
 - Training and Held-Out
 - N_r = Number of n-grams with frequency 'r' in Training data
 - T_r = sum of number of times all the n-grams with frequency r, as identified in the training data, appear in the Held-Out data

$$T_r = \sum_{\{w_1 \dots w_n : C_{ho}(w_1 \dots w_n) = r\}} C_{ho}(w_1 \dots w_n)$$

- Then $\{w_1 \dots w_n : C_{tr}(w_1 \dots w_n) = r\}$

$$P_{ho}(w_1 \dots w_n) = \frac{T_r}{N_r N}$$

HELD OUT ESTIMATION

Freq	Training Data (number of words with given frequency)	Held Out Data (Sum of Frequencies of words with r freq in Training)
1	N1	$T1 = (Nr * r^*)$
2	N2	T2
3	N3	T3
4	N4	T4
.		
.		
r	Nr (average frequency of these words = $Nr * r / Nr = r$)	Tr (average frequency of these words = $Tr / Nr = r^*$)
.		
.		

HELD-OUT ESTIMATION

- Probability is calculated keeping both training and held-out data in mind.
 - Is it the best estimate?
 - In this scenario?
 - If the role of training and held-out data is reversed ?
 - $P_{Ho}(w_1 \dots w_n) = \frac{T_r^{01}}{N_r^0 N}$
 - $P_{Ho}(w_1 \dots w_n) = \frac{T_r^{01}}{N_r^0 N}$

DELETED ESTIMATION (CROSS-ESTIMATION)

- Jelinek and Mercer 1985
 - $P_{Del}(w_1 \dots w_n) = \frac{T_r^{01} + T_r^{10}}{(N_r^0 + N_r^1)N}$
 - Performs really well
 - Still a way off for low frequency events

GOOD TURING SMOOTHING

- Word classes with similar frequency counts can be treated similarly
 - Or can help each other
- A word which is at frequency 1 is a rare word, it is just by chance that you have been able to see it (data property)
- We should be able to compute (approximately) probabilities of lower frequency words with the help of higher frequency words

$$r^* = (r + 1) \frac{E(N_r + 1)}{E(N_r)}$$

- Here r^* is the adjusted frequency
- The probability then is r^*/N

GOOD-TURING SMOOTHING

- Use Number of times n-grams of frequency r have occurred
 - To fit for both high frequency and low frequency terms, use a curve fitting function
- $P_{GT}(w_1 \dots w_n) = \frac{r^*}{N}$
 - Where $r^* = \frac{(r+1)S(r+1)}{S(r)}$
- For unseen events,
 - $P_{GT}(w_1 \dots w_n) = \frac{N_1}{N * N_0}$
- Simple Good Turing
 - $S() =$ Power Curve for high frequency terms
 - $N_r = ab^r$

EXAMPLE

- Consider the data
 - I like strong coffee
 - I bought some coffee
 - I drank black coffee
 - Good coffee is strong coffee
- Then the test sentence
 - I like black strong coffee
- The trigram <black strong coffee> is never seen before and will be assigned some small probability
 - But bigram <strong coffee> is seen and is very informative

BACK OFF

- Back off is “using lower order N-grams for estimating higher orders”
- Back-Off
 - Linear Interpolation
 - Katz Backoff
 - General linear Interpolation

BACK OFF

- So far, all we have done is get estimates for various n-grams
- Problems (Same old),
 - For n-grams with low frequency, estimates are not really good
- Solution,
 - If n-gram does not work, fall back to (n-1)-gram
 - Better still, combine estimators for n-gram, (n-1)-gram, (n-2)-gram ... unigram

KATZ BACK-OFF

- $P_{li}(w_n | w_1 \dots w_{n-1}) = \alpha * \frac{C(w_1 \dots w_n)}{C(w_1 \dots w_{n-1})}$ if Counts are greater than a certain k
- Else
- $P_{li}(w_n | w_1 \dots w_{n-1}) = (1 - \alpha) * P_{li}(w_n | w_2 \dots w_{n-1})$

LINEAR INTERPOLATION

- $P_{li}(w_n | w_1 \dots w_{n-1}) = \lambda_1 P(w_n) + \lambda_2 P_{li}(w_n | w_{n-1}) + \lambda_3 P_{li}(w_n | w_{n-1}, w_{n-2})$
- Works really well
- Reserve Probability mass for unseen items
- λ_i s need to be learned separately

GENERAL LINEAR INTERPOLATION

- *Why just back off to immediate lower order?*
- This method allows random back off schemes moderated by the weights λ_i s
- One might back off to any of the lower orders or to any other estimator as chosen by the designer
- $P_{li}(w_n|h) = \sum_{i=1}^k \lambda_i P_i(w_n|h)$

WITTEN BELL SMOOTHING

$$P_{wb}(w_i \mid w_{i-n+1} \dots w_{i-1}) = \lambda_{w_{i-n+1} \dots w_{i-1}} P_{wb}(w_i \mid w_{i-n+1} \dots w_{i-1}) \\ + (1 - \lambda_{w_{i-n+1} \dots w_{i-1}}) P_{wb}(w_i \mid w_{i-n+2} \dots w_{i-1})$$

Where,

$$(1 - \lambda_{w_{i-n+1} \dots w_{i-1}}) = \frac{|\{w_i \mid C(w_{i-n+1} \dots w_i) > 0\}|}{|\{w_i \mid C(w_{i-n+1} \dots w_i) > 0\}| + \sum_{w_i} C(w_{i-n+1} \dots w_i)}$$

SAMPLE DATA

- The boy ate a chocolate
 - The girl bought a chocolate
 - The girl then ate a chocolate
 - The boy bought a horse
 - The boy has a horse
 - The boy stole a chocolate
-
- Compute likelihood of
 - The girl stole a horse

QUESTIONS?