

Universidad del Istmo de Guatemala Facultad de Ingenieria Ing. en Sistemas Informatica 1

Prof. Ernesto Rodriguez - erodriguez@unis.edu.gt

# Hoja de trabajo #3

Samantha Rodas Chaluleu Adalí Garrán Jiménez 13 de Agosto, 2019

# Ejercicio #1

Utilizando la definicion de suma  $(\oplus)$  para los numeros naturales unarios, llevar a cabo la suma entre tres [s(s(s(0)))] y cuatro [s(s(s(s(0))))].

|  | $s(s(s(0))) \oplus s(s(s(s(0))))$      |
|--|--|
| a=s(x)<br>x=s(s(0))<br>b=s(s(s(s(0)))) | $s[s(s(0)) \oplus b]$                  |
| a=s(x)<br>x=s(0)<br>b=s(s(s(s(0))))    | $s[s(s(0)\oplus b)]$                   |
| a=s(x)<br>x=0<br>b=s(s(s(s(0))))       | $s[s(s(0\oplus b))]$                   |
| 0+b=b                                  | s[(s(s(b))] $[s(s(s(s(s(s(s(0)))))))]$ |
|  | = 7                                    |

#### Ejercicio #2

Definir inductivamente una función para multiplicar  $(\otimes)$  numeros naturales unarios.

$$a \otimes b := \begin{cases} 0 & \text{si } a = o \\ 0 & \text{si } b = o \\ a & \text{si } b = 1 \\ b & \text{si } a = 1 \\ b \oplus (x \otimes b) & \text{si } a = \sigma(x) \end{cases}$$

### Ejercicio #3

Verifique que su definición de multiplicación es correcta multiplicando los siguientes valores:

•  $\sigma(\sigma(\sigma(0))) \otimes 0$  $\sigma(\sigma(\sigma(0))) \otimes 0$  $a = \sigma(\sigma(\sigma(0)))$  $a \otimes 0 = 0$ •  $\sigma(\sigma(\sigma(0))) \otimes \sigma(0)$  $\sigma(\sigma(\sigma(0))) \otimes \sigma(0)$  $a = \sigma(x)$  $x = \sigma(\sigma(0))$  $b = \sigma(0)$  $b \oplus (x \otimes b)$  $= b \oplus (\sigma(\sigma(0)) \otimes b)$  $a = \sigma(x)$  $x = \sigma(0)$  $= b \oplus b \oplus (\sigma(0) \otimes b)$  $a = \sigma(x)$ x=0 $= b \oplus b \oplus b \oplus (0 \otimes b)$  $0 \otimes b=0$  $= b \oplus b \oplus b \oplus (0)$  $=b\oplus b\oplus b$  $= \sigma(0) \oplus \sigma(0) \oplus \sigma(0)$  $=\sigma(\sigma(\sigma(0)))$ =3•  $\sigma(\sigma(\sigma(0))) \otimes \sigma(\sigma(0))$  $\sigma(\sigma(\sigma(0))) \otimes \sigma(\sigma(0))$  $a = \sigma(x)$  $x = \sigma(\sigma(0))$  $b = \sigma \sigma((0))$ 

 $b \oplus (x \otimes b)$ 

$$= b \oplus (\sigma(\sigma(0)) \otimes b)$$

$$= \sigma(x)$$

$$x = \sigma(0)$$

$$= b \oplus b \oplus (\sigma(0) \otimes b)$$

$$= b \oplus b \oplus b \oplus (0 \otimes b)$$

$$= b \oplus b \oplus b \oplus (0)$$

$$= b \oplus b \oplus b \oplus (0)$$

$$= b \oplus b \oplus b$$

$$= \sigma(\sigma(0)) \oplus \sigma(\sigma(0)) \oplus \sigma(\sigma(0))$$

$$= \sigma(\sigma(\sigma(\sigma(\sigma(\sigma(0))))))$$

$$= 6$$

## Ejercicio #4

Demostrar utilizando inducción:

• 
$$a \oplus \sigma(\sigma(0)) = \sigma(\sigma(a))$$
  
Caso Base:  $a=0$ 

$$0 + \sigma(\sigma(0)) = \sigma(\sigma(0))$$
$$\sigma(\sigma(0)) = \sigma(\sigma(0))$$

Caso Inductivo: 
$$a=\sigma(x)$$

Hipótesis Inductiva: 
$$x+\sigma(\sigma(0)) = \sigma(\sigma(x))$$

$$\sigma(x + \sigma(\sigma(0))) = \sigma(\sigma(\sigma(0)))$$

$$\sigma(\sigma(\sigma(0)) + x) = \sigma(\sigma(\sigma(0)))$$

$$\sigma(\sigma(\sigma(0) + x)) = \sigma(\sigma(\sigma(0)))$$

$$\sigma(\sigma(\sigma(0 + x))) = \sigma(\sigma(\sigma(0)))$$

$$\sigma(\sigma(\sigma(x))) = \sigma(\sigma(\sigma(0)))$$

 $\sigma(x) + \sigma(\sigma(0)) = \sigma(\sigma(\sigma(0)))$ 

• 
$$a \otimes b = b \otimes a$$
  
Caso Base:  $a=0$ 

$$0 \otimes b = b \otimes 0$$
$$0 = 0$$

Caso Inductivo:  $a=\sigma(x)$ 

Hipótesis Inductiva:  $x \otimes b = b \otimes x$ 

$$\sigma(x) \otimes b = b \otimes \sigma(x)$$

$$b = \sigma(i)$$

$$b \oplus (x \otimes b) = b \otimes \sigma(x)$$

$$\sigma(i) \oplus (x \otimes \sigma(i)) = \sigma(i) \otimes \sigma(x)$$

$$\sigma(i) \oplus (x \otimes \sigma(i)) = \sigma(x) \oplus (i \otimes \sigma(x))$$

$$\sigma(i \oplus (x \otimes \sigma(i))) = \sigma(x \oplus (i \otimes \sigma(x)))$$

$$\sigma(i \oplus (x \otimes b)) = \sigma(x \oplus (i \otimes a))$$

Por Hipótesis inductiva:

$$x \otimes b = b \otimes x$$

$$i \otimes a = a \otimes i$$

$$\sigma(i \oplus (b \otimes x)) = \sigma(x \oplus (a \otimes i))$$

$$\sigma(i \oplus (\sigma(i) \otimes x)) = \sigma(x \oplus (\sigma(x) \otimes i))$$

$$\sigma(i \oplus x \oplus (i \otimes x)) = \sigma(x \oplus i \oplus (x \otimes i))$$

$$\sigma(x \oplus i \oplus (i \otimes x)) = \sigma(x \oplus i \oplus (x \otimes i))$$

$$x \otimes i = i \otimes x$$

$$\sigma(x \oplus i \oplus (x \otimes i)) = \sigma(x \oplus i \oplus (x \otimes i))$$

$$\mathbf{k} = x \oplus i \oplus (x \otimes i)$$

$$\sigma(k) = \sigma(k)$$

• 
$$a \otimes (b \otimes c) = (a \otimes b) \otimes c$$

Caso Base: a=0

$$0 \otimes (b \otimes c) = (0 \otimes b) \otimes c$$
$$0 \otimes (b \otimes c) = 0 \otimes c$$
$$0 = 0$$

Caso Inductivo:  $a=\sigma(x)$ 

Hipótesis Inductiva:  $x \otimes (b \otimes c) = (x \otimes b) \otimes c$ 

$$\sigma(x) \otimes (b \otimes c) = (\sigma(x) \otimes b) \otimes c$$
$$(b \otimes c) \oplus (x \otimes (b \otimes c)) = (\sigma(x) \otimes b) \otimes c$$
$$(b \otimes c) \oplus (x \otimes (b \otimes c)) = (b \oplus (x \otimes b)) \otimes c$$

Distribuir la multiplicación  $b \oplus (x \otimes b) \otimes c$  (ver siguiente demostración  $(a \oplus b) \otimes c = (a \otimes c) \oplus (b \otimes c)$ ):

$$(b \otimes c) \oplus (x \otimes (b \otimes c)) = (b \otimes c) \oplus ((x \otimes b) \otimes c)$$

$$(b \otimes c) \oplus [(x \otimes (b \otimes c)) = ((x \otimes b) \otimes c)] \oplus (b \otimes c)$$

Asumiendo que  $[(x \otimes (b \otimes c)) = ((x \otimes b) \otimes c)]$  es verdadero (Hipótesis Inductiva):

$$(b \otimes c) = (b \otimes c)$$

•  $(a \oplus b) \otimes c = (a \otimes c) \oplus (b \otimes c)$ 

Caso Base: a=0

$$(0 \oplus b) \otimes c = (0 \otimes c) \oplus (b \otimes c)$$

$$(b)\otimes c=(0)\oplus (b\otimes c)$$

$$b \otimes c = b \otimes c$$

Caso Inductivo:  $a=\sigma(x)$ 

Hipótesis Inductiva:  $(x \oplus b) \otimes c = (x \otimes c) \oplus (b \otimes c)$ 

$$(\sigma(x) \oplus b) \otimes c = (\sigma(x) \otimes c) \oplus (b \otimes c)$$

$$\sigma(x \oplus b) \otimes c = (\sigma(x) \otimes c) \oplus (b \otimes c)$$

$$c \oplus ((x \oplus b) \otimes c) = (\sigma(x) \otimes c) \oplus (b \otimes c)$$

$$c \oplus ((x \oplus b) \otimes c) = (c \oplus (x \otimes c)) \oplus (b \otimes c)$$

$$c \oplus [(x \oplus b) \otimes c) = (x \otimes c)) \oplus (b \otimes c)] \oplus c$$

Asumiendo que  $[(x \oplus b) \otimes c) = (x \otimes c) \oplus (b \otimes c)$  es verdadero (Hipótesis Inductiva):

$$c = c$$