

MAT185 Linear Algebra Assignment 2

Instructions:

Please read the **MAT185 Assignment Policies & FAQ** document for details on submission policies, collaboration rules and academic integrity, and general instructions.

1. **Submissions are only accepted by Gradescope.** Do not send anything by email. Late submissions are not accepted under any circumstance. Remember you can resubmit anytime before the deadline.
2. **Submit solutions using only this template pdf.** Your submission should be a single pdf with your full written solutions for each question. If your solution is not written using this template pdf (scanned print or digital) then your submission will not be assessed. Organize your work neatly in the space provided. Do not submit rough work.
3. **Show your work and justify your steps** on every question but do not include extraneous information. Put your final answer in the box provided, if necessary. We recommend you write draft solutions on separate pages and afterwards write your polished solutions here on this template.
4. **You must fill out and sign the academic integrity statement below;** otherwise, you will receive zero for this assignment.

Academic Integrity Statement:

Full Name: _____

Student number: _____

Full Name: _____

Student number: _____

I confirm that:

- I have read and followed the policies described in the document **MAT185 Assignment Policies & FAQ**.
- In particular, I have read and understand the rules for collaboration, and permitted resources on assignments as described in subsection II of the the aforementioned document. I have not violated these rules while completing and writing this assignment.
- I understand the consequences of violating the University's academic integrity policies as outlined in the [Code of Behaviour on Academic Matters](#). I have not violated them while completing and writing this assignment.

By signing this document, I agree that the statements above are true.

Signatures: 1) _____

2) _____

Preamble: An application of linear algebra to calculus.

Recall the technique of partial fractions decomposition to evaluate the integral of rational functions. For example, suppose we would like to evaluate the integral

$$\int \frac{7x^2 + 7}{(x^2 + 3)(x - 2)} dx$$

We look for scalars a, b , and c such that

$$\frac{7x^2 + 7}{(x^2 + 3)(x - 2)} = \frac{ax + b}{x^2 + 3} + \frac{c}{x - 2}$$

After some algebra, we find that $a = 2$, $b = 4$, and $c = 5$, and therefore,

$$\frac{7x^2 + 7}{(x^2 + 3)(x - 2)} = \frac{2x + 4}{x^2 + 3} + \frac{5}{x - 2}$$

Then,

$$\begin{aligned} \int \frac{7x^2 + 7}{(x^2 + 3)(x - 2)} dx &= \int \frac{2x + 4}{x^2 + 3} dx + \int \frac{5}{x - 2} dx \\ &= \ln(x^2 + 3) + \frac{4}{\sqrt{3}} \arctan\left(\frac{x}{\sqrt{3}}\right) + 5 \ln(x - 2) + C \end{aligned}$$

where C is a constant.

In Question 1, we will use the theory of basis and dimension in linear algebra to explain why the partial fractions decomposition

$$\frac{7x^2 + 7}{(x^2 + 3)(x - 2)} = \frac{ax + b}{x^2 + 3} + \frac{c}{x - 2}$$

exists, thereby allowing us to solve the integral.

1. Let

$$V = \left\{ \frac{dx^2 + ex + f}{(x^2 + 3)(x - 2)} \mid d, e, f \in \mathbb{R} \right\}$$

We define vector addition and scalar multiplication in V by the usual function addition and scalar multiplication. Then V is vector space.

(a) Prove that $\dim V = 3$. Then, explain why a partial fractions decomposition of the form

$$\frac{dx^2 + ex + f}{(x^2 + 3)(x - 2)} = \frac{ax + b}{x^2 + 3} + \frac{c}{x - 2}$$

is consistent with the dimension of V .

Use the page 3 to answer this question.

1(a)

By definition of dimension ($\dim V = 3$), there are three vectors in any of V 's bases. To prove this we will show there exists three linearly independent vectors that span V .

Assume these three vectors are $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3 \in V$:

$$\mathbf{v}_1 = \frac{x^2}{(x^2+3)(x-2)}, \mathbf{v}_2 = \frac{x}{(x^2+3)(x-2)}, \mathbf{v}_3 = \frac{1}{(x^2+3)(x-2)},$$

i. To show linear independence we will use its definition:

$$\begin{aligned} \lambda_1 \mathbf{v}_1 + \lambda_2 \mathbf{v}_2 + \lambda_3 \mathbf{v}_3 &= \mathbf{0} \\ \lambda_1 \frac{x^2}{(x^2+3)(x-2)} + \lambda_2 \frac{x}{(x^2+3)(x-2)} + \lambda_3 \frac{1}{(x^2+3)(x-2)} &= \mathbf{0} \end{aligned}$$

Multiplying both sides by the denominator:

$$\lambda_1 x^2 + \lambda_2 x + \lambda_3 = \mathbf{0}$$

???The only value satisfying this equation is $\lambda_1 = \lambda_2 = \lambda_3 = 0$ Therefore, $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$ are linearly independent.

ii. Proving span

$$\begin{aligned} \text{span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\} &= \{\mathbf{v} | \mathbf{v} = \sum_{i=1}^n \lambda_i \mathbf{v}_i, \lambda_i \in \mathbb{R}\} \\ \lambda_1 \mathbf{v}_1 + \lambda_2 \mathbf{v}_2 + \lambda_3 \mathbf{v}_3 &= \mathbf{0} \\ &= \lambda_1 \frac{x^2}{(x^2+3)(x-2)} + \lambda_2 \frac{x}{(x^2+3)(x-2)} + \lambda_3 \frac{1}{(x^2+3)(x-2)} = \mathbf{0} \\ &= \frac{\lambda_1 x^2 + \lambda_2 x + \lambda_3}{(x^2+3)(x-2)} \end{aligned}$$

By choosing $\lambda_1 = d, \lambda_2 = e, \lambda_3 = f$, we see any vector $\frac{dx^2+ex+f}{(x^2+3)(x-2)}$ can be formed. Therefore $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ spans V .

iii. Since $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ span V and are linearly independent, no vectors need to be removed or added to the set. By Proof of Construction, a basis has been formed by $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ where $\dim\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\} = 3$ since there are three linearly independent vectors.

Showing $\frac{ax+b}{x^2+3} + \frac{c}{x-2} | a, b, c \in \mathbb{R}$ also has dimension of 3:

$$\begin{aligned} \frac{ax+b}{x^2+3} + \frac{c}{x-2} &= \frac{(ax+b)(x-2)}{(x^2+3)(x-2)} + \frac{c(x^2+3)}{(x^2+3)(x-2)} \\ &= \frac{(c+a)x^2}{(x^2+3)(x-2)} + \frac{(b-2a)x}{(x^2+3)(x-2)} + \frac{(3c-2b)}{(x^2+3)(x-2)} \end{aligned}$$

This is the form of a linear combination of three vectors. Where $c+a = d, b-2a = e, 3c-2b = f$, the linear combination spans V . To show linear independence, we can multiply by the common denominator and represent the coefficient equations as a matrix.

$$\begin{aligned} \mathbf{Ax} &= \mathbf{b} \\ \begin{bmatrix} 1 & 0 & 1 \\ -2 & 1 & 0 \\ 0 & -2 & 3 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} &= \begin{bmatrix} d \\ e \\ f \end{bmatrix} \end{aligned}$$

Matrix \mathbf{A} has 3 linearly independent columns, and therefore a dimension of 3. As a result, the partial fraction decomposition forms three linearly independent vectors, which agree with the $\dim V = 3$.

1. Let

$$V = \left\{ \frac{dx^2 + ex + f}{(x^2 + 3)(x - 2)} \mid d, e, f \in \mathbb{R} \right\}$$

We define vector addition and scalar multiplication in V by the usual function addition and scalar multiplication. Then V is vector space.

(b) Using that $\dim V = 3$ from part (a), explain why we do not expect a partial fractions decomposition of the form

$$\frac{dx^2 + ex + f}{(x^2 + 3)(x - 2)} = \frac{a}{x^2 + 3} + \frac{b}{x - 2}$$

to exist.

By proving $\dim \frac{a}{x^2 + 3} + \frac{b}{x - 2} < 3$, we know a solution of this form can not span V since it can not have a smaller dimension than the basis.

$$\begin{aligned} \frac{a}{x^2 + 3} + \frac{b}{x - 2} &= \frac{a(x - 2)}{(x^2 + 3)(x - 2)} + \frac{b(x^2 + 3)}{(x - 2)(x^2 + 3)} \\ &= \frac{ax - 2a}{(x^2 + 3)(x - 2)} + \frac{bx^2 + 3b}{(x - 2)(x^2 + 3)} \\ &= \frac{ax - 2a + bx^2 + 3b}{(x^2 + 3)(x - 2)} \\ &= \frac{bx^2}{(x^2 + 3)(x - 2)} + \frac{ax}{(x^2 + 3)(x - 2)} + \frac{-2a + 3b}{(x^2 + 3)(x - 2)} \end{aligned}$$

Comparing to the form of V and multiplying out the denominator:

$$\begin{aligned} \frac{dx^2}{(x^2 + 3)(x - 2)} + \frac{ex}{(x^2 + 3)(x - 2)} + \frac{f}{(x^2 + 3)(x - 2)} &= \frac{bx^2}{(x^2 + 3)(x - 2)} + \frac{ax}{(x^2 + 3)(x - 2)} + \frac{-2a + 3b}{(x^2 + 3)(x - 2)} \\ dx^2 + ex + f &= bx^2 + ax + (-2a + 3b) \end{aligned}$$

To span V , we must show $b = d$, $a = e$, $-2a + 3b = f$. Representing these equations in Matrix form:

$$\begin{aligned} \mathbf{Ax} &= \mathbf{b} \\ \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ -2 & 3 & 0 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} &= \begin{bmatrix} d \\ e \\ f \end{bmatrix} \end{aligned}$$

Matrix \mathbf{A} has a linearly dependent column of 0s, and therefore $\dim A < 3$. This means the linear combination of the partial fraction decomposition also has a dimension less than 3.

$$\dim\left(\left\{\frac{a}{x^2 + 3}, \frac{b}{x - 2}\right\}\right) < 3$$

Therefore the partial fraction decomposition can not span V and the equality does not hold. This is numerical seen as any case where $f \neq 3b - 2a$, and there will be no solution to the $\mathbf{Ax} = \mathbf{b}$ system.

2. Suppose that W_1 and W_2 are both three dimensional subspaces of \mathbb{R}^4 . In this question, we will show that $W_1 \cap W_2$ contains a plane.

Let $\mathbf{w}_1, \mathbf{w}_2, \mathbf{w}_3$ be a basis for W_1 , and let $\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3$ be a basis for W_2 .

(a) If $\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3$ all belong to W_1 explain why $W_1 \cap W_2$ contains a plane.

Given u_1, u_2, u_3 form a basis for W_2 , they must be linearly independent and span W_2 by definition. Given $u_1, u_2, u_3 \in W_1$, $W_2 \subset W_1$ since any vector $\mathbf{x} \in W_2$ and $\mathbf{x} \in \text{span}\{u_1, u_2, u_3\}$ by closure under vector addition and scalar multiplication. If $W_2 \subset W_1$, it follows $W_1 \cap W_2 = W_2$ and therefore $u_1, u_2, u_3 \in W_1 \cap W_2$.

The dimension of plane is 2, as it is a surface spanned by two linearly independent vectors. In other words, a plane's bases are formed by two linearly independent vectors.

Take two linearly independent vectors $u_1, u_2 \in W_1 \cap W_2$, which are also linearly independent being a basis for W_2 . $\text{span}\{u_1, u_2\}$ forms a plane because it is all linear combinations of two linearly independent vectors and $\text{span}\{u_1, u_2\} \subset W_1 \cap W_2$ as previously stated.

(b) Now suppose that not all of $\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3$ belong to W_1 . Say $\mathbf{u}_1 \notin W_1$. Prove that $\mathbf{w}_1, \mathbf{w}_2, \mathbf{w}_3, \mathbf{u}_1$ is a basis for \mathbb{R}^4 . Since w_1, w_2, w_3 form a basis for W_1 , every vector $w \in W$ can be expressed by a linear combination:

$$w = \lambda_1 w_1 + \lambda_2 w_2 + \lambda_3 w_3, \lambda_1, \lambda_2, \lambda_3 \in \mathbb{R}$$

Assume u_1 can be formed by a linear combination of w_1, w_2, w_3 .

$$u = \lambda_1 w_1 + \lambda_2 w_2 + \lambda_3 w_3$$

Then, $u_1 \in W_1$ since all linear combinations of the basis vectors must be in W . Given $u_1 \notin W_1$, by contradiction, u_1 can not be formed by a linear combination of w_1, w_2, w_3 or $u_1 \notin \text{span}\{w_1, w_2, w_3\}$.

Consider the following linear combination where $\alpha_1, \lambda_1, \lambda_2, \lambda_3 \in \mathbb{R}$ and we assume $\alpha \neq 0$:

$$\begin{aligned} \alpha_1 u_1 + \lambda_1 w_1 + \lambda_2 w_2 + \lambda_3 w_3 &= 0 \\ \lambda_1 w_1 + \lambda_2 w_2 + \lambda_3 w_3 &= -\alpha_1 u_1 \\ -\frac{\lambda_1}{\alpha_1} w_1 - \frac{\lambda_2}{\alpha_1} w_2 - \frac{\lambda_3}{\alpha_1} w_3 &= u_1 \end{aligned}$$

Since $-\frac{\lambda_x}{\alpha_1} \in \mathbb{R}$, this equation is a linear combination of w_1, w_2, w_3 that forms u_1 . By the previous contradiction, this is not possible, and therefore, $\alpha = 0$. Additionally, $\lambda_1 = \lambda_2 = \lambda_3 = 0$ since they are linearly independent by the definition of a basis.

Therefore, u_1, w_1, w_2, w_3 are four linearly independent vectors. Since $\dim \mathbb{R}^4 = 4$, there are four linearly independent vectors in the basis of \mathbb{R}^4 . By the Algebra Triangle Theorem, the set $\{u_1, w_1, w_2, w_3\}$ must be a basis for W_1 since the four vectors are linearly independent.

2. Suppose that W_1 and W_2 are both three dimensional subspaces of \mathbb{R}^4 . In this question, you will show that $W_1 \cap W_2$ contains a plane.

Let $\mathbf{w}_1, \mathbf{w}_2, \mathbf{w}_3$ be a basis for W_1 , and let $\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3$ be a basis for W_2 .

(c) Using the assumption and conclusion from part (b), find two vectors in $W_1 \cap W_2$ and then prove that these two vectors span a plane.

From (b), there can only be 4 linearly independent vectors in \mathbb{R}^4 . Since $W_1 \subset \mathbb{R}^4$ and $W_2 \subset \mathbb{R}^4$, only 4 vectors of $w_1, w_2, w_3, u_1, u_2, u_3$ can be linearly independent.

We know w_1, w_2, w_3, u_1 are linearly independent from (b). Therefore, u_2 and u_3 must be linearly dependent to two different vectors from w_1, w_2, w_3, u_1 since u_2 and u_3 are linearly independent themselves. We also know u_2 and u_3 can not be dependent to u_1 since by the definition of a basis, u_1, u_2, u_3 are linearly independent. Furthermore, $u_1 \notin W_1 \cap W_2$ given $u_1 \notin W_1$.

Let's pick w_2 to be linearly dependent to u_2 and w_3 to be linearly dependent to u_3 . Since W_1, W_2 is closed under scalar multiplication, where $\alpha \in \mathbb{R}^4$, $\alpha w_2 \in W_1$ and $u_2 \in W_2$. As w_2 and u_2 are linearly dependent, u_2 can be expressed as $u_2 = \alpha w_2$. Therefore $u_2 \in W_1$. By the same argument with u_3 and w_3 , $u_3 \in W_1$.

By the reverse, $w_2 = \frac{1}{\alpha} u_2 \in W_1$. Therefore, $w_2 \in W_1$ and $w_3 \in W_2$.

Since $w_2, w_3, u_2, u_3 \in W_1$ and $w_2, w_3, u_2, u_3 \in W_2$, $w_2, w_3, u_2, u_3 \in W_1 \cap W_2$.

A \mathbb{R}^4 plane through the origin is formed from two linearly independent vectors $\in \mathbb{R}^4$ as $\dim \text{plane} = 2$. To satisfy the condition these two vectors must be $\in W_1 \cap W_2$, either $\{w_2, w_3\}$ or $\{u_2, u_3\}$ can be picked.

Finally $\text{span}\{w_2, w_3\} \in W_1 \cap W_2$, and being two linearly independent vectors, must be a basis for a plane by the Algebra Triangle Theorem. Therefore, they must also span a plane by the definition of basis.