

Linear Algebra Foundations for Systems

Prepared by Staples Education

Condensed Cheat Sheet: Span, Linear Maps, Determinants, and Eigenvalues

Unit 14 at a Glance: Linear Algebra for Systems

Before coupled differential equations can be solved, the language of linear algebra has to be in place. A matrix is read geometrically: its columns say where the basis vectors land, its determinant says how it scales area and whether it flips orientation, and its eigenvectors mark the special directions it merely stretches. The payoff is the eigenvalue method — guessing $\mathbf{x} = e^{\lambda t}\mathbf{v}$ turns the system $\mathbf{x}' = A\mathbf{x}$ into the algebraic eigenvalue problem $A\mathbf{v} = \lambda\mathbf{v}$.

- **Vectors and transformations** — linear combinations, span, basis, and the matrix whose *columns are the images of the basis vectors*.
- **The determinant** — the signed area-scaling factor $ad - bc$; zero means collapse, negative means an orientation flip.
- **Change of basis** — the similarity transform $P^{-1}AP$ rewrites the same map in new coordinates; an eigenbasis makes it diagonal.
- **Eigentheory** — $A\mathbf{v} = \lambda\mathbf{v}$, the characteristic equation $\det(A - \lambda I) = 0$, and the trace/determinant shortcut for 2×2 .
- **Eigenvalues through ODEs** — $\mathbf{x} = e^{\lambda t}\mathbf{v}$ decouples $\mathbf{x}' = A\mathbf{x}$ into scalar modes, the gateway to Unit 15.

Part I — Condensed Cheat Sheet

14.1 Vectors, Span, and Linear Transformations

Linear combinations, span, and basis; linear transformations and the matrices that encode them; columns as images of the basis vectors; composition as matrix product.

Linear Combination, Span, and Basis

$$a\mathbf{v} + b\mathbf{w} \quad (a, b \text{ scalars}); \quad \text{span}\{\mathbf{v}_i\} = \left\{ \sum_i c_i \mathbf{v}_i \right\}; \quad \text{basis} \iff \text{independent and spanning.}$$

Linearity Condition

$$T(a\mathbf{u} + b\mathbf{v}) = aT(\mathbf{u}) + bT(\mathbf{v}) \quad \text{for all scalars } a, b \text{ and vectors } \mathbf{u}, \mathbf{v}.$$

The Matrix of a Transformation

$$A = [T(\mathbf{e}_1) \mid T(\mathbf{e}_2) \mid \cdots], \quad T(\mathbf{x}) = A\mathbf{x}, \quad (T_2 \circ T_1) \leftrightarrow A_2 A_1.$$

Method — Build the Matrix of a Linear Map

1. Apply T to each standard basis vector $\mathbf{e}_j$.
2. Record each image $T(\mathbf{e}_j)$ as the j -th *column*.
3. Assemble the columns into the matrix A .
4. Read off the action as $T(\mathbf{x}) = A\mathbf{x}$ and verify on a test vector.

Key Takeaway

A linear transformation is **completely determined by where it sends the basis**. The columns of A are exactly the images $T(\mathbf{e}_j)$, and composing two maps corresponds to *multiplying* their matrices (right-to-left).

Common Pitfall

A map with a nonzero shift, $T(\mathbf{u}) = \mathbf{u} + \mathbf{c}$ with $\mathbf{c} \neq \mathbf{0}$, is **affine, not linear** — it moves the origin. Linearity *requires* $T(\mathbf{0}) = \mathbf{0}$. Also, matrix multiplication is not commutative: $A_2A_1 \neq A_1A_2$ in general, so composition order matters.

14.2 The Determinant and Change of Basis

The determinant as a signed area-scaling factor; the formula $ad - bc$; what zero and negative determinants mean; change-of-basis matrices and the similarity transform $P^{-1}AP$.

The 2×2 Determinant

$$\det \begin{pmatrix} a & b \\ c & d \end{pmatrix} = ad - bc = \text{signed factor by which the map scales area.}$$

Reading the Determinant Geometrically

$|\det A|$ = area scaling, $\det A = 0 \Rightarrow$ collapse (dependent columns), $\det A < 0 \Rightarrow$ orientation reversed.

Change of Basis and Similarity

$B = P^{-1}AP$, columns of P = new basis vectors in the old coordinates; eigenbasis $\Rightarrow B$ diagonal.

Method — Re-express a Map in a New Basis

1. Place the new basis vectors (written in old coordinates) as the columns of P .
2. Compute the inverse P^{-1} .
3. Form the similarity transform $B = P^{-1}AP$.
4. If the columns of P are eigenvectors of A , then B is diagonal with the eigenvalues on the diagonal.

Key Takeaway

The determinant packages two facts in one number: its **magnitude is the area-scaling factor** and its **sign records orientation**. Because a change of basis only relabels coordinates, *similar* matrices $P^{-1}AP$ share the same determinant, trace, and eigenvalues.

Common Pitfall

Do not confuse a **negative** determinant with a **zero** determinant. Negative still means invertible — the plane is merely flipped. Only $\det A = 0$ collapses a dimension and makes A singular (columns linearly dependent).

14.3 Eigenvalues and Eigenvectors

The eigenvalue equation $A\mathbf{v} = \lambda\mathbf{v}$; the characteristic equation $\det(A - \lambda I) = 0$; trace as the sum and determinant as the product of eigenvalues; the trace/determinant shortcut for 2×2 matrices.

The Eigenvalue Equation

$A\mathbf{v} = \lambda\mathbf{v}, \quad \mathbf{v} \neq \mathbf{0} : \quad A$ only *stretches* $\mathbf{v}$ along its own line, by the factor λ .

The Characteristic Equation

$$\det(A - \lambda I) = 0; \quad 2 \times 2 : \quad \lambda^2 - (\operatorname{tr} A)\lambda + \det A = 0.$$

Trace–Determinant Shortcut

$$\lambda = m \pm \sqrt{m^2 - p}, \quad m = \frac{1}{2} \operatorname{tr} A, \quad p = \det A; \quad \operatorname{tr} A = \lambda_1 + \lambda_2, \quad \det A = \lambda_1 \lambda_2.$$

Method — Find Eigenvalues and Eigenvectors

1. Form $A - \lambda I$.
2. Set $\det(A - \lambda I) = 0$ and solve for λ (for 2×2 , use $\lambda^2 - (\operatorname{tr} A)\lambda + \det A = 0$).
3. For each λ , solve $(A - \lambda I)\mathbf{v} = \mathbf{0}$ for the eigenvector $\mathbf{v}$.
4. Verify $A\mathbf{v} = \lambda\mathbf{v}$.

Key Takeaway

Eigenvectors are the **special directions a matrix leaves invariant** — it only scales them, by the eigenvalue. In the eigenbasis the matrix becomes diagonal, which is exactly what makes powers, exponentials, and differential equations easy.

Common Pitfall

The characteristic equation is $\det(A - \lambda I) = 0$, **not** $\det A - \lambda = 0$, and the 2×2 form carries a *minus* sign on the trace: $\lambda^2 - (\text{tr } A)\lambda + \det A$. Handy check: for diagonal or triangular matrices the eigenvalues sit on the diagonal — the off-diagonal entries do not change them.

14.4 Eigenvalues Through Differential Equations

The trial solution $\mathbf{x} = e^{\lambda t} \mathbf{v}$ that reduces $\mathbf{x}' = A\mathbf{x}$ to the eigenvalue problem; one exponential mode per eigenpair; the general solution by superposition; growth, decay, and oscillation from the sign and type of the eigenvalues.

The Eigen-Ansatz

$$\text{Try } \mathbf{x} = e^{\lambda t} \mathbf{v} : \quad \mathbf{x}' = \lambda e^{\lambda t} \mathbf{v} = A e^{\lambda t} \mathbf{v} \Rightarrow A \mathbf{v} = \lambda \mathbf{v}.$$

General Solution (Distinct Eigenvalues)

$$\mathbf{x}(t) = c_1 e^{\lambda_1 t} \mathbf{v}_1 + c_2 e^{\lambda_2 t} \mathbf{v}_2 \quad (\text{superposition of the eigen-modes}).$$

Stability from the Eigenvalues

$$\lambda < 0 : \text{ decay}, \quad \lambda > 0 : \text{ growth}, \quad \lambda = \alpha \pm i\beta : \text{ oscillation (decays if } \alpha < 0).$$

Method — Solve $\mathbf{x}' = A\mathbf{x}$ by Eigenvalues

1. Find the eigenvalues of A from $\det(A - \lambda I) = 0$.
2. Find an eigenvector $\mathbf{v}_i$ for each eigenvalue λ_i .
3. Form the eigen-modes $e^{\lambda_i t} \mathbf{v}_i$.
4. Superpose: $\mathbf{x} = \sum_i c_i e^{\lambda_i t} \mathbf{v}_i$, then fix the c_i from the initial conditions.

Key Takeaway

The eigenvalue method **decouples a coupled linear system into independent scalar modes**. Along each eigenvector A acts as multiplication by λ , so $\mathbf{x}' = A\mathbf{x}$ collapses to $x' = \lambda x$ with solution $e^{\lambda t}$ — the foundation of every technique in Unit 15.

Common Pitfall

In the superposition each exponential must ride its **own** eigenvector: $c_1 e^{\lambda_1 t} \mathbf{v}_1 + c_2 e^{\lambda_2 t} \mathbf{v}_2$, never a crossed pairing. And long-term behavior is set by the sign of $\text{Re}(\lambda)$ — a single positive eigenvalue makes the whole system unstable.