

Methods and Results

Annual weather effects on Black-tern abundance and trends

We designed a model to estimate the effects of annual weather patterns on abundance of Black-terns on BBS surveys, while accounting for medium- and long-term population trends. We used two annual weather covariates: one that represented the local spring moisture using the 3-month (April - June) Standardized Precipitation Evapotranspiration Index (SPEI (Beguería et al., n.d.)); and, a second that represented the North Atlantic Oscillation Index (NAOI, (Hurrell 1995), accessed through <https://www.cpc.ncep.noaa.gov/products/precip/CWlink/pna/nao.shtml>) from the previous winter (January - June, on a 1-year lag, 18 - 12 months before the BBS surveys were conducted; following (Davis et al. 2023)).

The model was based on the GAMYE models described in (A. C. Smith and Edwards 2020) and (A. Smith et al. 2023). In the GAMYE model, the population trajectory (pattern of annual abundance through time) is modeled as an additive combination of a non-linear smooth (i.e., the GAM component) and year-effects that model the annual fluctuations around the smooth (the YE component). We added a sub-model to estimate the year-effects using two predictors, so that the annual fluctuations in stratum- i and year- t ($\gamma_{i,t}$) were an additive combination of the effect of SPEI ($\beta_{1_i} SPEI_{i,t}$), the effect of the NAOI in year $t-1$ ($\beta_{2_i} NAOI_{t-1}$), and an additional random variate ($\varepsilon_{i,t}$).

$$\gamma_{i,t} = \beta_{1_i} SPEI_{i,t} + \beta_{2_i} NAOI_{t-1} + \varepsilon_{i,t}$$

Each of the stratum-specific parameters for the effects of SPEI (β_{1_i}) and NAOI (β_{2_i}), were estimated as a combination of spatially-varying random effects (β'_{1_i}), centered on a mean hyperparameter (β'_1).

$$\beta_{1_i} = \beta'_1 + \beta''_{1_i}$$

We estimated the spatially-varying component (β''_{1_i}) using the same intrinsic spatial conditional autoregressive (iCAR) neighbourhood matrix used to estimate the spatially varying population trends ((A. Smith et al. 2023)). This iCAR modelling structure estimates the stratum-level component for each covariate as a normally distributed variate, centered on the mean of the

parameters in the surrounding strata, with an estimated among-strata variation. For example, the local component of the effect of SPEI on the annual relative abundance in stratum- i (β''_{1_i}) is a normal random variate centered on the mean of the effect in the set of n -strata that are direct neighbours of stratum- i ($n \in N_i$), and that has a standard deviation ($\sigma_{\beta''_1}$) estimated across the neighbourhood matrix for all strata.

$$\beta''_{1_i} \sim Normal \left(\frac{\sum_{n \in N_i} \beta''_{1_i}}{N_i}, \frac{\sigma_{\beta''_1}}{N_i} \right)$$

Exploring the spatially varying effects of annual weather

The overall mean effect of SPEI on annual abundance was positive ($\beta'_1 = 0.12$ [0.09-0.16]). The effects of SPEI varied across the species' range. It was strongest and clearly positive in the core of the species' range across the southern and central prairie pothole region. In the western and eastern periphery of the species' range the effect was only weakly positive or even slightly negative.




Figure 1: Spatially varying effect of spring moisture (June, 3-month SPEI) on the annual relative abundance of Black Tern observed during BBS surveys. The colours in the map represent the mean effect in each of the 1-degree longitude by 1-degree latitude strata, where blue colours represent positive effects of moisture on abundance, red represents negative effects.

The overall mean effect of NAOI on annual abundance was weakly negative but with relatively high uncertainty ($\beta'_2 = -0.04 [-0.09-0.02]$). NAOI had the strongest negative effect in the western Great Lakes region and in the Canadian portion of the Prairie Potholes region. By contrast, in the southern part of the species' range, NAOI had a largely positive effect on annual abundance.

References Cited

::: {#refs}

- Beguería, Santiago, Sergio M. Vicente Serrano, Fergus Reig-Gracia, and Borja Latorre Garcés. n.d. "SPEIbase v.2.9 [Dataset]."
- Davis, Kayla L., Sarah P. Saunders, Stephanie Beilke, Erin Rowan Ford, Jennifer Fuller, Ava Landgraf, and Elise F. Zipkin. 2023. "Breeding Season Management Is Unlikely to Improve Population Viability of a Data-Deficient Migratory Species in Decline." *Biological Conservation* 283 (July): 110104. <https://doi.org/10.1016/j.biocon.2023.110104>.
- Hurrell, James W. 1995. "Decadal Trends in the North Atlantic Oscillation: Regional Temperatures and Precipitation." *Science* 269 (5224): 676–79. <https://doi.org/10.1126/science.269.5224.676>.
- Smith, Adam C, and Brandon P M Edwards. 2020. "North American Breeding Bird Survey Status and Trend Estimates to Inform a Wide Range of Conservation Needs, Using a Flexible Bayesian Hierarchical Generalized Additive Model." *The Condor*, no. duaa065 (December). <https://doi.org/10.1093/ornithapp/duaa065>.
- Smith, Adam, Allison Binley, Lindsay Daly, Brandon Edwards, Danielle Ethier, Barbara Frei, David Iles, Timothy Meehan, Nicole Michel, and Paul Smith. 2023. "Spatially Explicit Bayesian Hierarchical Models for Avian Population Status and Trends." <https://doi.org/10.32942/X2088D>.




Figure 2: Maps representing the uncertainty in the effect of the SPEI. Left map represents the lower limit of the 90% Credible Interval and the right map represents the upper limit. Regions that are similar colours in both maps have local effects of SPEI where at least a 95% of the posterior probability is in the same direction (e.g., if regions are blue in both maps, at least 95% of the posterior probability of the effect is positive). In general, these maps show that the local effects of SPEI are clearly positive in the core of the species' range, but weaker and less certain in the western and eastern edges of the range.

Effect of Jan–June NAO (1–year lag) on annual abundance




Figure 3: Spatially varying effect of the North Atlantic Oscillation (January - June in year $t-1$) on the annual relative abundance of Black Tern observed during BBS surveys. The top map shows the mean effect in each of the 1-degree longitude by 1-degree latitude strata, where blue colours represent positive effects of NAOI on abundance, red represents negative effects, and white represents no effect.




Figure 4: Maps representing the uncertainty in the effect of the North Atlantic Oscillation. Left map represents the lower limit of the 90% Credible Interval and the right map represents the upper limit. Regions that are similar colours in both maps have local effects of NAOI where at least a 95% posterior probability is in the same direction (e.g., if regions are blue in both maps, at least 95% of the posterior probability of the effect is positive). In general, these maps show that the local effects of NAOI are somewhat uncertain